

ELISA Test on Cordia myxa L. Leaf Extract for α -Glucosidase Inhibitor

by Virsa Handayani

Submission date: 29-May-2023 09:00AM (UTC+0800)

Submission ID: 2104061515

File name: L_ELISA_Test_on_Cordia_myxa_L._Leaf_Extract_for_-Glucosidase.pdf (444K)

Word count: 2484

Character count: 13866

ELISA Test on *Cordia myxa* L. Leaf Extract for α -Glucosidase Inhibitor

Ahmad Najib^{1,*}, Aktsar Roskiana Ahmad¹, Virsa Handayani²

ABSTRACT

Aimed: Determine the potential of *Cordia myxa* L. leaf on inhibited α -glucosidase. **Material:** ELISA Kit, Ethanol 96%, Column Chromatography, n-hexane, ethyl acetate, Glucobay®. **Method:** Sample from *Cordia myxa* L. leaf extracted by ethanol 96% then evaporated to get the sticky extract. The sticky extract of *Cordia myxa* L. leaf fractionated by column chromatography with n-hexane, n-hexane: ethyl acetate (90:10; 80:20; 75:25; 70:30; 65:35; 60:40; 55:45; 50:50) **Assay:** The fractions assayed by ELISA (Enzyme-Linked Immunosorbent Assay) with acarbose (Glucobay®) as the comparator. **Result:** The results showed that the n-hexane fraction is the highest potency on inhibited α -glucosidase with the noncompetitive mechanism. The IC₅₀ of n-hexane fraction is 0.53 ppm been while the acarbose is 6.85 ppm. **Conclusion:** The n-hexane fraction of *Cordia myxa* L. leaf has the highest potency to use for possible decrease blood glucose level.

Key words: *Cordia myxa* L., ELISA, α -Glucosidase, IC₅₀, Acarbose.

Ahmad Najib^{1,*}, Aktsar Roskiana Ahmad¹, Virsa Handayani²

¹Phytochemistry Division-Pharmacognosy-Phytochemistry Laboratory, Faculty of Pharmacy Universitas Muslim Indonesia, Makassar- INDONESIA.

²Pharmacognosy Division-Pharmacognosy-Phytochemistry Laboratory, Faculty of Pharmacy Universitas Muslim Indonesia, Makassar- INDONESIA.

Correspondence

Mr. Ahmad Najib

Universitas Muslim Indonesia, Laboratory of Pharmacognosy-Phytochemistry, Faculty of Pharmacy Kampus II UMI, Makassar- INDONESIA.

Phone no : +62 81524045514

E-mail: ahmad.najib@umi.ac.id

History

- Submission Date: 31-10-2018;
- Review completed: 27-11-2018;
- Accepted Date: 30-11-2018

DOI : 10.5530/pj.2019.11.54

Article Available online

<http://www.phcogj.com/v11/i2>

Copyright

© 2019 Phcog.Net. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



INTRODUCTION

Cordia myxa L. belongs to family Boraginaceae, this plant family around 2740 species distributed in 148 genera. The genus *Cordia* is one of the most presenting of this family. *Cordia* is a genus of trees or shrubs the borage family.¹ About 300 species have been identified worldwide that use as folk medicine in Indonesia.² The phytochemical qualitative analysis showed that *Cordia myxa* L. contain the presence of oil, glycosides, flavonoids, sterols, saponins, terpenoids, alkaloids, phenolic acids, coumarins, tannins, resins, gums and mucilage.³ The methanolic extracts from this plants species have reported as the antioxidant and α -amylase and α -glucosidase enzyme inhibitory activity.⁴

Nondependent insulin Diabetes Mellitus (NIDM) or type 2 diabetes mellitus (DM) is correlated with the α -glucosidase inhibitors, this enzyme which is present in the brush border of enterocytes in the intestinal villi. Overall, the α -glucosidase inhibitors reduce postprandial insulin concentrations through the attenuated rise in postprandial glucose levels.⁵

The mechanism of α -glucosidase inhibitors base on inhibited the enzyme substrate react with an enzyme on room temperature.⁶ Data resources from the absorbance measurement by the spectrophotometer on specific maximum wave length.⁵ The resulting base on the IC₅₀ value of sample compares with the IC₅₀ of positive control.⁷ The on this case IC₅₀ value will show the best activity if equal or less than IC₅₀ value of positive control.⁸ On this method the researcher will need relatively more sample to investigate⁵ so to reduce the needs of the sample it will use by another method such

as ELISA.⁹ This research will use it to investigate the inhibitory of α -glucosidase by *Cordia myxa* L. leaf.

EXPERIMENTAL METHOD

Sample Preparation

Cordia myxa L. leaf from Enrekang regency South Celebes-Indonesia. The taxonomic identification by the Botanical division of Pharmacognosy-Phytochemistry Laboratory, Faculty of Pharmacy, Universitas Muslim Indonesia. The 2.15 Kg dried sample by room temperature was grind then extracted by 5 liters' ethanol 95%, this process was done four times. The liquid extract then evaporated to give the gummy extract.¹⁰

Sample Fractionation

The gummy extract fractionated by column chromatography with silica gel G. 60 (Merck®) with n-hexane, n-hexane: ethyl acetate (90:10; 80:20; 75:25; 70:30; 65:35; 60:40; 55:45; 50:50). Results of fractions identify by Thin Layer Chromatography (TLC)¹¹ to investigate the chromatogram profile.¹²

ELISA Test on Fractions

Enzyme Linked Immuno Sorbent Assay (ELISA)¹³ on fraction base on scale down from previous research⁵ using the reaction mixture consisting 25 μ L of 2 mM p-nitrophenyl α -D-glucopyranoside (Sigma Chemical Co.), 49.5 μ L phosphate buffer (pH 7.0) adding to flask contains 0.5 μ L of sample dissolved

Cite this article: Najib A, Ahmad AR, Handayani V. ELISA Test on *Cordia myxa* L. Leaf Extract for α -Glucosidase Inhibitor. Pharmacogn J. 2019;11(2):358-61.

in DMSO at various concentration (0.31-2.5 $\mu\text{g mL}^{-1}$). The reaction mixture was pre-incubated for 5 min at 37°C, the reaction was started by adding 25 μL α -glucosidase (Sigma) incubation was continued from 30 min. The reaction stopped by adding 1 ml of 0.01 M Na_2CO_3 . The activity of α -glucosidase was determined by measuring the release of p-nitro phenol at 400 nm. Acarbose (Glucobay®) positive control of α -glucosidase.⁵

Kinetics of Inhibition Against α -glucosidase

Inhibition of sample with α -glucosidase activities was measured by increasing concentration of p-nitrophenyl α -D-glucopyranoside as a substrate in the absence or presence of samples at different concentrations. Inhibition type was determined by Lineweaver-Burk plot analysis of the data, which were calculated from the result according to Michaelis-Menten kinetics.¹⁴

RESULT AND DISCUSSION

After four times maceration with ethanol 96%, the result of extraction shown Table 1.

This research used ethanol because it can extract more active compound than the others.¹⁵ Water and various concentrations (50%, 75% and 100%) This solvent has the lower toxicity and record experiments results were investigated to find that ethanol concentration, solvent/solid ratio and time to increase the extraction yield.¹⁶ The effect of different solvents (methanol, ethanol, acetone and distilled water) and the extraction yield increases with increasing polarity of the solvent used in extraction.¹⁵ Water and various concentrations (50%, 75% and 100%)

After evaporated and fractioned with n-hexane and n-hexane: ethyl acetate on various concentration then obtained to investigate the chromatogram¹⁷ as shown in Figure 1.

Table 1: The results of the calculations percent yield of ethanol extract of *Cordia myxa* L. leaf.

Solvent	The weight of the sample (Kg)	The weight of extract (g)	Yield (%)
Ethanol 96 %	2.15	100	4.65



Figure 1: Chromatogram on a fraction.

Description: Stationary phase = Silica gel F₂₅₄
Mobile phase = n-hexane: ethyl acetate (7:3)
Apparatus: Camag Nanomat 4
Detector: UV light 366 nm




Figure 2: UV-Vis Spectrum on Sample.

Table 2: The results of the ELISA assay on a fraction of *Cordia myxa* L. leaf.

Sample	Comparison of solven (n-hex : EtoAc)	Activity from IC ₅₀ value (ppm)
Ethanol extract	-	4.44
Acarbose (Glucobay®)	-	6.85
1 st Fraction	n-hex	0.53
2 nd Fraction	90:10	5.81
3 rd Fraction	80:20	0.89
4 th Fraction	75:25	2.43
5 th Fraction	70:30	0.91
6 th Fraction	65:35	7.55
7 th Fraction	60:40	9.28
8 th Fraction	55:45	12.60
9 th Fraction	50:50	16.52

Chromatogram showed that there are 10 fraction obtained some spot representing of many compound.¹⁸ To clarify the amount of compound on the sample we use UV-Vis spectrophotometer¹⁹ as shown in Figure 2.

Analysis of UV-Vis spectrum on sample identified 4 peaks. It important to know that sensitivity of this equipment can detect the compound which is cannot detect by TLC.²⁰ The data above will assist to investigated for furthermore for the compounds from the sample by elucidation method to determinate the potential compound that can inhibit the α -glucosidase.²¹ ELISA assay on the sample as showed on Table 2.

The results on Table 2 regarding the potential for inhibitory of α -glycosidase have been proofed on sample²² even the IC₅₀ there are more potent than acarbose. Since the 6th fraction showed that the activity has been decreasing; it mean the active compound are reduced from the fractions before.

The mechanism of kinetic reaction base on Lineweaver-Burk plot analysis from the sample has been calculated and plotted on Figure 3, below

Regarding on graphic above the inhibitory base on noncompetitive mechanism because of intercept on each line not on Y axis.²³ This mechanism makes a chance to investigate on *in silico* method on our next research as we have been determining on another sample.²⁴




Figure 3: Lineweaver-Burke plot of the reaction α -glucosidase in the presence of sample.

CONCLUSION

The n-hexane fraction of *Cordia myxa* L. leaf have a potential activity to inhibit α -glucosidase related by the highest potency to use for decrease blood glucose level blood with IC_{50} value 0.53 ppm with kinetic mechanism of inhibitory from sample base on noncompetitive.

ACKNOWLEDGEMENT

Thank you to Directorate Research, Technology and High Education, Ministry of Research, Technology and High Education Republic Indonesia for funding this research for grant Basic Research for University Institution (PDUPT-RISTEK DIKTI) 2018.

CONFLICT OF INTEREST

The authors declare that their is no conflict of interest.


REFERENCES

- Abdel-aleem ER, Seddik FEF, Samy MN. Botanical studies of the leaf of *Cordia myxa* L. J of Pharmacognosy and Phytochemistry. 2017;6(6):2086-91.
- Malik A, Ahmad AR. Antidiabetic Effect of Standardized Extract of Indonesian Kanunang Leaves (*Cordia myxa* L.). Int J Pharmtech Res. 2016;9(8):268-75.
- Al-snafi PAE. The Pharmacological and therapeutic importance of *Cordia myxa*-A review. IOSR J of Pharmacy. 2017.
- Marini G, Graikou K, Zengin G, Karikas GA, Gupta MP. Phytochemical Analysis and Biological Evaluation of Three Selected *Cordia* species from Panama. Ind Crop Prod. 2018;120:84-9.
- Najib A, Hartati S, Elya B. *In vitro* bioassay of n-buthanol isolate of *Acorus calamus* L. on inhibitory of activity α -glucosidase. Int J PharmTech Res. 2011;3.
- Nurcholis W, Munshif AA, Ambarsari L. Xanthorrhizol contents, α -glucosidase

inhibition and cytotoxic activities in ethyl acetate fraction of *Curcuma zanthorrhiza* accessions from indonesia. Brazilian J Pharmacogn. 2018;28(1):44-9.

- Wu H, Zeng W, Chen L, Yu B, Guo Y, Chen G, et al. Integrated multi-spectroscopic and molecular docking techniques to probe the interaction mechanism between maltase and 1-deoxyxojirimycin, an α -glucosidase inhibitor. Int J Biol Macromol. 2018;114:1194-202.
- Najib A, Risdia, Waris, Esti, Dian PN. Radical Scavenging Activity of Leaf Extract of Edible Hibiscus. Int J Pharmacol. 2016;9(6):343-7.
- Wielkoszynski T, Moghaddam A, Bäckman A, Broden J, Piotrowski R, Mond-Paszek R, et al. Novel diagnostic ELISA test for discrimination between infections with *Yersinia enterocolitica* and *Yersinia pseudotuberculosis*. Eur J Clin Microbiol Infect Dis. 2018;37(12):2301-6.
- Najib A, Alam G, Halidin M. Isolation and identification of antibacterial compound from diethyl ether extract of *Plantago major* L. Pharmacogn J. 2013;4(31):59-62.
- Hamidu L, Ahmad AR, Najib A. Qualitative and quantitative test of total flavonoid buni fruit (*Antidesma bunius* (L.) Spreng) with UV-Vis spectrophotometry method. Pharmacogn J. 2018;10(1).
- Malik A, Kurniawan A, Najib A. Comparative study of HPTLC fingerprint of β -Asarone content between leaves and rhizome of *Acorus calamus* L. Int J PharmTech Res. 2014;6(2):829-33.
- Sahli H, et al. An advanced intelligent ELISA test for bovine tuberculosis diagnosis. Biomed Signal Process Control. 2018;46:59-66. doi:10.1016/j.bspc.2018.05.031
- Hartati S, Elya B, Najib A. n-Buthanol fraction of *Acorus calamus* rhizome extract to inhibit the activity of β -glucosidase. Journal of Tropical Medicinal Plants. 2010;11(2):201-3.
- Do QD, et al. Effect of extraction solvent on total phenol content, total flavonoid content and antioxidant activity of *Limnophila aromatica*. J Food Drug Anal. 2014;22(3):296-302. doi:10.1016/j.jfda.2013.11.001
- Ballesteros LF, Teixeira JA, Mussatto SI. Selection of the Solvent and Extraction Conditions for Maximum Recovery of Antioxidant Phenolic Compounds from Coffee Silverskin. Food Bioprocess Technol. 2014;7(5):1322-32. doi:10.1007/s11947-013-1115-7
- Mohammad A, et al. Detection reagents used in on-plate identification of amino acids by thin layer chromatography: A review. Journal of Liquid Chromatography and Related Technologies. 2018;1-9. doi:10.1080/10826076.2018.1485035
- Jung JW, et al. Three new isoprenylated flavonoids from the root bark of *Morus alba*. Molecules. 2016;21(9):2-11.
- D'Archivio AA, Maggi MA. Geographical identification of saffron (*Crocus sativus* L.) by linear discriminant analysis applied to the UV-visible spectra of aqueous extracts. Food Chem. 2017;219:408-13. doi:10.1016/j.foodchem.2016.09.169
- Schwack W, Pellissier E, Morlock G. Analysis of unauthorized Sudan dyes in food by high-performance thin-layer chromatography. Anal Bioanal Chem. 2018;1. doi:10.1007/s00216-018-0945-6
- Zhao Y, Kongstad KT, Jäger AK, Nielsen J, Staerk D. Quadruple high-resolution α -glucosidase/ α -amylase/PTP1B/radical scavenging profiling combined with high-performance liquid chromatography-high-resolution mass spectrometry-solid-phase extraction-nuclear magnetic resonance spectroscopy for identification of antidiabetic constituents in crude root bark of *Morus alba* L. J Chromatogr A. 2018. doi:10.1016/j.chroma.2018.04.041
- Parandiyal AK, Kaushal R, Chaturvedi OP. Forest and Fruit Trees-based Agroforestry Systems for Productive Utilization of Ravine Lands. In Ravine Lands: Greening for Livelihood and Environmental Security. Springer, Singapore. 2018.
- Gong Y, et al. Inhibitory effect of hesperetin on α -glucosidase: Molecular dynamics simulation integrating inhibition kinetics. Int J Biol Macromol. 2017. doi:10.1016/j.ijbiomac.2017.03.072
- Dewi Y, Mursalin, Najib A. *In silico* screening of chemical compounds from Sweet flag (*Aracus calamus* L.) as α -Glucosidase inhibitor. Int Res J Pharm. 2013;4(3):110-2.

GRAPHICAL ABSTRACT



SUMMARY

- Since *Cordia myxa* L. use to cure disease related to diabetes mellitus as a folk medicine this research finding the data to support the empirical base. This research focused on the mechanism of α -glucosidase inhibition one of three mechanisms that can cure the diabetic patient. By Enzyme-Linked Immunosorbent Assay (ELISA) this research has conducted. There are many fractions have been assayed thus resulted that n-hexane fraction has the best inhibition with IC_{50} 0.53 ppm.

ABOUT AUTHORS



Ahmad Najib: Phytochemistry Division, Pharmacognosy-Phytochemistry Laboratory Faculty of Pharmacy Universitas Muslim Indonesia, Makassar-Indonesia.



Aktsar Roskiana Ahmad: Phytochemistry Division: Pharmacognosy-Phytochemistry Laboratory Faculty of Pharmacy Universitas Muslim Indonesia, Makassar-Indonesia



Virsa Handayani: Pharmacognosy Division Pharmacognosy-Phytochemistry Laboratory Faculty of Pharmacy Universitas Muslim Indonesia, Makassar-Indonesia.

Cite this article: Najib A, Ahmad AR, Handayani V. ELISA Test on *Cordia myxa* L. Leaf Extract for α -Glucosidase Inhibitor. Pharmacog J. 2019;11(2):358-61.

ELISA Test on Cordia myxa L. Leaf Extract for α -Glucosidase Inhibitor

ORIGINALITY REPORT

16%
SIMILARITY INDEX

8%
INTERNET SOURCES

11%
PUBLICATIONS

9%
STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

1%
★ Maria G. Sande, Débora Ferreira, Joana L. Rodrigues, Luís D. R. Melo et al. "Electrochemical Aptasensor for the Detection of the Key Virulence Factor YadA of Yersinia enterocolitica", Biosensors, 2022
Publication

Exclude quotes On
Exclude bibliography On

Exclude matches Off