

**MELAKUKAN SUBMITTER AWAL KE
JURNAL BIOFLUX**

Aquaculture Management of Blue Swimming Crab using Integrated Submerged Net Cage in Pangkep Regency Waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Adimu HE

¹Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi;

²Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi;

³Graduates in Coastal and Ocean Resource Management, IPB University, West Java, Bogor 16680, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the suitability cultivation area and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analysed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen (5.1-10.7 mg/l), salinity (7-35 ‰), temperature (32-34.4 °C), and pH (7-8) respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which located close to the shoreline and cluster 2 (172,953 ha) which located far from the coastal line. The management of crab cultivation integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm of width. The net used made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied made from nylon Polyethylene (PE) number 12 with mesh size 1.0 cm. All parts of the submerged net-cage materials built from same materials with large size of net-cages.

Key Word: Aquaculture, management of blue swimming crab, using integrated, submerged.

Introduction. Blue swimming crab is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. The blue swimming crab (*Portunus pelagicus*) is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of blue swimming crab also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve blue swimming crab as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, have been the major issue of the blue swimming crab stock depletion and land utilization conflicts. Several studies related to the excessive catching of blue swimming crab in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2016; Wagiyo et al 2019).

In order to enhance severely depleted stocks of blue swimming crab, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. To achieve industrial scale production of soft swimming crabs, the option of using closed systems has increased in recent years. This type of system offers several

advantages, such as greater control over environmental variables, greater ease of installation, higher storage densities, greater ease of monitoring ecdysis occurrence and, especially, the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of 1). Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency, 2). Assessing the use of submerged net cage model for crab cultivation.

Materials and Methods

Description of the study sites. This study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia

Data collection. Materials and equipment used during the period of study including blue swimming larvae, submerged net cage, current meter, hand refractometer, Dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and

chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Table 1, Table 2, and Table 3).

Tabel 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	unsuitable (N)
X1 – X2 (= X1 + Ci)	acceptable (S2)
X2 – X3 (= max value)	suitable (S1)

Description: X0 = lower limit of assessment scale;
X1 = limit between unsuitable and acceptable class;
X2 = limit between acceptable and suitable class;
X3 = upper limit of assessment scale;

Tabel 2
Suitability matrix for adult blue swimming crab

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ -36‰	2	0.60	< 31or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or>6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00 – 1.33	not suitable (N)
1.34 – 1.67	fairly suitable (S2)
1.68 – 2.00	suitable (S1)

Data analysis. Data were descriptively and quantitatively analysed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency are presented in the following (table 4).

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Kriteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. This location is also very shallow in depth. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5 - 2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest regarding the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be done. The map of integration of crab cultivation using submerged net cage with other activities is presented below:




Figure 3. Map of management of crab cultivation in submerged net karamba integrated in the waters of Pangkep Regency.

Water quality measurement. For the determination of clusters 1 and cluster 2 of crab cultivation was done using the aspects of physics and chemistry of oceanography. The results of water temperature measurements at 28 stations during survey period; temperature 27 - 34 °C; salinity 28 - 34 ppt; pH (6.50 - 8.00). The optimal water pH for fish and shrimp life is 6.5 - 8 (neutral). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 is detrimental effect on aquatic animal caused high mortality.

DO measurements in the concentration range of 5.17 - 7.97 mg/L. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg/L which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32 °C (Adiwijaya et al 2002). However, the crabs are very tolerant of water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village; salinity 30-43 ppt, DO (4-10 mg/L), pH 4-8 and temperatures 27-34 °C.

Submerged net cage model for blue swimming crab farming. A rectangular model of submerged net cage was used: 1) submerged net cage size is 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size is 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapped using net and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of blue swimming crab can be done in the earthen pond with one log pond and two grow-out ponds [10]. The crablets reared for 1-2 months until the crabs can be distinguished by sex. Furthermore, the grow-out of crab is carried out by

applying a floating cage system or a net installed near the beach. The rearing ponds size is 50x100 meters. Ponds of this size can accommodate up to 10,000 of crab. The submerged net cage is shown in the following figure:



Figure 4. Submerged Net Cage Model for blue swimming crab farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers

To simplify the submerged and lift process, a plastic PE no 7 mm is used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 meters. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope is approximately 10 meters. So that, during tide, the submerged net cage is visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.


Acknowledgement. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education who has funded this research through Applied Research for Product and researcher colleagues who have provided significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. Reviews in Fisheries Science 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. Majalah Ilmu Kelautan. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. Jurnal Penelitian Perikanan Indonesia. 25 (2): 79-92. [In Indonesia]
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. Tropical Life Sciences Research 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]

*** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

HASIL USULAN PERTAMA DIMINTA PENGELOLAN JURNAL MEMBERBAIKI
NASKAH, UNTUK MENDAPATKAN PERSETUJUAN NASKAH DITERIMA
SEKALIGUS JUGA MELAKUKAN PEMBAYARAN



Aquaculture management of blue swimming crab *Portunus Pelagicus* (Linnaeus, 1758) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Department of Fisheries Resources Utilization, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Department of Marine Science, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal husbandry, Sembilanbelas November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages.

Key Words: Cultivation, Cluster, farming, fishermen, production,

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, have been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

Comment [L1]: Based on <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A2]: Please also display Latin name <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [L3]:

Comment [L4]:

Comment [A5]: Please display full name

Comment [L6]: has been chaged

Comment [A7]: The Abstract should also contain the conclusions of the study.

Comment [L8]: Conclusion

Comment [A9]: According to the international standards in scientific writing it is desirable to display 5 key words which do not appears in the title. This will increase findings via key words and implicit citations.

Comment [L10]: Has been changed

Comment [A11]: At the first mention please also display full Latin name according to WoRMS: <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A12]: After the first mention only the abbreviated Latin name should be displayed.

Comment [L13]: has been changed to *P. pelagicus*

Comment [A14]: Sara et al 2029?

Comment [L15]: has been changed to Sara et al 2019

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of *P. pelagicus* using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are ease of control, efficiency and fast monitoring, and the possibility of combining several forms of management systems, (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [L16]: has been changed

Comment [A17]: Plagiarism detected. Please reformulate.

Comment [L18]: has been repaired

Comment [A19]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *P. pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Table 4
The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates	Criteria
-------------	----------

Comment [L20]: has been repaired

Comment [A21]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	Cluster 1
119° 28' 49.413" E	4° 52' 25.889" S	
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	Cluster 2
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Comment [L22]: Has been repaired

Comment [A23]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

Comment [L24]: Karamba is floating net cages

Comment [A25]: What "karamba" means? Please explain.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism is evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *P. pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon)

Comment [L26]: OK

Comment [A27]: Please sustain with reference Citation needed.

Comment [L28]: [10] this figure is not included in the writing

Comment [A29]: What represents [10]? Please explain.

Comment [A30]: sexual dimorphism is evident

Comment [L31]: has been changed

PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. Reviews in Fisheries Science 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. Majalah Ilmu Kelautan. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. Jurnal

- Penelitian Perikanan Indonesia. 25 (2): 79-92. [In Indonesia]
- Wardhany V.A., Yuliandoko H., Subono., Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. International Electronics Symposium on Engineering Technology and Applications (IES-ETA), Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. Tropical Life Sciences Research 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:


Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

Comment [A32]: Please display all authors with their full names one by one with their affiliation including: University, Faculty, Department, County, Postal Code, City, Street Name, Street No., e-mail address. Thank you!

HASIL KOREKSI KEDUA PENULIS MEMBERPAKI NASKAH SESUAI CATATAN
YANG ADA MELAMPIRKAN BUKTI TELAH MELAKUKAN PEMBAYARAN



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Graduates in Coastal and Ocean Resource Management, IPB University, West Java, Bogor 16680, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [A1]: Please note: Images can contain only English legends/description. Please translate **Kec.**

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying

Comment [A3]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora.,

- Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. *Seminar Nasional Pendidikan, Sains dan Teknologi* ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id


Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

HASIL KOREKSI KETIGA PENULIS MEMBERPAKI NASKAH SESUAI CATATAN
YANG DIKIRIM MELALUI EMAIL JURNAL BIOFLUX



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal Husbandry, Nineteen November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Comment [L1]: Affiliation was changed because he worked at this institution. thank you

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Comment [L3]: Ok

chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) - X1 (= X0 + Ci)	Unsuitable (N)
X1 - X2 (= X1 + Ci)	Acceptable (S2)
X2 - X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00-1.33	Not suitable (N)
1.34-1.67	Fairly suitable (S2)
1.68-2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of aquaculture region and marine space management was based on the efficiency and the

effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [L4]: Ok

Comment [A5]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

Comment [A6]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

Comment [L7]: Ok

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water

- commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young, G., Eric G., Johnson, Jana L. D., Davis, Anson H., Hines, Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id


Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Campus B Central Buton, Mawasangka, e-mail: hasan.adimu@usn.ac.id

HASIL KOREKSI KE EMPAT PENULIS MEMBERPAKI NASKAH SESUAI CATATAN
YANG DIKIRIM MELALUI EMAIL JURNAL BIOFLUX



Aquaculture management of blue swimming crab using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Adimu

HE

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Graduates in Coastal and Ocean Resource Management, IPB University, West Java, Bogor 16680, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages.

Key Words: Aquaculture, management of blue swimming crab, using integrated, submerged.

Introduction. Blue swimming crab is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. The blue swimming crab (*Portunus pelagicus*) is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of blue swimming crab also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve blue swimming crab as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, have been the major issue of the blue swimming crab stock depletion and land utilization conflicts. Several studies related to the excessive catching of blue swimming crab in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2016; Wagiyono et al 2019).

In order to enhance severely depleted stocks of blue swimming crab, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be done using a composite culture of crabs with other cultured commodities such as

Comment [A1]: Please also display Latin name
<http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A2]: Please display full name

Comment [A3]: The Abstract should also contain the conclusions of the study.

Comment [A4]: According to the international standards in scientific writing it is desirable to display 5 key words which do not appears in the title. This will increase findings via key words and implicit citations.

Comment [A5]: At the first mention please also display full Latin name according to WoRMS:
<http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A6]: After the first mention only the abbreviated Latin name should be displayed.

Comment [A7]: Sara et al 2029?

finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. To achieve industrial scale production of soft swimming crabs, the option of using closed systems has increased in recent years. This type of system offers several advantages, such as greater control over environmental variables, higher ease of installation, higher storage densities, greater ease of monitoring ecdysis occurrence and, especially, the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included blue swimming larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study.

Comment [A8]: Plagiarism detected. Please reformulate.

Comment [A9]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult **blue swimming crab**

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of **blue swimming crab farming.** The management of **blue swimming crab** farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of aquaculture region and marine space management was based on the efficiency and the effectiveness of **blue swimming crab** using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab

cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Table 4
The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	

Comment [A10]: Please redo figure. Writing is small and not intelligible.
Please note: Images can contain only English legends/description.

Coordinates		Criteria
X	Y	
119° 27' 25.498" E	4° 45' 22.629" S	Cluster 2
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged net karamba integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and

Comment [A11]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

Comment [A12]: What "karamba" means? Please explain.

Comment [A13]: Please sustain with reference. Citation needed.

lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for blue swimming crab farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of blue swimming crab can be done in the earthen pond with one log pond and two grow-out ponds [10]. The crablets were reared for 1-2 months until the crabs can be distinguished by sex. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.

Comment [A14]: What represents [10]? Please explain.

Comment [A15]: sexual dimorphism is evident



Figure 4. Submerged net cage model for blue swimming crab farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length

of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References


- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. Reviews in Fisheries Science 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. Majalah Ilmu Kelautan. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. Jurnal Penelitian Perikanan Indonesia. 25 (2): 79-92. [In Indonesia]
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. Tropical Life Sciences Research 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]

- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Comment [A16]: Please display all authors with their full names one by one with their affiliation including: University, Faculty, Department, Country, Postal Code, City, Street Name, Street No., e-mail address. Thank you!

HASIL KOREKSI KE ENAM PENULIS MEMBERPAKI NASKAH SESUAI CATATAN
YANG DIKIRIM MELALUI EMAIL JURNAL BIOFLUX



Aquaculture management of blue swimming crab *Portunus Pelagicus* (Linnaeus, 1758) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Department of Fisheries Resources Utilization, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Department of Marine Science, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal husbandry, Sembilanbelas November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages.

Key Words: Cultivation, Cluster, farming, fishermen, production,

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, have been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

Comment [L1]: Based on <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A2]: Please also display Latin name <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [L3]:

Comment [L4]:

Comment [A5]: Please display full name

Comment [L6]: has been chaged

Comment [A7]: The Abstract should also contain the conclusions of the study.

Comment [L8]: Conclusion

Comment [A9]: According to the international standards in scientific writing it is desirable to display 5 key words which do not appears in the title. This will increase findings via key words and implicit citations.

Comment [L10]: Has been changed

Comment [A11]: At the first mention please also display full Latin name according to WoRMS: <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A12]: After the first mention only the abbreviated Latin name should be displayed.

Comment [L13]: has been changed to *P. pelagicus*

Comment [A14]: Sara et al 2029?

Comment [L15]: has been changed to Sara et al 2019

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of *P. pelagicus* using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are ease of control, efficiency and fast monitoring, and the possibility of combining several forms of management systems, (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [L16]: has been changed

Comment [A17]: Plagiarism detected. Please reformulate.

Comment [L18]: has been repaired

Comment [A19]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *P. pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Table 4
The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates	Criteria
-------------	----------

Comment [L20]: has been repaired

Comment [A21]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	Cluster 1
119° 28' 49.413" E	4° 52' 25.889" S	
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	Cluster 2
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Comment [L22]: Has been repaired

Comment [A23]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

Comment [L24]: Karamba is floating net cages

Comment [A25]: What "karamba" means? Please explain.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. [The optimal water pH for fish and shrimp life is 6.5-8 (neutral)] (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism is evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *P. pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon)

Comment [L26]: OK

Comment [A27]: Please sustain with reference Citation needed.

Comment [L28]: [10] this figure is not included in the writing

Comment [A29]: What represents [10]? Please explain.

Comment [A30]: sexual dimorphism is evident

Comment [L31]: has been changed

PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. Reviews in Fisheries Science 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. Majalah Ilmu Kelautan. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. Jurnal

- Penelitian Perikanan Indonesia. 25 (2): 79-92. [In Indonesia]
- Wardhany V.A., Yuliandoko H., Subono., Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. International Electronics Symposium on Engineering Technology and Applications (IES-ETA), Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. Tropical Life Sciences Research 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:


Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

Comment [A32]: Please display all authors with their full names one by one with their affiliation including: University, Faculty, Department, County, Postal Code, City, Street Name, Street No., e-mail address. Thank you!

HASIL KOREKSI KE 7 PENULIS MEMBERPAKI NASKAH SESUAI CATATAN YANG
DIKIRIM MELALUI EMAIL JURNAL BIOFLUX



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal Husbandry, Nineteen November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Comment [L1]: Affiliation was changed because he worked at this institution. thank you

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Comment [L3]: Ok

chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) - X1 (= X0 + Ci)	Unsuitable (N)
X1 - X2 (= X1 + Ci)	Acceptable (S2)
X2 - X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00-1.33	Not suitable (N)
1.34-1.67	Fairly suitable (S2)
1.68-2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of aquaculture region and marine space management was based on the efficiency and the

effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [L4]: Ok

Comment [A5]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

Comment [A6]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

Comment [L7]: Ok

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water

- commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young, G., Eric G., Johnson, Jana L. D., Davis, Anson H., Hines, Oded Zmora, Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id


Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Campus B Central Buton, Mawasangka, e-mail: hasan.adimu@usn.ac.id

HASIL KOREKSI KE 8 PENULIS MEMPERBAIKI NASKAH SESUAI CATATAN YANG
DIKIRIM MELALUI EMAIL JURNAL BIOFLUX



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Graduates in Coastal and Ocean Resource Management, IPB University, West Java, Bogor 16680, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [A1]: Please note: Images can contain only English legends/description. Please translate **Kec.**

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying

Comment [A3]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora.,

- Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id


Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

HASIL KOREKSI KE 8 PENULIS MEMPERBAIKI NASKAH SESUAI CATATAN YANG
DIKIRIM MELALUI EMAIL JURNAL BIOFLUX DENGAN
PEMERIKSAAN DETAIL



Aquaculture management of blue swimming crab *Portunus Pelagicus* (Linnaeus, 1758) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Department of Fisheries Resources Utilization, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Department of Marine Science, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal husbandry, Sembilanbelas November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages.

Key Words: Cultivation, Cluster, farming, fishermen, production,

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, have been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

Comment [L1]: Based on <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A2]: Please also display Latin name <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [L3]:

Comment [L4]:

Comment [A5]: Please display full name

Comment [L6]: has been chaged

Comment [A7]: The Abstract should also contain the conclusions of the study.

Comment [L8]: Conclusion

Comment [A9]: According to the international standards in scientific writing it is desirable to display 5 key words which do not appears in the title. This will increase findings via key words and implicit citations.

Comment [L10]: Has been changed

Comment [A11]: At the first mention please also display full Latin name according to WoRMS: <http://www.marinespecies.org/aphia.php?p=taxdet&id=1061743> cording to WoRMS

Comment [A12]: After the first mention only the abbreviated Latin name should be displayed.

Comment [L13]: has been changed to *P. pelagicus*

Comment [A14]: Sara et al 2029?

Comment [L15]: has been changed to Sara et al 2019

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of *P. pelagicus* using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are ease of control, efficiency and fast monitoring, and the possibility of combining several forms of management systems, (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [L16]: has been changed

Comment [A17]: Plagiarism detected. Please reformulate.

Comment [L18]: has been repaired

Comment [A19]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *P. pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Table 4
The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates	Criteria
-------------	----------

Comment [L20]: has been repaired

Comment [A21]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Comment [L22]: Has been repaired

Comment [A23]: Please redo figure. Writing is small and not intelligible. Please note: Images can contain only English legends/description.

Comment [L24]: Karamba is floating net cages

Comment [A25]: What "karamba" means? Please explain.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. [The optimal water pH for fish and shrimp life is 6.5-8 (neutral)] (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism is evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *P. pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon)

Comment [L26]: OK

Comment [A27]: Please sustain with reference Citation needed.

Comment [L28]: [10] this figure is not included in the writing

Comment [A29]: What represents [10]? Please explain.

Comment [A30]: sexual dimorphism is evident

Comment [L31]: has been changed

PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young., Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora., Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. Reviews in Fisheries Science 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. Majalah Ilmu Kelautan. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. Jurnal

- Penelitian Perikanan Indonesia. 25 (2): 79-92. [In Indonesia]
- Wardhany V.A., Yuliandoko H., Subono., Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. International Electronics Symposium on Engineering Technology and Applications (IES-ETA), Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. Tropical Life Sciences Research 29 (1): 1–15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:


Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

Comment [A32]: Please display all authors with their full names one by one with their affiliation including: University, Faculty, Department, County, Postal Code, City, Street Name, Street No., e-mail address. Thank you!

HASIL KOREKSI KE 10
PENULIS MEMPERBAIKI NASKAH SESUAI CATATAN YANG
DIKIRIM MELALUI EMAIL JURNAL BIOFLUX
DENGAN PROOFREADING



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Graduates in Coastal and Ocean Resource Management, IPB University, West Java, Bogor 16680, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and

Comment [A1]: Please note: Images can contain only English legends/description. Please translate **Kec.**

secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68–2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of

aquaculture region and marine space management wss based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4. The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying

Comment [A3]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
Alicia C., Young, Eric G., Johnson., Jana L. D., Davis., Anson H., Hines., Oded Zmora.,

- Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. *Seminar Nasional Pendidikan, Sains dan Teknologi* ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id


Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail:eldin.kahedupa@gmail.com

HASIL KOREKSI KE 11
PENULIS MEMPERBAIKI NASKAH SESUAI CATATAN YANG
DIKIRIM MELALUI EMAIL JURNAL BIOFLUX
DENGAN PROOFREADING



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan Eldin Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal Husbandry, Nineteen November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Comment [L1]: Affiliation was changed because he worked at this institution. thank you

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their customer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Alicia et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and

Comment [A2]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Comment [L3]: Ok

chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1
Criteria for determining the suitability of an area

X0 (= nilai value) - X1 (= X0 + Ci)	Unsuitable (N)
X1 - X2 (= X1 + Ci)	Acceptable (S2)
X2 - X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2
Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28°C - 31°C	2	0.80	<28°C or >31°C	1	0.40
Salinity	0.30	31‰ - 36‰	2	0.60	< 31 or 36‰	1	0.30
Dissolved Oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6,78 - 8,0	2	0.20	<6,78 or >8,0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3
Determination of area suitability category

Interval class	Category
1.00-1.33	Not suitable (N)
1.34-1.67	Fairly suitable (S2)
1.68-2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of aquaculture region and marine space management was based on the efficiency and the

effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

Comment [L4]: Ok

Comment [A5]: Please note: Images can contain only English legends/description. Please translate **Kec.**

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not Suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

Comment [A6]:

Please note: Images can contain only English legends/description.

Please translate **SELAT MAKASSAR**

Kab.

Kec.

Comment [L7]: Ok

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

Adiwijaya D., Jaya S., Sugeng., Sutikno E., 2002 Business opportunity for brackish water

- commodity: blue swimming crab (*Portunus pelagicus*) cultured in Earthen Pond. BBPBAP, Jepara 13-20. [In Indonesia]
- Alicia C., Young, G., Eric G., Johnson, Jana L. D., Davis, Anson H., Hines, Oded Zmora, Yonathan Z., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter. *Reviews in Fisheries Science* 16:254-261.
- Hamid A., Wardiatno Y., Lumbanbatu D.T.F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. *Asian Fisheries Science* 29: 206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. *AACL Bioflux* 8(5):729-739.
- Ihsan., 2015 [Sustainable utilisation of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University. [In Indonesia]
- Ihsan., Asbar., Asmidar., 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017. Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar. [In Indonesia]
- Juwana S., 1993 Effect of light intensity salinity and Temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds. *Majalah Ilmu Kelautan*. 16:194-204. [In Indonesia]
- Perkins F.J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York. PP 670 - 678.
- Sara L., Astuti O., Muzuni., Safilu., 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. *AACL Bioflux* 12 (5):1909-1917.
- Tavares C.P.S., Silva U.A.T., Pereira L.A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. *Reviews in Aquaculture* 0:1-11
- Wagiyo K., Tirtadanu., Ernawati T., 2019 Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay. *Jurnal Penelitian Perikanan Indonesia*. 25(2):79-92. [In Indonesia].
- Wardhany V.A., Yuliandoko H., Subono, Harun M.U., A.R., Astawa I.G.P., 2018 Temperature Control System Based on Fuzzy Logic, pH and Water Salinity in Vanammei Shrimp Farms. *International Electronics Symposium on Engineering Technology and Applications (IES-ETA)*, Bali. PP 145-149. [In Indonesia].
- Wiyono E.S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi, Indonesia. *Tropical Life Sciences Research* 29 (1): 1-15.
- Yuniati., Rachman F., 2016 Potential Cluster for Fisheries in Public Waters, East Java 2016. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN:978-602-61599-6-0.[In Indonesia]
- *** ME (Ministry of Environment), 2004 Minister Decree number 51 2004 about Water Quality Standard for Sea water. [In Indonesia]
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. [In Indonesia]

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail:ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmaleh@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id

Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Campus B Central Buton, Mawasangka, e-mail: hasan.adimu@usn.ac.id

SURAT PERNYATAAN
SELURUH PENULIS MEMBUAT SURAT PERNYATAAN
SEBELUM TERBIT JURNAL



Submission letter

Article title: **Aquaculture Management of Blue Swimming Crab using Integrated Submerged Net Cage in Pangkep Regency Waters, South Sulawesi, Indonesia**


Name of the authors: Ihsan, Kasmawati, Andi Asni, Ernaningsih, Asbar, Asmidar, Hasan E. Adimu

Hereby I would like to submit the manuscript entitled “**aquaculture management of blue swimming crab using integrated submerged net cage in pangkep regency waters, South Sulawesi, Indonesia**” to Aquaculture, Aquarium, Conservation & Legislation - International Journal of the Bioflux Society.


This manuscript was not submitted or published to any other journal.

The authors declare that the manuscript is an original paper and contain no plagiarised text. All authors declare that they are not currently affiliated or sponsored by any organization with a direct economic interest in subject of the article. My co-authors have all contributed to this manuscript and approve of this submission.


Author




Ihsan




Kasmawati




Andi Asni




Ernaningsih



Asbar



Asmidar



Hasan E. Adimu

*First author: *ihsan.ihsan@umi.ac.id*

BUKTI PEMBAYARAN JURNAL
BIOFLUX

aplikasi setoran/transfer/kliring/inkaso deposit/transfer/clearing/collection form



kepada to PT Bank Mandiri (Persero) Tbk

harap dilakukan transaksi berikut please do this transaction:

transaksi transaction ☐ setoran deposit ☐ TT tt ☐ RTGS rtgs ☐ SKNBI sknbi ☐ Kliring-Inkaso clearing-collection ☐ Bank Draft bank draft

harap ditulis dengan huruf cetak please fill in with block letters

VALIDASI validation

15216 1521657 1521654 29 04 24/09/2021 10:03:01 AM 4512
152-00-1488142-5 IHSAN IDR 4,715,000.00 DR
09-15216-0009106-02 USD 300.00 CR
25.00 1.0000000 14.400.0000000
PEMBAYARAN JURNAL BIOFLUX
TANGGAL EFEKTIF 24/09/2021

PENERIMA (wajib diisi) beneficiary ☐ perorangan individual ☒ perusahaan company ☐ pemerintah government

Status kependudukan resident status ☐ penduduk resident ☐ bukan penduduk non-resident

Nama name Bioflux SRL

Nomor rekening account number 213USD CRT00L2881401

Bank Bank Banca Transilvania

Alamat & telp penerima beneficiary address & phone no Romania

Jenis & Nomor Identitas ID type & number SWIFT Code of the bank BTR

TUJUAN TRANSAKSI purpose of transaction (wajib diisi) ☐ Tabungan / Investasi savings / investment ☐ Pembayaran payment ☐ Biaya hidup personal expenses ☐ Bisnis business purpose ☐ Pembelian barang / jasa purchase of goods / services ☐ Donasi / amal donation

BERITA TRANSAKSI transaction remarks Pembayaran Jurnal Bioflux

diisi oleh Bank filled out by the Bank

Jumlah transfer amount of transfer	
Komisi commission	
Biaya Pengiriman transfer fee (SWIFT/RTGS/SKNBI)	00001000
Biaya Koresponden correspondent charge	
Sub Total	
Kurs rate	
Total	24 SEP 2021

Pemohon dengan ini menyetujui syarat-syarat dan ketentuan yang tercantum dibalik formulir aplikasi ini
the applicant hereby accepts all terms and conditions stated on the reverse side of this transaction form

Pengesahan Bank bank's authorization 179266 Tanda tangan pemohon applicant's signature

Nama name

PENGIRIM (wajib diisi) applicant ☒ nasabah customer ☐ non nasabah walk in customer (WIC)

NIK / Paspor (WNA) ID number

Informasi pengirim applicant information ☒ perorangan individual ☐ perusahaan company ☐ pemerintah government

Status kependudukan resident status ☒ penduduk resident ☐ bukan penduduk non-resident

Nama name Dr Ir Ihean, M.S.

Alamat & nomor telepon address & telephone number Perum Dewi Kumala
Ae 9/4 Mahassa
085212304040

METODE TRANSAKSI (wajib diisi) method of transaction ☐ tunai cash ☒ debet rekening: 152001488142 ☐ cek/bilyet giro cheque

Bank Tertarik drawee bank	No. cek/BG cheque number	Valuta currency	Nominal amount
	1520014881425		

Jumlah setoran/transfer/kliring/inkaso deposit/transfer/clearing/collection/amount 300 USD

Terbilang in words Tiga ratus dollar

SUMBER DANA TRANSAKSI (wajib diisi) source of fund ☐ Gaji / penghasilan salary / income ☐ Tabungan / hasil investasi savings / investment ☐ Warisan inheritance ☐ Dana pemerintah Government Funds ☐ Hibah / hadiah Grants / gifts ☐ Penjualan aset sale of assets ☐ Hasil usaha business proceed ☐ Sumbangan contribution

BIAYA TRANSAKSI transaction fee ☐ Tunai cash ☐ debet rekening: ☐ debet account:

Biaya bank koresponden correspondent charge ☐ Pengirim applicant ☐ Penerima beneficiary ☐ Lainnya others



diisi apabila pembawa formulir bukan Pengirim filled out if the bearer of this form is not the applicant



Nama name



Alamat & nomor telepon address & telephone number



NIK / Paspor (WNA) ID number



KORESPONDENSI AUTOH BY E-MAIL
HASIL KORESPONDENSI SELAMA PROSES PENULISAN SAMPAI TERBIT
TERBIT JURNAL














Certificate/Letter of preliminary acceptance

This certificate shows that your paper:

**Aquaculture Management of Blue Swimming Crab using Integrated Submerged
Net Cage in Pangkep Regency Waters, South Sulawesi, Indonesia**

Authors:

Ihsan, Kasmawati, Andi Asni, Ernaningsih, Asbar, Asmidar, Adimu HE

at a preliminary evaluation is qualified for processing (reviewing process) in order to be
published with revision in volume 13 (2020)
of the scientific/academic journal: Aquaculture, Aquarium, Conservation & Legislation –
International Journal of the Bioflux Society.

AACL Bioflux is covered by Thomson ISI Web of Knowledge via:


- Zoological Record (Biosis) and
- CAB Abstracts (CABI)

Thank you for publishing with us!

Sincerely yours,
Editor Senior Researcher
Miklos Botha, PhD



NASKAH TELAH DIBERBITKAN
BIOFLUX



Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia

¹Ihsan, ¹Kasmawati, ¹Andi Asni, ¹Ernaningsih, ²Asbar, ²Asmidar, ³Hasan E. Adimu

¹ Fisheries Resources Utilization Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ² Marine Science Study Program, Faculty of Fisheries and Marine Science, Muslim University of Indonesia, Makassar, South Sulawesi; ³ Department of Marine Science, Faculty of Agriculture Fisheries and Animal Husbandry, Nineteen November University of Kolaka, Southeast Sulawesi, Indonesia. Corresponding author: Ihsan, ihsan.ihsan@umi.ac.id

Abstract. The present study aimed to determine the cultivation area suitability and submerged net cage model of blue swimming crab integrated with other aquaculture activity. This study was conducted from April to November 2017. Primary and secondary data were collected and analyzed for water suitability and land utilization using mapping and geographic information system analysis (GIS). The water quality parameters suggested different results: dissolved oxygen 5.1-10.7 mg L⁻¹, salinity 7-35 ‰, temperature 32-34.4°C, and pH 7-8 respectively. The findings suggested two large clustering aquaculture management area; cluster 1 (the area of 26,342 ha) which is located close to the shoreline and cluster 2 (172,953 ha) which is located far from the coastal line. The management of crab cultivation was integrated with other utilization using submerged net cage with seaweed farming. The suitable submerged net cage model for integrated crab cultivation are rectangular cage with 2 different sizes: (1) large size cages of 150 cm length, 100 cm of height and 100 cm width. The net used was made from nylon (PE Number 12), the material of the main pillar was threaded iron of number 12 and the width of number 7. The net applied was made from nylon Polyethylene (PE) number 12 with mesh size of 1.0 cm. All parts of the submerged net-cage materials were built from the same materials like the large size net-cages. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cage cultures.

Key Words: cultivation, cluster, farming, fishermen, production.

Introduction. Blue swimming crab (*Portunus pelagicus*) is a sought-after aquaculture commodity contributing significantly to local people and the national income of Indonesia. *P. pelagicus* is one of the major commodities for export with regards to the Indonesian fisheries sector (Wiyono & Ihsan 2018). The increase in market prices of *P. pelagicus* also has been attributed in part to increasing volumes of exports and domestic demands. The value of Indonesian swimming crab and crab exports in 2012-2017 increased each year by 0.67% and 6.06% (MMAF 2019). Many restaurants and hotels serve *P. pelagicus* as its seafood dish for their costumer. As a result, it attracts the attention of local fishermen to start catching crabs and farming crabs as a source of income. However, improper utilization management of crabs resources, as well as land utilization, has been the major issue of the *P. pelagicus* stock depletion and land utilization conflicts. Several studies related to the excessive catching of *P. pelagicus* in Indonesia were revealed by (Hamid & Wardiatno 2015; Hamid et al 2016; Sara et al 2019; Wagiyo et al 2019).

In order to enhance severely depleted stocks of *P. pelagicus*, a proper utilization marine-based culture management is required. The deficits may be minimized through conditioning strategies leading to enhance crab production (Young et al 2008). It can be

done using a composite culture of crabs with other cultured commodities such as finfish and seaweed. It may prevent the conflicts of spatial use in the coastal areas of Pangkep Regency.

Aquaculture management of blue swimming crab using integrated submerged net cage is one of the management strategies that may increase the production of blue swimming crab as well as maintaining the stock of crab and the crab supply for the market. These management system changes have increased in recent years. The advantages of this management are easy control, efficiency, fast monitoring, and the possibility of incorporating several forms of automation (Tavares et al 2017).

The management of the crab cultivation is expected to be carried out in an integrated manner with other land utilization such as seaweed farming and other finfish aquaculture. Therefore, it is able to solve the conflicts of space utilization for the crab cultivation. The most important aspect of the management of integrated crab cultivation is the submerged net cages models. In line with this, the research objective consisted of: 1) Determining the management of crab cultivation in an integrated manner with other spatial utilization activities in the waters of Pangkep Regency; 2) Assessing the use of submerged net cage model for crab cultivation.

Material and Method

Description of the study sites. The present study was conducted at Pangkep waters from April to November 2017 (Figure 1).




Figure 1. The location of Pangkep waters, South Sulawesi, Indonesia.

Data collection. Materials and equipment used during the period of study included *P. pelagicus* larvae, submerged net cage, current meter, hand refractometer, dissolved oxygen (DO) meter, underwater camera, thermometer, global positioning system, label, stationery, cold-boxes, Secchi disk, check-list form. Data consisted of primary and secondary data. Primary data was obtained in situ observation to determine the clustering aquaculture area for crab cultivation based on the aspects of physics and

chemistry of oceanography. Twenty eight of sampling sites were used for data collection. The Global Positioning System (GPS) was used to determine the observation location or station. The secondary data was obtained from relevant agencies with the present study. Matrix and conformity criteria were applied for management suitability analysis, according to Ihsan (2015), (Tables 1-3).

Table 1

Criteria for determining the suitability of an area

X0 (= score value) – X1 (= X0 + Ci)	Unsuitable (N)
X1 – X2 (= X1 + Ci)	Acceptable (S2)
X2 – X3 (= max value)	Suitable (S1)

X0 - lower limit of assessment scale; X1 - limit between unsuitable and acceptable class; X2 - limit between acceptable and suitable class; X3 - upper limit of assessment scale.

Table 2

Suitability matrix for adult *Portunus pelagicus*

Parameters	Suitable category				Unsuitable category		
	Weight	Description of suitable parameters	Score	Total Score	Description of unsuitable parameters	Score	Total Score
Temperature	0.40	28-31°C	2	0.80	<28 or >31°C	1	0.40
Salinity	0.30	31-36‰	2	0.60	< 31 or 36‰>	1	0.30
Dissolved oxygen	0.20	4-6 ppm	2	0.40	<4 ppm or >6 ppm	1	0.20
pH	0.10	6.78 – 8.0	2	0.20	<6.78 or >8.0	1	0.10
	1.00	Total maximum score		2.00	Total minimum score		1.00

Table 3

Determination of area suitability category

Interval class	Category
1.00–1.33	Not suitable (N)
1.34–1.67	Fairly suitable (S2)
1.68-2.00	Suitable (S1)

Data analysis. Data were descriptively and quantitatively analyzed including the suitability data of the aquatic environment. The geographic information systems (GIS) with ArcView method was applied.

Results

Integrated management of *P. pelagicus* farming. The management of *P. pelagicus* farming using submerged net cage in an integrated manner in the waters of Pangkep Regency has been recognized as aquaculture region. Due to efficiency and effectiveness in crab farming with submerged net cage, it is necessary to determine the suitability of the crab aquaculture in Pangkep waters based on location of crab cultivation. The object of observation can be grouped based on the characteristics possessed by the object of observation. Grouping observational objects was performed by considering the homogeneity level of the observation object in one cluster while maximizing heterogeneity between clusters (Yuniati & Rachman 2016). Clustering the location of aquaculture region and marine space management was based on the efficiency and the effectiveness of *P. pelagicus* using submerged net cage with consideration of distance of location from coastal line or islands (Table 1).

The results showed that the clustering aquaculture area can be divided into two large clusters, namely cluster 1 (the area of 26,342 ha) which the location of crab cultivation is integrated with the location of seaweed cultivation, located close to the shoreline (Figure 2). Most methods of seaweed farming in the waters of Pangkep Regency included long line method. The aquaculture method is integrated where on the surface of the water seaweed farming and crab cultivation is resided in the bottom of the waters of seaweed cultivation. Meanwhile, cluster 2 (172,953 ha) where the location of crab cultivation integrated with the location of seaweed cultivation, located far from the coastal line (Figure 2). It leads to an increase in the operational cost for monitoring and feeding. Nonetheless, the crab cultivation using a submerged net cage does not need much operational cost because the fishermen can do both monitoring and fishing at the same time. The cluster area 2 was not influenced significantly by water quality changes particularly, declining of temperature and salinity. There are no much pollution inputs from upper land such as river, sedimentation, and effluent waste.




Figure 2. Suitable area for blue swimming crab farming using submerged net cage in Pangkep Regency.

In addition, the coordinate points of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency is presented in Table 4.

In the present study, the unsuitable area of 3,590.74 ha for crab farming is considered. This location is directly adjacent to the shoreline leading to poor water quality due to freshwater input from rivers and sedimentations. The water at this location is also very shallow. It is not suitable for submerged net cage. The depth requirements of the location of crab cultivation for submerged net cage should be 1.5-2 times of the size of submerged net cage (Ihsan et al 2017).

Integrated management of aquaculture is intended to prevent the conflicts interest concerning the use of area for aquaculture and other purposes. Not only seaweed cultivation can be integrated with crab farming but also other finfish aquaculture using floating net cages can also be performed. The map of integration of crab cultivation using submerged net cage with other activities is presented in Figure 3.

Table 4

The coordinates of the location of the suitability of the crab cultivation land after clustering in the waters of Pangkep Regency

Coordinates		Criteria
X	Y	
119° 35' 14.662" E	4° 34' 37.872" S	Not suitable
119° 33' 36.552" E	4° 37' 9.734" S	
119° 31' 25.095" E	4° 40' 16.154" S	
119° 29' 1.241" E	4° 45' 39.504" S	
119° 28' 40.538" E	4° 49' 7.916" S	
119° 30' 6.122" E	4° 51' 57.196" S	
119° 28' 49.413" E	4° 52' 25.889" S	Cluster 1
119° 27' 36.596" E	4° 49' 6.426" S	
119° 27' 25.498" E	4° 45' 22.629" S	
119° 30' 24.496" E	4° 39' 45.164" S	
119° 32' 38.749" E	4° 36' 39.597" S	
119° 34' 16.040" E	4° 34' 17.208" S	
119° 23' 1.902" E	4° 34' 44.186" S	Cluster 2
119° 20' 55.266" E	4° 45' 4.512" S	
119° 20' 42.457" E	4° 55' 45.986" S	




Figure 3. Map of management of crab cultivation in submerged floating net cages integrated in the waters of Pangkep Regency.

Water quality measurement. For the suitability determination of clusters 1 and cluster 2 for crab cultivation was done using the physical and chemical oceanographic aspects. The results of water quality measurements at 28 stations during survey period revealed: temperature of 27-34°C, salinity of 28-34 ppt, and pH of 6.50-8.00. The optimal water

pH for fish and shrimp life is 6.5-8 (neutral) (Wardhany et al 2018). This pH level promotes optimal balance between oxygen and carbon dioxide which inhibits the growth of various harmful microorganisms in the water. The optimal pH water for crab cultivation is 6.5 and 8.5 (Juwana 1993). Furthermore at pH 9.5-11.0 and 4.0-6.0 lead to low production and lower than 4.0 or higher 11.0 have detrimental effect on aquatic organisms causing high mortality.

DO measurements revealed a concentration range of 5.17-7.97 mg L⁻¹. DO-dissolved oxygen content is an important limiting factor for biota, and the range of values is 5.1-9.2 mg L⁻¹ which is considered optimal and supports the life of marine biota (ME 2004). The optimal water temperature for crab cultivation is 26-32°C (Adiwijaya et al 2002). However, the crabs are very tolerant concerning water temperatures ranging between 17 and 37°C (Perkins 1974). Study on crab cultivation in ponds in Laikang Village revealed the following parameters: salinity 30-43 ppt, DO 4-10 mg L⁻¹, pH 4-8 and temperatures of 27-34°C.

Submerged net cage model for *P. pelagicus* farming. A rectangular model of submerged net cage was used: 1) submerged net cage size was 150 cm length and 100 cm width and 100 cm height; 2) small submerged net cage size was 75 cm length, 50 cm width and 50 cm height using a net made of nylon (PE No. 12). For the net cage framework wrapping net was used and tied properly. The use of wood as a framework for making submerged net cage is much durable compared to iron which much prone to be corrosive.

The cultivation of *P. pelagicus* can be done in the earthen pond with one log pond and two grow-out ponds. The crablets were reared for 1-2 months until the sexual dimorphism was evident. Furthermore, the grow-out of crab was carried out by applying a floating cage system or a net installed near the beach. The rearing ponds size was 50 x 100 m. Ponds of this size can accommodate up to 10,000 crabs. The submerged net cage model used is presented in Figure 4.



Figure 4. Submerged net cage model for *Portunus pelagicus* farming. (A) Large submerged net cage of 150 cm length, 100 cm main pole and small submerged net cage of 75 cm length, height 50 cm of threaded steel number 12; (B) Large submerged net crab of 100 cm width of main pole and a small cage of 50 cm steel size 7 (normal); (C) Nets (nylon) PE / 12 mesh size 1.0 cm; (D) submerged net cage door size 30 x 50 cm (E) submerged net cage door leaf size 30 x 50, frame made of steel and use a PA / 12 size net; (F) Plastic strap No PE/7 mm, submerged net cage hangers.

To simplify the submerged and lift process, a plastic PE no 7 mm was used. The length of the plastic strap depends on the depth of the water. The length of plastic rope used during the study was 5-10 m. Besides that, it is expected that the depth of the water with the length of the rope must be different. If the water depth is 7 meters, the length of the rope was approximately 10 m. So that, during tide, the submerged net cage was visible from the surface of the water.

Conclusions. Management of submerged net cage can be integrated with other aquaculture activity, such as seaweed cultivation, and other finfish floating net cages. The submerged net cage system is considered suitable for integrated crab cultivation with a rectangular model; 1) large size cages, 2) small size cages, with the same size and models.

Acknowledgements. The authors would like to thank the Directorate of Research and Community Service, the Directorate General for Research and Development of the Ministry of Research, Technology and Higher Education for funding the present research through Applied Research for Product. Our gratitude are also extended to researcher colleagues for their significant contributions to this paper.

References

- Adiwijaya D., Jaya S., Sugeng, Sutikno E., 2002 [Business opportunity for brackish water commodity: blue swimming crab (*Portunus pelagicus*) cultured in earthen pond]. BBPBAP, Jepara 13-20. [In Indonesian].
- Hamid A., Wardiatno Y., Lumbanbatu D. T. F., Riani E., 2016 Stock status and fisheries exploitation of blue swimming crab *Portunus pelagicus* (Linnaeus 1758) in Lasongko Bay Central Buton, Indonesia. Asian Fisheries Science 29:206-219.
- Hamid A., Wardiatno Y., 2015 Population dynamics of the blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) in Lasongko Bay, Central Buton, Indonesia. AACL Bioflux 8(5):729-739.
- Ihsan, 2015 [Sustainable utilization of blue swimming crab (*Portunus pelagicus*) in Pangkep Regency Sulawesi Selatan]. Dissertation, Bogor (ID): Graduate School, Bogor Agricultural University, Bogor, Indonesia. [In Indonesian].
- Ihsan, Asbar, Asmidar, 2017 Fisheries management of blue swimming crab based on spatial and marine culture in submerged net cage for national food security. Final Report 2017, Penelitian terapan unggulan perguruan tinggi (PTUPT). Universitas Muslim Indonesia, Makassar, Indonesia. [In Indonesian].
- Juwana S., 1993 [Effect of light intensity salinity and temperature on survival rate and growth performance of blue swimming crab (*Portunus pelagicus*) seeds]. Majalah Ilmu Kelautan 16:194-204. [In Indonesian].
- Perkins F. J., 1974 The biology of estuarine and coastal waters. Academic Press Inc. (London) Ltd. New York, pp. 670-678.
- Sara L., Astuti O., Muzuni, Safilu, 2019 Status of blue swimming crab (*Portunus pelagicus*) population in Southeast Sulawesi waters, Indonesia. AACL Bioflux 12(5):1909-1917.
- Tavares C. P. S., Silva U. A. T., Pereira L. A., Ostrensky A., 2017 Systems and techniques used in the culture of soft-shell swimming crabs. Reviews in Aquaculture 10(4):913-923.
- Wagiyo K., Tirtadanu, Ernawati T., 2019 [Fishery and population dynamics of blue swimmer crabs (*Portunus pelagicus* Linnaeus, 1758) In Jakarta Bay]. Jurnal Penelitian Perikanan Indonesia 25(2):79-92. [In Indonesian].
- Wardhany V. A., Yuliandoko H., Subono, Harun A. M. U., Astawa I. G. P., 2018 [Temperature control system based on Fuzzy Logic, pH and water salinity in *Vanammei* shrimp farms]. International Electronics Symposium on Engineering Technology and Applications (IES-ETA), Bali, pp. 145-149. [In Indonesian].
- Wiyono E. S., Ihsan 2018 Abundance, fishing season and management strategy for blue swimming crab (*Portunus pelagicus*) in Pangkajene Kepulauan, South Sulawesi,

- Indonesia. Tropical Life Sciences Research 29(1):1–15.
- Young A. C., Johnson E. G., Davis J. L. D., Hines A. H., Zmora O., Zohar Y., 2008 Do hatchery-reared blue crabs differ from wild crabs, and does it matter? Reviews in Fisheries Science 16(1-3):254-261.
- Yuniati, Rachman F., 2016 [Potential cluster for fisheries in public waters, East Java 2016]. Seminar Nasional Pendidikan, Sains dan Teknologi ISBN: 978-602-61599-6-0 Faculty of Math and Biology Science, Muhammadiyah University of Semarang. ISBN: 978-602-61599-6-0. [In Indonesian].
- *** ME (Ministry of Environment), 2004 [Ministry Decree No. 51/2004 about water quality standard for sea water]. Ministry of Environment, Indonesia. [In Indonesian].
- *** MMAF (Ministry of Marine Affairs and Fisheries), 2019 Prospects of Indonesian Rajungan and Crab Export Market to China. Ministry of Marine Affairs and Fisheries, Indonesia. [In Indonesian].

Received: 10 July 2020. Accepted: 27 October 2020. Published online: 03 November 2020.

Authors:

Ihsan, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ihsan@umi.ac.id

Kasmawati, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: kasmawati@gmail.com

Andi Asni, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asni.andi@yahoo.co.id

Ernaningsih, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: ernaningsih36@yahoo.co.id

Asbar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asbar_umi1964@yahoo.com

Asmidar, Muslim University of Indonesia, Faculty of Fisheries and Marine Science, Department of Fisheries Resources Utilization, Indonesia, South Sulawesi 90231, Makassar, Jl. Urip Sumoharjo No.5, Panaikang, Kec. Panakkukang, e-mail: asmidar.darwis@gmail.com

Hasan Eldin Adimu, Sembilanbelas November University of Kolaka, Faculty of Agriculture Fisheries and Animal husbandry, Department of Marine Science, Indonesia, Southeast Sulawesi 93561, Kampus B Buton Tengah, Kec. Mawasangka, e-mail: eldin.kahedupa@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Ihsan, Kasmawati, Asni A., Ernaningsih, Asbar, Asmidar, Adimu H. E., 2020 Aquaculture management of blue swimming crab (*Portunus pelagicus*) using integrated submerged net cage in Pangkep Regency waters, South Sulawesi, Indonesia. AACL Bioflux 13(6):3279-3286.