

DNA barcoding of Telmatherinidae family in Lake Towuti, South Sulawesi, Indonesia

by St Hadijah


Submission date: 10-Jul-2023 08:35AM (UTC+0800)

Submission ID: 2128722761

File name: 2019.1208-1215.pdf (117.46K)

Word count: 4308

Character count: 23817



9

DNA barcoding of Telmatherinidae family in Lake Towuti, South Sulawesi, Indonesia

¹Jayadi Jayadi, ¹Ilmiah Ilmiah, ¹Siti Hadijah, ¹Muhammad Kasnir,
²Dewi I. Roslim

8

¹Department of Aquaculture, Faculty of Fishery and Marine Science, Indonesia Muslim University, Makassar, Indonesia; ²Department of Biology, Faculty of Mathematics and Nature Sciences, Universitas Riau, Pekanbaru, Indonesia. Corresponding author: J. Jayadi, jayadi_fatril@yahoo.com; jayadi.jayadi@umi.ac.id

7

Abstract. Telmatherinidae is a family of endemic fish in South Sulawesi. The study is aimed at identifying molecular gene of the endemic fish of the Telmatherinidae family in Towuti Lake, South Sulawesi, to identify and analyze genetic diversity, genetic marking, genetic distances, genetic characterization, and dendrogram of the fish. DNA barcoding in this research used the Cytochrome c Oxidase I (COI) gene. Amplification of mitochondrial COI gene regions was conducted by using COI Fish F2 and COI Fish R 2 primer. Data analysis, total digestion of DNA, Polymerase Chain Reaction (PCR), Electrophoresis, gel purification, and sequencing Basic Local Alignment Search Tool (BLAST) were performed. The results showed that DNA sequence was 681 bp. Meanwhile, analysis of dendrogram suggested that the fish of Telmatherinidae family in Towuti Lake are similar to the fish in the *Paratherina*, *Telmatherina* and *Tominanga* genera. The genus *Telmatherina* including *Telmatherina celebensis*, *Telmatherina bonti*, and *Telmatherina opudi* showed 85% significant shared similarity with *Atherina* sp. Only 84% similarity of genus *Telmatherina* was found in *Pristigenys alta*, *Parexocoetus brachypterus*, *Cypselurus hiraii*, and *Hypoatherina tsurugae*. Meanwhile, genus *Paratherina*: *Paratherina striata* and *Paratherina wolterecki* have 83-84% similarity with *Scorpius lineolata*, *Hyporhamphus affinis*, *Cypselurus hiraii*, *Parexocoetus brachypterus*, and *Pristigenys alta*. In addition, genus *Tominanga*: *Tominanga sanguicauda* has 84% similarity with *Hypoatherina tsurugae*, *Cypselurus hiraii*, *Prognichthys sealei*, *Hyporhamphus affinis*, and *Pristigenys alta*.

Key Words: genetic diversity, Telmatherinidae, COI gene, dendrogram, Lake Towuti.

22

Introduction. Lake Towuti is located in Nuha sub-district, East Luwu regency South Sulawesi, Indonesia, and is interconnected with Lake Matano and Lake Mahalona. These three lakes are included in an ancient Lake Complex (Wijaya et al 2009; Samuel et al 2009; Nasution et al 2010; Umar et al 2012). Lake Towuti has been an important habitat for several endemic freshwater fish (Kottelat et al 1993; Wirjoatmodjo et al 2003; Samuel et al 2009; Parenti 2011; Parenti & Ebach 2013) such as the fish families from Telmatherinidae, Gobidae, Adrianichthyidae and Hemiramphidae (Herder et al 2006; Nasution et al 2007; Stelbrink et al 2014; Hutama et al 2016). They are economically important for many local communities around Lake Towuti (Wijaya et al 2009).

Potential productivity of fish in Lake Towuti is equal to ± 195 ton year⁻¹ (Wijaya et al 2009). Some endemic fish species are utilized for consumption and ornamental purposes. Consequently, the endemic fish population continues to decline yearly, and conservation measures have not been carried out properly (Mamangkey et al 2007; Wijaya et al 2009; Samuel et al 2009; Nasution et al 2010; Umar et al 2012). The uncontrolled exploitation of endemic fish can lead endemic fish into extinction, even in their natural habitat (Jayadi et al 2015, 2016; Nasution et al 2015).

Another threat that causes a decline of endemic fish populations in Lake Towuti is water pollution from domestic waste disposal. Pollution detrimentally affects the environment as well as endemic fish. Moreover, land clearing for settlement and agriculture around the lake leads to erosion and sedimentation. This is exacerbated by introducing invasive fish species into Lake Towuti (Prianto et al 2014).

1

Several endemic fish species included in the category of vulnerable species in Lake Towuti include Termatherinidae family of genera *Telmatherina*, *Paratherina*, and *Tominanga* (Kottelat et al 1993; IUCN 2003; Suwelo 2005; Herder et al 2006; Nasution et al 2007; Stelbrink et al 2014; Hutama et al 2016). The Telmatherinidae family was referred to as sailfin silversides fish (Kottelat 1991; Stelbrink et al 2014; Hutama et al 2016). Therefore, it is necessary to conduct sustainable management, such as genetic conservation of species and habitat.

Genetic conservation is one of sustainable management measures that can be applied by analyzing the genetic diversity of particular species (Mamangkey et al 2007; Hadijah et al 2014; Jayadi et al 2015). Genetic diversity analysis provides the short and long term information related to a particular species' population (Ferguson et al 1995). For instances, genetic diversity can be used in considering biological resource management (Yusron 2005; Jayadi et al 2015; Nugroho et al 2017). Furthermore, it can be used for improving fish stock (Islam et al 2011), domestication, and aquaculture (Lante et al 2011; Iskandariah et al 2015; Lorenzen et al 2012; Jayadi et al 2016). To analyze the genetic diversity, barcoding DNA using COI can be applied (Ward et al 2005; Muchlisin et al 2013; Jusmaldi et al 2014; Hubert et al 2015; Nuryanto et al 2017; Pramono et al 2017; Abbas et al 2017). The purpose of this study is to determine genetic diversity using molecular identification such as the COI DNA gene in the endemic species of the Telmatherinidae family in Lake Towuti as well as genetic marking, genetic characters, and dendrogram.

Material and Method

Fish sampling. Fish sampling for Telmatherinidae fish family was conducted in Lake Towuti, South Sulawesi, Indonesia from January to June, 2018, using fishing net size of 1 mm. Morphological identification was carried out according to Herder et al (2006), Kottelat (1991), Kottelat et al (1993), and Said & Hidayat (2015). In addition, fish muscle (from 15 fish) in the tail was taken (1 cm x 1 cm) and then placed into a 96% ethanol solution and stored in the freezer.

Total DNA isolation. D NEasy Blood and tissue kit (cat. No. 69504 and 69506 Qiagen) was applied for total DNA. The total DNA isolation process was conducted based on the existing protocol. The electrophoresis technique was performed to measure the quality and the quantity of the total DNA.

Polymerase Chain Reaction (PCR). Total DNA was amplified by PCR technique using a pair of universal primers for COI genes in fish, namely Fish F2: 5'-TCG ACT AAT CAT AAA GAT ATC GGC AC-3' and FishR2: 5' TCA ACT GGG TGA CCG AAG AAT CAG AA -3' (Ward et al 2005). The PCR components include 1X of Supreme NZY Taq, 2X Green Master Mix, 2.4 µM of forward primer, 2.4 µM reverse primer, 1 µL of total DNA, and dH₂O until 50 µL of PCR volume. The PCR program includes pre-PCR at 94°C for 5 minutes; PCR was 35 cycles consisting of denaturation at 94°C for 45 seconds, annealing at 50°C for 45 seconds, elongation at 72°C for 1 minute, and post-PCR at 72°C for 10 minutes. PCR success was detected by electrophoresis techniques.

Electrophoresis. Electrophoresis was performed to detect the success of DNA and PCR isolation. Total DNA and PCR products were moved to a 1% agarose gel in a 1X TBE buffer at 50 voltage for 45 minutes. DNA tape was colored using 5 µg mL⁻¹ of ethidium bromide, visualized on a UV transilluminator lamp, and then photographed using a UV filtered digital camera.

DNA sequencing. DNA sequencing was performed at PT Genetika Science in Jakarta as a channeling agent. Gene purification and sequencing were carried out at 1st Base in Malaysia. The PCR product was 40 µL and each primer was 30 µL.

Data analysis. DNA sequence data was used forward and reverse primers which were put together or aligned using BioEdit7 software. The BLAST (Basic Local Alignment Search Tool) analysis at <http://www.ncbi.nlm.nih.gov/BLAST> (Madden 2013) was carried out on DNA sequences from each fish sample to determine its similarity to DNA COI sequences in the GenBank database. Accessions were of high similarity, and sequences were downloaded to make phylogenetic trees using MEGA (molecular evolutionary genetics analysis) version 6.17.5 (build #: 6140226) (Tamura et al 2013) based on the Kimura 2-parameter model and UPGMA (Unweighted Pair Group Method with Arithmetic mean) with 1000 bootstrap.

Results and Discussion. Amplification of fish samples from the Telmatherinidae family using primary Fish_F2/Fish_R2 produced 750 bp of DNA tape (Figure 1). The DNA band that has been produced indicates that the sequencing process was done properly. The DNA sequence obtained from 20 samples was 681 bp and registered in Gen Bank (Table 1).




Figure 1. Profiles of DNA bands that use Fish_F2 / Fish_R2 primers. L = 1 kb DNA Ladder (Thermo Scientific).

Table 1
Registration number of DNA sequence results of 20 fish samples from the family of Telmatherinidae in Lake Towuti

No	Fish spesies	Sample label	Registration number
1	<i>Telmatherina celebensis</i>	A01	MH568798
2	<i>Telmatherina celebensis</i>	A02	MH568799
3	<i>Paratherina striata</i>	B01	MH568800
4	<i>Paratherina striata</i>	B02	MH568801
5	<i>Tominanga sanguicauda</i>	C01	MH568802
6	<i>Tominanga sanguicauda</i>	C02	MH568803
7	<i>Tominanga sanguicauda</i>	C03	MH568804
8	<i>Paratherina wolterecki</i>	D10	MH568805
9	<i>Paratherina wolterecki</i>	D11	MH568806
10	<i>Paratherina wolterecki</i>	D12	MH568807
11	<i>Telmatherina bonti</i>	E13	MH568808
12	<i>Telmatherina bonti</i>	E14	MH568809
13	<i>Paratherina cyanea</i>	F15	MH568810
14	<i>Paratherina cyanea</i>	F16	MH568811
15	<i>Paratherina cyanea</i>	F17	MH568812
16	<i>Telmatherina opudi</i>	G18	MH568813

DNA barcodes of endemic fish in the Telmatherinidae family in Lake Towuti using primary COI Fish_F2 / Fish_R2 found long DNA band fragments after mt-DNA amplification to around 681 bp. The COI primer has been applied to identify the genus *Thunnus* and

genus *Squalus* with a fragment length of 655 bp (Ward et al 2005), the genus *Kryptopterus* around 707 bp (Jusmaldi et al 2014), and the genus *Mystus* at 697 bp (Pramono et al 2017). The length of fragments in endemic fish in South Sulawesi such as *Glossogobius matanensis* was 500 bp (Mamangkey et al 2007), *Glossogobius aureus* was 600 bp (Hadijah et al 2014), *Telmatherina ladigesii* was 600 bp (Jayadi et al 20015), *Pterygoplichthys* sp. was 650 bp (Rosnaeni et al 2017), Family Sparinidae was 650 bp (Abbas et al 2017), and *Harpodon nehereus*, *Harpodon microchir* and *Harpodon squamosus* were 618 bp (Nugroho et al 2017).

DNA barcoding is used to assign a biological specimen to a species (Ardura et al 2010; Fahmi et al 2016; Nuryanto et al 2017; Pramono et al 2017). Furthermore, DNA barcoding contributes to science and promotes more sustainable practices in taxonomy and the development of new molecular tools for species identification (Hubert et al 2015).

The results of the Telmatherinidae family dendrogram analysis obtained three genera, namely: *Paratherina*, *Telmatherina*, *Tominanga* (Figure 2).




Figure 2. Dendrogram in the Telmatherinidae family in Lake Towuti.

Phylogenetic results show that three genera from the Telmatherinidae family (*Paratherina*, *Telmatherina* and *Tominanga*) were found in Lake Towuti. Species of the genus *Paratherina* are *Paratherina striata*, *Paratherina wolterecki*, *Paratherina cyanea*, and the genus *Telmatherina* is represented by *Telmatherina celebensis*, *Telmatherina bonti*, *Telmatherina opudi*, and the genus *Tominanga* is represented by *Tominanga sanguicauda* (Figure 2). All types of fish have been registered in Gen Bank (Table 1). Those fish population are known as native and endemic in Lake Towuti (Kottelat 1991; Parenti 2011; Stelbrink et al 2014; Hutama et al 2016).

The population of fish species from the Telmatherinidae family in Lake Towuti has decreased (Kottelat et al 1993; Suwelo 2005; Herder et al 2006; Nasution et al 2007; Stelbrink et al 2014; Hutama et al 2016) and have become vulnerable species (IUCN 2003). Consequently, inbreeding in the population leads to lower genetic variation of Telmatherinidae family. Low value of genetic diversity shows that a narrow level of migration can provide opportunities for limited gene exchange with other populations (Sugama et al 1996). Genetic variation of fish populations in nature can reduce due to damaged habitat, limited migration, isolated from other populations, depression inflation and decreased reproductive ability (Jayadi et al 2015). Changes in genes can occur due to individual gene migration in a population (Ezilrani & Christopher 2015).

In addition, genetic variation is important for evaluating fish resources in the wild. Genetic variations have a direct or indirect potential impact on populations, communities, and ecosystems (Hughes et al 2008). As said by Ezilrani & Christopher (2015), the genetic structure of a fish species is an illustration of the long changes due to biological and environmental factors. The results of the BLAST-GEN analysis for endemic fish of Telmatherinidae family in the Gen Fish Bank are presented in Table 2.

Table 2

Analysis of BLAST-GEN in the Telmatherinidae family

Fish species	Description	Max score	Total score	Query cover	E value	Indent	Accession
<i>Telmatherina celebensis</i>	Atherinidae sp.	673	673	95%	0.0	85%	MF409492.1
	<i>Scorpius lineolata</i>	685	685	99%	0.0	84%	AP011063.1
	<i>Pristigenys alta</i>	673	673	99%	0.0	84%	MG856600.1
	<i>Hypoatherina tsurugae</i>	673	673	99%	0.0	84%	AP004420.1
	<i>Cypselurus hiraii</i>	667	667	99%	0.0	84%	AB182653.1
<i>Paratherina striata</i>	<i>Scorpius lineolata</i>	673	673	99%	0.0	84%	AP011063.1
	<i>Hyporhamphus affinis</i>	656	656	98%	0.0	84%	KJ013045.1
	<i>Cypselurus hiraii</i>	656	656	99%	0.0	83%	AB182653.1
	<i>Parexocoetus brachypterus</i>	650	650	99%	0.0	83%	KY067947.1
	<i>Pristigenys alta</i>	644	644	99%	0.0	83%	MG856600.1
<i>Tominanga sanguicauda</i>	<i>Hypoatherina tsurugae</i>	685	685	99%	0.0	84%	AP004420.1
	<i>Cypselurus hiraii</i>	679	679	99%	0.0	84%	AB182653.1
	<i>Prognichthys sealei</i>	673	673	99%	0.0	84%	KY067951.1
	<i>Hyporhamphus affinis</i>	667	667	98%	0.0	84%	KJ013045.1
	<i>Pristigenys alta</i>	662	662	99%	0.0	84%	MG856600.1
<i>Paratherina wolterecki</i>	<i>Pristigenys alta</i>	685	685	99%	0.0	84%	MG856600.1
	<i>Cypselurus hiraii</i>	685	685	99%	0.0	84%	AB182653.1
	<i>Hypoatherina tsurugae</i>	679	679	99%	0.0	84%	AP004420.1
	<i>Cheilopogon doederleinii</i>	673	673	99%	0.0	84%	AP017897.1
	<i>Cheilopogon arcticeps</i>	673	673	99%	0.0	84%	KU360728.1
<i>Telmatherina bonti</i>	<i>Scorpius lineolata</i>	673	673	95%	0.0	85%	MF409492.1
	<i>Hyporhamphus affinis</i>	679	679	99%	0.0	84%	MG856600.1
	<i>Cypselurus hiraii</i>	679	679	99%	0.0	84%	KY067947.1
	<i>Parexocoetus brachypterus</i>	673	673	99%	0.0	84%	AB182653.1
	<i>Pristigenys alta</i>	673	673	99%	0.0	84%	AP004420.1
<i>Paratherina cyanea</i>	<i>Pristigenys alta</i>	685	685	99%	0.0	84%	MG856600.1
	<i>Cypselurus hiraii</i>	685	685	99%	0.0	84%	AB182653.1
	<i>Hypoatherina tsurugae</i>	679	679	99%	0.0	84%	AP004420.1
	<i>Cheilopogon doederleinii</i>	673	673	99%	0.0	84%	AP017897.1
	<i>Cheilopogon arcticeps</i>	673	673	99%	0.0	84%	KU360728.1
<i>Telmatherina opudi</i>	Atherinidae sp.	673	673	95%	0.0	85%	MF409492.1
	<i>Pristigenys alta</i>	679	679	99%	0.0	84%	MG856600.1
	<i>Parexocoetus brachypterus</i>	679	679	99%	0.0	84%	KY067947.1
	<i>Cypselurus hiraii</i>	673	673	99%	0.0	84%	AB182653.1
	<i>Hypoatherina tsurugae</i>	673	673	99%	0.0	84%	AP004420.1

Table 2 illustrates that the genus *Telmatherina* including *Telmatherina celebensis*, *Telmatherina bonti* and *Telmatherina opudi* showed 85% shared significant similarity to Atherinidae sp. Only 84% similarity of genus *Telmatherina* was found in *Pristigenys alta*, *Parexocoetus brachypterus*, *Cypselurus hiraii*, and *Hypoatherina tsurugae*. Meanwhile, genus *Paratherina*: *Paratherina striata*, *Paratherina wolterecki* have similarities between 83 and 84% with *Scorpius lineolata*, *Hyporhamphus affinis*, *Cypselurus hiraii*, *Parexocoetus brachypterus*, and *Pristigenys alta*. In addition, genus *Tominanga*: *Tominanga sanguicauda* has 84% similarity to *Hypoatherina tsurugae*, *Cypselurus hiraii*, *Prognichthys sealei*, *Hyporhamphus affinis*, and *Pristigenys alta*.

Conclusions. The use of the cytochrome oxidase I (COI) DNA has been successful in identifying the family of endemic fish from Telmatherinidae family in Lake Towuti, South Sulawesi Indonesia. The research finding shows that the genus *Paratherina* is represented by the species *Paratherina striata*, *Paratherina wolterecki*, *Paratherina cyanea*, the genus *Telmatherina* - *Telmatherina celebensis*, *Telmatherina bonti*, *Telmatherina opudi*, and the genus *Tominanga* - *Tominanga sanguicauda*.

Acknowledgements. The authors would like to thank the Directorate Research and Community Service, Ministry of Research, Technology and Higher Education for funding this research through Scheme on Basic Research of Higher Education Research Grants in 2018. We also thank to institutions of Research and Community Service, Muslim Indonesia University and Department of Biology Faculty of Mathematics and Nature Science of Riau University for analysis of genetics.

References

- Abbas E. M., Soliman T., Al-Magd M. A., Farrag M. S. M., Ismail R. F., Kato M., 2017 Phylogeny and DNA barcoding of the family Sparidae inferred from mitochondrial DNA of the Egyptian waters. *Jurnal of Fisheries and Aquatic Science* 12(2):73-81.
- Ardura A., Linde A. R., Moreira J. C., Garcia-Vazquez E., 2010 DNA barcoding for conservation and management of Amazonian commercial fish. *Biological Conservation* 143:1438-1443.
- Ezilrani P., Christopher J. G., 2015 Genetic variation and differentiation in African catfish, *Clarias gariepinus*, assessed by heterologous microsatellite DNA. *Indian Journal of Biotechnology* 14:388-393.
- Fahmi M. R., Prastio A. B., Kusumah R. V., 2016 [Barcoding of DNA of ornamental fishes in peatlands]. *Jurnal Riset Akuakultur* 11(2):137-145. [in Indonesian]
- Ferguson A., Taggart, Prodohl T. B., McMeel O., Thompson C., Stone C., McGinnity P., Hynes R. A., 1995 The application of molecular markers to study and conservation of fish population, with special reference to salmo. *Journal of Fish Biology* 47:103-126.
- Hadijah S., Tuwo A., Jusoff K., 2014 Genetic diversity population of Beloso (*Glossogobius aureus*) in Lake Tempe, Sulawesi, Indonesia. *World Journal of Fish and Marine Sciences* 6(1):87-97.
- Herder F., Schwarzer J., Pflander J., Hadiaty R. K., Schliwen U. K., 2006 Preliminary checklist of sailfin silversides (Teleostei: Telamtherinidae) in the Malili Lakes on Sulawesi. *Verhandlungen der Gesellschaft für Ichthyologie Band 5*:159-163.
- Hubert N., Kadarusman, Wibowo A., Busson F., Caruso D., Sulandari S., Nafqoh N., Pouyaud L., Ruber L., Avarre J. C., Herder F., Hanner R., Keith P., Hadiaty R. K., 2015 DNA barcoding Indonesia freshwater fishes: challenges and prospects. *DNA Barcodes* 3:144-169.
- Hughes A. R., Imouye B. D., Johnson M. T. J., Underwood N., Vellend M., 2008 Ecological consequences of genetic diversity. *Ecology Letters* 11:609-623.
- Hutama A. A., Hadiaty R. K., Hubert N., 2016 Biogeography of Indonesian freshwater fishes: current progress. *Treubia* 43:17-30.
- Iskandariah, Soelistyowati D. T., Gustiano R., Kusmini I. I., Huwoyon G. H., 2015 [Genetic variability of three Siamese sepat populations (*Trichopodus pectoralis* Regan; Osphronemidae) from Borneo using RAPD analysis and measurement of morphometric truss]. *Berita Biologi* 14(1):57-68. [in Indonesian]
- Islam M. N., Basak A., Ashrafullah, Ala M. S., 2011 Genetic diversity in wild and hatchery populations of stinging catfish (*Heteropneustes fossilis* Bloch) revealed by RAPD Analysis. *Journal of Biological Science* 19:81-87.
- IUCN, 2003 The IUCN Red List of Threatened Species. Version 2018-1. Available at: www.iucnredlist.org. Accessed: October, 2018.
- Jayadi, Tamsil A., Hadijah S. T., 2015 [Studies of genetic variation of Celebes rainbow (*Telmatherina ladigesii*) with method random amplified polymorphism DNA (RAPD)]. In: *Proceedings of the National Seminar on Health and Environmental Microbiology*. Hafsan, Nur F., Muthiadin C., Wahida B. F., Aziz I. R. (eds), Makassar 29 January 2015, pp. 21-27. [in Indonesian]
- Jayadi J., Husma A., Nursahran, Ardiansyah, Sriwahidah, 2016 Domestication of Celebes rainbow fish (*Marosatherina ladigesii*). *AACL Bioflux* 9(5):1067-1077.
- Jusmaldi, Duryadi D., Affandi R., Rahardjo M. F., Gustiano R., 2014 [Barcoding DNA of catfish species genus *Kryptopterus* from Sungai Mahakan Kalimantan Timur]. *Jurnal Ikhtiologi Indonesia* 14(3):191-199. [in Indonesian]
- Kottelat M., 1991 Saifin silversides (Pisces: Telamtherinidae) of Lake Matano, Sulawesi, Indonesia, with descriptions of six new species. *Ichthyological Exploration of Freshwaters* 1(4):321-344.
- Kottelat M., Whitten A. J., Kartikasari S. N., Wirjoatmodjo S., 1993 *Freshwater fishes of western Indonesia and Sulawesi*. Periplus Editions, Hong Kong, 221 pp.

- Lante S., Tenriulo A., Parenrengi A., Ravhmansyah, Malina A. C., 2011 [Genetic diversity of baronang fish populations (*Siganus guttatus*) in the Makassar Strait and Bone Bay using the Random Amplified Polymorphic DNA (RAPD) method]. *Jurnal Riset Akuakultur* 6(2):211-224. [in Indonesian]
- Lorenzen K., Malcolm C. M., Beveridge, Mengel M., 2012 Cultured fish: integrative biology and management of domestication and interaction with wild fish. *Biological Reviews* 87:639-660.
- Madden T., 2013 The NCBI handbook. 2nd edition, Bethesda (MD): National Center for Biotechnology Information (US), 478 pp.
- Mamangkey J. J., Sulistiono, Safei D. S., Soedharma D., Sukimin S., Nugroho E., 2007 [Genetic diversity of endemic Butini fish (*Glossogobius matanensis*) based on Random Amplified Polymorphism DNA (RAPD) marker in Lake Towuti South Sulawesi]. *Jurnal Riset Akuakultur* 2(3):385-393. [in Indonesian]
- Muchlisin Z. A., Thomy Z., Fadli N., Sarong M. A., Siti-Azizah M. N., 2013 DNA barcoding of freshwater fishes from Lake Laut Tawar, Aceh Province, Indonesia. *Acta Ichthyologica et Piscatoria* 43(1):21-29.
- Nasution S. H., Sulistiono, Syafei D. S., Hayani G. S., 2007 [Spatial and temporal distributions of rainbow selebensis, *Telmatheina celebensis* Boulenger, the endemic fish in lake Towuti, South Sulawesi]. *Jurnal Penelitian Perikanan Indonesia* 13(2):95-104. [in Indonesian]
- Nasution H. S., Muschsin I., Sulistiono, 2010 [Potential recruitment of bonti-bonti fish (*Paratherina striata* Aurich) in Lake Towuti, South Sulawesi]. *Bawal* 3(1):45-55. [in Indonesian]
- Nasution S. H., Sulastri S., Muchlisin Z. A., 2015 Habitat characteristics of Lake Towuti, South Sulawesi, Indonesia - the home of endemic fishes. *AACL Bioflux* 8(2):213-223.
- Nugroho E. D., Nawir D., Amin M., Lestari U., 2017 DNA barcoding of nomei fish (Synodontidae: *Harpadon* sp.) in Tarakan Island, Indonesia. *AACL Bioflux* 10(6):1466-1474.
- Nuryanto A., Pramono H., Sastranegara M. H., 2017 Molecular identification of fish larvae from East Plawangan of Segar Anakan, Cilacap, Central Java, Indonesia. *Biosaintifika* 9(1):33-40.
- Parenti L. R., 2011 Endemism and conservation of the native freshwater fish fauna of Sulawesi, Indonesia. In: *Prosiding Seminar Nasional Ikan VI & Kongres Masyarakat Iktiologi Indonesia III*. Simanjuntak C. P. H., Zahid A., Rahardjo M. F., Hadiaty K. H., Krismono, Haryono, Tjakrawidjaja A.H. (eds), Cibinong 8-9 June 2010, pp. 1-10. [in Indonesian]
- Parenti L. R., Ebach M. C., 2013 Evidence and hypothesis in biogeography. *Journal of Biogeography* 40:813-820.
- Pramono T. B., Arfiat D., Widodo M. S., Yanuhar U., 2017 [Identification of *Mystus* genus fish with genetic approach]. *Jurnal Sumberdaya Akuatik Indopasifik* 1(2):123-132. [in Indonesian]
- Prianto E., Kartamihardja E. K., Husnah, Umar C., Kasim K., Zulfia N., Budi E. K., 2014 [Policy to increase fish production and conservation of resources in the public waters of the Zone Wallacea]. Fisheries Management Research Center and Fish Resource Conservation. Technical report, 70 pp. [in Indonesian]
- Rosnaeni, Elfidasari D., Fahmi M. R., 2017 DNA barcodes of the pleco (Loricariidae, Pterygoplichthys) in the Ciliwung River. *International Journal of Advanced Research* 5(2):33-45.
- Said D. S., Hidayat, 2015 [101 Freshwater ornamental fish of the archipelago]. LIPI Press, Jakarta, pp. 121-166. [in Indonesian]
- Samuel, Husnah, Makmur S., 2009 [Capture fisheries in Lake Matano, Mahalona, and Towuti, South Sulawesi]. *Jurnal Penelitian Perikanan Indonesia* 15(1):123-131. [in Indonesian]

- Stelbrink B., Stoger I., Hardiaty R. K., Schliewen U. K., Herder F., 2014 Age estimates for an adaptive lake fish radiation, its mitochondrial introgression, and an unexpected sister group: saifin silversides of the Malili Lakes system in Sulawesi. *BMC Evolutionary Biology* 14(94):1-14.
- Sugama K., Haryanti, Cholik F., 1996 Biochemical genetics of tiger shrimp *Penaeus monodon* description electrophoresis detectable loci. *Indonesian Fisheries Research Journal* 11:19-28.
- Suwelo I. S., 2005 [The rare and threatened fish species should be protected by law]. *Jurnal Ilmu-Ilmu Perairan dan Perikanan Indonesia* 12(2):161-168. [in Indonesian]
- Tamura K., Stecher G., Peterson D., Filipski A., Kumar S., 2013 MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30:2725-2729.
- Yusron E., 2005 [Utilization of genetic diversity in the management of marine living resources]. *Oseana* 30(2):29-34. [in Indonesian]
- Umar M. T., Suwarni, Salam R., Omar S. B. A., 2012 [Study of bonti-bonti fish growth (*Paratherina striata* Aurich, 1953) in Lake Towuti, South Sulawesi]. Seminar Nasional Ikan Universitas Gajah Mada, Yogyakarta, pp. 1-12. [in Indonesian]
- Ward R. D., Zemlak T. S., Innes B. H., Last P. R., Hebert P. D. D., 2005 DNA barcoding Australia's fish species. *Philosophical Transaction of the Royal Society B* 360:1847-1857.
- Wijaya D., Samuel, Masak P. R. P., 2009 [Study of water quality and fish resource production potential in Lake Towuti, South Sulawesi]. *Bawal* 2(6):291-297. [in Indonesian]
- Wirjoatmodjo S., Sulistiono, Rahardjo M. F., Suwelo I. S., Hadiyati R. K., 2003 Ecological distribution of endemic fish species in lake Poso and Malili Complex, Sulawesi Island. Funded by Asean Regional Center for Biodiversity Conservation and the European Commission, 30 pp.

Received: 01 April 2019. Accepted: 23 June 2019. Published online: 21 August 2019.

Authors:

Jayadi Jayadi, Department of Aquaculture, Faculty of Fishery and Marine Science, Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km 05, Makassar 9023, Indonesia, e-mail: jayadi_fatrial@yahoo.com; jayadi.jayadi@umi.ac.id

Ilmiah Ilmiah, Department of Aquaculture, Faculty of Fishery and Marine Science, Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km 05, Makassar 9023, Indonesia, e-mail: ilmiah@umi.ac.id

Siti Hadijah, Department of Aquaculture, Faculty of Fishery and Marine Science, Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km 05, Makassar 9023, Indonesia, e-mail: siti.hadijah@umi.ac.id

Muhammad Kasnir, Department of Aquaculture, Faculty of Fishery and Marine Science, Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km 05, Makassar 9023, Indonesia, e-mail: Muhammad.kasnir@umi.ac.id

Dewi Indriyani Roslim, Department of Biology, Faculty of Mathematics and Nature Sciences, Universitas Riau, Pekanbaru, Indonesia, e-mail: dewiindriyaniroslim@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Jayadi J., Ilmiah I., Hadijah S., Kasnir M., Roslim D. I., 2019 DNA barcoding of Telmatherinidae family in Lake Towuti, South Sulawesi, Indonesia. *AACL Bioflux* 12(4):1208-1215.

DNA barcoding of Telmatherinidae family in Lake Towuti, South Sulawesi, Indonesia

ORIGINALITY REPORT

19%
SIMILARITY INDEX

16%
INTERNET SOURCES

14%
PUBLICATIONS

5%
STUDENT PAPERS

PRIMARY SOURCES

1 lib.unnes.ac.id 2%
Internet Source

2 e-journal.biologi.lipi.go.id 1%
Internet Source

3 lkcnhm.nus.edu.sg 1%
Internet Source

4 ore.exeter.ac.uk 1%
Internet Source

5 discovery.ucl.ac.uk 1%
Internet Source

6 Submitted to Universiti Putra Malaysia 1%
Student Paper

7 K P Bandjolu, E Madiyono, M Herjayanto, A Gani, M Nur, S Laheng, M T Gundo. "Checklist of endemic (Adrianichthyidae, Gobiidae, Zenarchopteridae) and introduced fish in Lake Poso, Sulawesi, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021 1%
Publication

8	idosi.org Internet Source	1 %
9	www.fisheriesjournal.com Internet Source	1 %
10	e-journal.unair.ac.id Internet Source	1 %
11	Nicolas Hubert, Daniel Lumbantobing, Arni Sholihah, Hadi Dahruddin et al. "Revisiting species boundaries and distribution ranges of <i>Nemacheilus</i> spp. (Cypriniformes: Nemacheilidae) and <i>Rasbora</i> spp. (Cypriniformes: Cyprinidae) in Java, Bali and Lombok through DNA barcodes: implications for conservation in a biodiversity hotspot", <i>Conservation Genetics</i> , 2019 Publication	1 %
12	journals.sagepub.com Internet Source	1 %
13	Melta Rini Fahmi, Anjang Bangun Prasetyo, Ruby Vidia Kusumah, Erma Primanita Hayuningtyas, Idil Ardi. "BARCODING DNA IKAN HIAS LAHAN GAMBUT", <i>Jurnal Riset Akuakultur</i> , 2016 Publication	1 %
14	www.neliti.com Internet Source	1 %

15	Submitted to Panjab University Student Paper	1 %
16	jxb.oxfordjournals.org Internet Source	<1 %
17	lib.dr.iastate.edu Internet Source	<1 %
18	Nursyahrhan Nursyahrhan et al.. "Food Habits and Condition Factor of the Yellow-Finned Medaka (<i>Oryzias profundicola</i>) from Lake Towuti, South Sulawesi, Indonesia.", Egyptian Journal of Aquatic Biology and Fisheries, 2022 Publication	<1 %
19	fjfsdata01prod.blob.core.windows.net Internet Source	<1 %
20	lume.ufrgs.br Internet Source	<1 %
21	Arif Wibowo, Anthony Sisco Panggabaian, Achmad Zamroni, Asep Priatna, Helman Nur Yusuf. "USING DNA BARCODE TO IMPROVE THE IDENTIFICATION OF MARINE FISH LARVAE, CASE STUDY COASTAL WATER NEAR JAKARTA AND BANDA SEA, INDONESIA", Indonesian Fisheries Research Journal, 2018 Publication	<1 %
22	ijcrbp.com Internet Source	<1 %

23

www.ncbi.nlm.nih.gov

Internet Source

<1 %

24

www.swst.org

Internet Source

<1 %

25

Iskandariah Iskandariah, Irin Iriana Kusmini, Otong Zenal Arifin, Rudhy Gustiano. "VARIASI GENETIK HASIL PERSILANGAN NILA BEST DENGAN RED NIFI DAN NIRWANA MENGGUNAKAN PENANDA RAPD", Jurnal Riset Akuakultur, 2011

Publication

<1 %

26

Meesbah Jiwaji, Rónán Daly, Kshama Pansare, Pauline McLean, Jingli Yang, Walter Kolch, Andrew R Pitt. "The Renilla luciferase gene as a reference gene for normalization of gene expression in transiently transfected cells", BMC Molecular Biology, 2010

Publication

<1 %

27

researcherslinks.com

Internet Source

<1 %

28

sabraojournal.org

Internet Source

<1 %

29

virologyj.biomedcentral.com

Internet Source

<1 %

30

www.iucnredlist.org

Internet Source

<1 %

31 Sarmini, Ketut Prasetya, Ulin Nadiroh. <1 %
"Urgency of increasing the quantity and
quality of student creativity program", Journal
of Physics: Conference Series, 2018
Publication

32 umt-ir.umt.edu.my:8080 <1 %
Internet Source

33 www.scielo.br <1 %
Internet Source

Exclude quotes On

Exclude matches < 8 words

Exclude bibliography On