

ICONETOS PAPER PUBLICATION PROGRESS

Inbox



ICONETOS Admin <iconetos@uin-malang.ac.id> Wed, Feb 10, 6:18 PM

to hanahalinurr, luthfin, riachyaoktavia, Prilya, Muhammad, Lilik, tutikimia88, me, surur.math

Dear ICONETOS presenters/authors,

On behalf of the 2nd International Conference on Engineering, Technology, and Social Sciences (ICONETOS 2020) Committee, we would like to say thank you for joining the conference and patiently waiting for the paper publication.

The editing process of your paper is already finished and all paper has been sent to the publisher. We will give you the information soon after we get an update from the publisher.

Thank you.

Best regards
ICONETOS organizing committee

ICONETOS 2020 paper progress and payment receipt

Inbox



ICONETOS Admin <iconetos@uin-malang.ac.id> Thu, Jan 28, 9:25 PM

to me

Dear St. Subaedah,

On behalf of the 2nd International Conference on Engineering, Technology, and Social Sciences (ICONETOS 2020) Committee, we would like to say thank you for joining the conference and patiently waiting for the paper publication.

The editing process of your paper is already finished and ready for publication. You can check your camera ready paper in the attachment, and you can also find the payment receipt attached to this email.

Thank you.

Best regards
ICONETOS organizing committee

Penyesuaian standar publikasi di Atlantis Press

Inbox



Imam Rofiki <imam.rofiki@uin-malang.ac.id> Sun, Nov 15, 2020, 7:03 PM

to me

Kepada
Yth. St. Subaedah,

Bersama ini kami memberitahukan bahwa paper Anda perlu perbaikan dalam penulisan referensi dan beberapa bagian untuk kutipan. Mohon disesuaikan dengan panduan penulisan/template. Paper revisian bisa dikirimkan besok.
Terima kasih atas kerjasamanya.

Hormat kami,

Panitia Iconetos

Application of Arbuscular Mycorrhizae and Compost on Growth and Yield of Soybean

St. Subaedah^{1,*}, Andi Ralle²,

^{1,2)} Departement Agrotechnology Universitas Muslim Indonesia
Urip Sumoharjo Road No. 226, Makassar 90231, Indonesia

* Corresponding author's e-mail: st.subaedah@umi.ac.id

ABSTRACT

National soybean demand continues to increase in line with the increase in population, while soybean production is still very low. The low production is caused by several factors and one of them is because the development is faced with lands with low soil fertility. Therefore, it is necessary to improve soil fertility, including the use of biological fertilizers such as arbuscular mycorrhizae along with the provision of compost. This research was conducted with the aim to determine the effect of arbuscular mycorrhizal applications and compost on the growth and production of soybean plants. This research was prepared based on a randomized block design consisting of 5 treatments, namely: control (without mycorrhizal and without compost), mycorrhizal application, mycorrhizal + compost 5 g/plant, mycorrhizae + compost 10 g/plant, mycorrhizae + compost 15 g/plant. After the treatment was repeated three times, so there were 15 experimental units. The results showed that mycorrhizal application accompanied by compost resulted in higher growth and yield of soybean than without mycorrhizal application and mycorrhizal application without compost.

Keywords: *Arbuscular mycorrhizae, compost, soybean, growth, yield*

1. INTRODUCTION

Soybean (*Glycine max* (L.) Merrill) is one commodity crops are urgently needed in Indonesia, because it can be consumed in a variety of processed food products such as tofu, tempeh, milk, and many other processed products. In addition to soy it also has a relatively high nutrient content, among others, 35 g protein, 53 g carbohydrate 18 g fat and 8 g of water in 100 g of food, even for specific high yielding varieties, protein content is 40-43 g [1].

Efforts to increase the availability of soybeans have become one of the targets of national development in agriculture. This is because the need for soybeans continues to increase in line with the increase in population. In 2018, soybean demand reached 2.5 million tons, while soybean production only reached 82.598 tons [2], so that soybean imports can not be avoided. Soybean development efforts are faced with low soil fertility due to erosion and land management that does not pay attention to its sustainability aspects [3]. Therefore, efforts are needed to improve land management that can support the sustainability of soybean crop production. One effort that can be done is the use of biological fertilizers, such as arbuscular mycorrhizae. Arbuscular mycorrhiza is a mutually beneficial symbiosis between fungi and plant roots [4].

Arbuscular mycorrhizae infect the root system of the host plant which will intensively produce hyphae interwoven, so that mycorrhizal plants will be able to increase nutrient and water absorption capacity [5; 6], and increase plant resistance to soil salinity stress [7].

The effectiveness of mycorrhizae in increasing nutrient availability to improve plant growth depends on the type of mycorrhizae itself, the type of host plant, and also very much depends on soil types, especially soil pH and fertility levels and their interactions [8]. Furthermore, [9] suggests that the activity of microorganisms in the soil is influenced, among others, by the level of organic matter in the soil, because organic matter is used by microorganisms as their energy source. To increase the level of soil organic matter can be reached by using organic fertilizers, such as compost. Based on this description, a research was conducted on the effect of mycorrhizal and compost applications on the growth and yield of soybean plants.

2. RESEARCH OF METHODS

This research was conducted in an experimental form at the Green House of the Faculty of Agriculture and Soil Laboratory of the Universitas Muslim Indonesia, Makassar, South Sulawesi, Indonesia. This research took place from April to August 2020.

The tools used in this study were polybags, shovels, analytical scales, hand sprayer, plant labels, oven, sieve, bucket, tape measure, camera and writing instruments. While the materials used in this study were soil, sand, compost, soybean seeds, mycorrhizal, distilled water and alcohol.

This research was prepared based on a randomized block design (RBD) which consisted of 5 treatments, namely:

M0: Control (without mycorrhizal and without compost)

M1: Mycorrhiza

M2: Mycorrhizae + compost 5 g / plant

M3: Mycorrhizae + compost 10 g / plant

M4: Mycorrhizae + compost 15 g / plant

Each treatment was repeated three times in order to obtain 15 experimental units.

2.1. Trial Implementation

This research was conducted in several stages, namely the preparation of planting media, planting and mycorrhizal inoculation, maintenance and observation. Each stage will be described as follows:

2.1.1. Preparation of Planting Media

The planting medium used was soil and sand with a ratio of 2: 1 (v / v). The media is then weighed as much as 8 kg / polybag, before being put into a poly bag, for treatment of M2, M3 and M4 the planting media is mixed with compost according to the specified treatment dose. After all the planting media has been put in a poly bag, it is then arranged according to the predetermined research plan.

2.1.2. Seed Planting and Mycorrhizal Inoculation

Soybean seeds are planted in the planting medium that has been provided. One polybag is planted with 3 soybean seeds, with the aim of anticipating that if one seed dies, there are still two living seeds. Furthermore, when the plants were two weeks old, 10 g / plant of mycorrhizae were administered.

2.1.3. Maintenance

The maintenance stage is carried out for approximately 3 months. Soybean plants are placed in a place that gets enough sunlight. Watering is carried out according to soil moisture conditions. NPK fertilization with a low dose of 150 kg / ha.

The variables observed in this study were: plant height, number of leaves, number of branches, number of pods per plant, weight of pods per plant and weight of seeds per plant.

Data from observations in the field were analyzed using the randomized block design method and then if the treatment had a significant effect, then it was followed by the LSD test at the α 0.05 level.

3. RESULTS AND DISCUSSION

3.1. Results

3.1.1. Plant height

The observation result of soybean plant height at 8 weeks after planting (WAP) showed that the application of mycorrhiza and compost did not significantly affect plant height. The average height of soybean plants at the age of 8 MST which is presented in Figure 1 shows that the treatment of mycorrhizal applications (M1) without compost tends to obtain higher soybean plants, namely 42.00 cm compared to other treatments.




Figure 1. Effect of Arbuscular Mycorrhizae and Compost Application on Soybean Plant Height at Age 8 WAP

3.1.2. Number of Leaves

Mycorrhizal and compost applications showed a significant effect on the number of soybean leaves at the age of 8 WAP. The results of the LSD further test in Table 1 show that the application of mycorrhizae with the addition of 5 g of compost (M2) obtained a higher number of leaves, namely 28.67 strands and significantly different from other treatments (M0, M3 and M4) but not different from the treatment of mycorrhizal applications without compost (M1).

Table 1. The Effect of Mycorrhizal and Compost Applications on the Number of Soybean Plant Leaves at 8 WAP

Treatment	Number of Leaves	LSD α 0.05
M0: Control	13.33 b	8.52
M1: Mycorrhizae	21.00 ab	
M2: M+ 5g C	28.67 a	
M3: M+10g C	20.00 b	
M4: M+15g C	16.00 b	

Note: The values followed by the same letters are not significantly different based on the LSD level of α 0.05

3.1.3. Number of Branches

The results of data analysis on the number of branches showed that the application of mycorrhizal and compost did not significantly affect the number of soybean branches. However, the results of the average value of the number of branches presented in Figure 2 indicate that the treatment of mycorrhizal administration with the addition of 5 g of compost (M2) tended to obtain more branches, namely 11 branches compared to other treatments.




Figure 2. Effect of Mycorrhizal and Compost Applications on the Number of Branches

3.1.4. Number of Pods

Observation of the number of pods showed that the application of mycorrhizal and compost significantly affected the number of soybean pods produced. The average number of soybean pods per plant is presented in Table 2 shows that the highest number of pods was obtained in the mycorrhizal application treatment with the addition of 10 g (M3) compost, namely 57.67 pods followed by mycorrhizal application treatment with the addition of 5 g of compost (M2). and significantly different from the number of soybean pods produced in the control treatment, namely without mycorrhizal and without compost (M0) but not significantly different from other treatments (M1, M3 and M4).

Table 2. Average Number of Soybean Pods with Mycorrhizal and Compost Applications

Treatment	Number of Soybean Pods	LSD α 0.05
M0: Control	36.67 b	12.04
M1: Mycorrhizae	46.00 ab	
M2: M+ 5g C	57.00 a	

M3: M+10g C	57.67 a
M4: M+15g C	51.67 a

Note: The values followed by the same letters are not significantly different based on the LSD level of α 0.05

3.1.5. Pod Weight

The results of data analysis on soybean pod weight per plant showed that the application of mycorrhizal and compost had a significant effect. In Table 3, the average weight of soybean pods per plant is presented and shows that the application of mycorrhizal treatment and the addition of 5 g of compost per plant (M2) obtained the heaviest weight of soybean pods per plant, namely 52.28 g and significantly different from other treatments (M0, M1 and M4) but not significantly different from the application of mycorrhizal treatment and the addition of 10 g of compost (M3).

Table 3. Average pod weight per plant (g) with mycorrhizal and compost application

Treatment	Pod Weight (g)	LSD α 0.05
M0: Control	19.89	13,85
M1: Mycorrhizae	31.99	
M2: M+ 5g C	52.28	
M3: M+10g C	43.22	
M4: M+15g C	34.20	

Note: The values followed by the same letters are not significantly different based on the LSD level of α 0.05

3.1.6. Seed Weight Per Plant

The results of observations on the weight of soybean seeds per plant showed that the application of mycorrhizae and compost had a significant effect. The results of the LSD follow-up test in Table 4 show that the heaviest weight of soybean seeds per plant was obtained in the application of mycorrhizal application of 5 g of compost per plant (M2), which was 28.04 g and was significantly different from the control treatment (without mycorrhizal and without compost) and also significantly different from the application of mycorrhizal treatment without compost (M1).

Table 4. Average weight of soybean seeds (g) per plant using mycorrhizal and compost applications

Treatment	Seed Weight (g)	LSD α 0,05
M0: Control	13.35 c	9.19
M1: Mycorrhizae	19.23 bc	
M2: M+ 5g C	28.04 a	
M3: M+10g C	23.04 ab	
M4: M+15g C	23.63 ab	

Note: The values followed by the same letters are not significantly different based on the LSD level of α 0.05

3.2. Discussion

The results of the data analysis showed that the application of mycorrhizae and compost had a significant effect on the growth and production of soybean as indicated by the parameters of the number of leaves, number of branches, pod weight and seed weight per plant. The results of this study are in line with the findings of [10], who found that soybeans given mycorrhizal treatment affected growth, yield components and

soybean yield. Likewise, the results of [11] study found that the application of mycorrhizae resulted in better soybean plant growth.

The good effect of mycorrhizal applications on the growth and production of soybean plants is due to the potential for mycorrhizae to infect the root system of the host plant and will produce hyphae intensively so that mycorrhizal plants will be able to increase their capacity to absorb nutrients and water [5]. Several studies have shown that mycorrhizae can improve plant growth and nutrient uptake [12; 13; 14]. From this study it was also found that mycorrhizal applications increased the availability of phosphorus by 34% compared to those without mycorrhizal applications (without mycorrhizal application, the availability of phosphorus was 4.03 ppm, and increased to 5.41 ppm with mycorrhizal applications. Increased availability of phosphorus is very significant for plant growth, because Phosphorus is one of the important macro nutrients in plant growth and development; it functions as a constituent of metabolites in complex compounds, as an activator, cofactor or fiber enzyme unifier and plays a role in physiological processes and is also a structural component. Of a number of important compounds, the energy transfer molecules ADP and ATP [15].

The results of this study also showed that mycorrhizal application treatment accompanied by the application of compost as much as 5 g per plant showed a higher production as indicated by the pod weight per plant and seed weight per plant compared to without compost and compost application with a higher dose. This is due to the mycorrhizal effectiveness, among others, influenced by soil fertility, particularly soil P content. The application of compost can increase the soil organic matter content which in turn will increase the movement and availability of P nutrients in the soil, so that with more compost doses it allows the availability of more P nutrients and results in decreased effectiveness of mycorrhizae. This is consistent with the findings of [16] who stated that mycorrhizal associations tend to decrease when the P concentration in plants increases. Higher P availability can reduce spore production [17] and reduce secondary external hyphae [18]. Therefore, increasing the P status of the roots would reduce hyphae branching and mycorrhizal associations.

4. CONCLUSION

The application of mycorrhizae accompanied by 5 g of compost per plant obtained the highest number of leaves and pods, as well as the heaviest pod weight and seed weight.

ACKNOWLEDGMENTS

Thanks to the Ministry of Research and Technology and Higher Education (RISTEK-DIKTI) of the Republic of Indonesia for providing financial assistance, and Universitas Muslim Indonesia so that this research can be carried out.

ACKNOWLEDGMENTS

REFERENCES

- [1]. Suprpto. 2001. Bertanam Kedelai. Penebar Swadaya. Jakarta. 74p.
- [2]. CBS, 2019. Indonesian Statistics. <http://www.bps.go.id>
- [3]. Subaedah, St. 2018. Agroteknologi Lahan Kering. Nas Media Pustaka, Makassar . 200p
- [4]. Smith, S.E., F.A. Smith. 2012. Fresh perspectives on the roles of arbuscular mycorrhizal fungi in plant nutrition and growth. Mycologia, 104(1): 1–13
- [5]. Talanca, A.H. 2010. Status cendawan Mikoriza Vesicular-Arbuskular (MVA) pada tanaman. In Prosiding Pekan Serelia Nasional. 3:353-357
- [6]. Buechel, T. and E. Bloodnick, 2016. Mycorrhizae: description of types, benefits and uses. Plant Health. <https://gpnmag.com/article/mycorrhizae-description-of-types-benefits-and-uses>

- [7]. Damaiyanti, D.R.R., N.Aini, R. Soelistyono. 2015. Effects of arbuscular mycorrhiza inoculation on growth and yield of tomato (*Lycopersicum esculentum* Mill) under salinity stress. J. Of Degraded and Mining Lands Management, 3 (1):447-452
- [8]. Bruce, A., Smith, W. E. and Tester, M. 1994. The development of mycorrhizal infection in cucumber: effects of P supply on root growth, formation of entry points and growth of infection units. New Phytol. 127: 507-514.
- [9]. Sutanto, R. 2005. Pertanian Organik. Kanisius. Yogyakarta. 219 p.
- [10]. Haryoko, W., dan Y.M.Zen. 2003. Pengaruh Flavonoid dan Cendawan Mikoriza Arbuskula Terhadap Pertumbuhan dan Hasil Kedelai Pada Ultisol. J. Stigma. 4: 352-356
- [11]. Turmuktini, T. 2009. Interaksi antara dosis fungi mikoriza arbuskula terhadap pertumbuhan, kuantitas dan kualitas tiga kultivar kedelai. Jurnal Hayati, 3:79-83
- [12]. Graham, J. H. dan D. M. Eissenstat. 1998. Field evidence for the carbon cost of citrus mycorrhizas. New Phytology 140: 103-110
- [13]. Fidelibus, M. W., C. A. Martin, G. C. Wright dan J. C. Stutz. 2000. Effect of arbuscular mycorrhizal (AM) fungal communities on growth of Volkamer lemon in continually moist or periodically dry soil. Scientia Horticulturae 84: 127-140.
- [14]. Subaedah, S., S.S. Netty, A. Ralle. 2020. Growth and Yield of Two Soybean Varieties by Phosphate Fertilization and Arbuscular Mycorrhizal Application. J. Biological of Sciences. 20 (4): 147-152

ICONETOS 2020 PAPER NOTIFICATION

External

Inbox



ICONETOS Admin <iconetos@uin-malang.ac.id> Oct 24, 2020, 8:43 PM

to me

Dear Participant,

We are pleased to inform you that your paper submitted to The 2nd International Conferences on Engineering, Technology and Social Science (ICONETOS 2020) is **ACCEPTED** for presentation. Attached you will find the LoA and the review comments. As for the publication, there will be another review round after the conference. So Please keep up the update even after the conference is over.

It is mandatory to prepare the camera-ready paper as per the instructions listed on <http://iconetos.uin-malang.ac.id/> and your paper will not be published unless the following are done:

1. Fill in the registration form through this link <https://bit.ly/3jj773N> before October 26th, 2020
(For those who have already filled the form, you don't have to do it again). You will receive the program book and the conference link after filling the form.
2. The oral presenters have to make a presentation video (can be converted from a ppt file), while the poster presenters have to make a poster for presentation. You may choose one of them. The video length must not be more than 10 minutes. The poster template layout will be available on the conference website. Please submit it before October 29th, 2020. The video/poster submission is in this link: <https://bit.ly/2IYv57v>
3. Format your paper as per guidelines and Submit your full paper to the link (in word document format) before November 6th, 2020. Please submit the revised paper in this link: <https://bit.ly/37zHWqM>
4. The accepted similarity level is a maximum of 25%
5. Make a payment (for the presenters from outside UIN Malang)

Conference fees:

Indonesian: IDR 250.000

Outside Indonesia: USD 50

The Publication fees can be paid later after the revised paper is submitted.

Publication Fees :

Indonesian: IDR 1.300.000

Outside Indonesia: USD 100

Bank Transfer to:

Bank: BRI (Bank Rakyat Indonesia)

Account Name: RPL 032 UIN UNTUK DANA KELOLAAN

Account Number: 0344-01-000456-30-6

Swift Code: BRINIDJA

Bank Code: 002

City : Malang

Country: Indonesia

Via PayPal:

<https://www.paypal.me/haridasamudro>

6. The conference will be held online on October 31st, 2020. The conference link will be sent to you after the registration form is completed.

All related conference materials can be found at <http://iconetos.uin-malang.ac.id/>

Please let us know if you have any questions regarding the registration process.

With a warmest regard,

-On behalf of the ICONETOS 2020 Committee

(TPC Chair of ICONETOS 2020)

2 Attachments



International Conference on Engineering, Technology, and Social Sciences

MANUSCRIPT REVIEWING KITS

PAPER ID: Application of Arbuscular Mycorrhizae and Compost on Growth and Yield of Soybean					
PLAGIARISM CHECK					22%
RATING (Referee to rate the article) 1=poor, 2=fair, 3=good, 4=excellent					
NO	Criteria	1	2	3	4
1	Clarity of formal structure			√	
2	Suitability of title to the contents			√	√
3	Abstract sufficiently informative			√	√
4	Originality/Novelty			√	√
5	Methodology sufficiently described			√	
6	Methodology appropriate to study			√	
7	Result/conclusions support by data analysis			√	
8	Clarity in the presentation of findings			√	
9	Language			√	
10	Bibliography adequate and relevant			√	
SPECIFIC COMMENTS BY THE REFEREE					
The reviewer should identify and comment on major strengths and weakness of the study design and methodology and provide the author(s) with useful suggestions for improvement of the manuscript. Please provide your comments in the box below.					
1. Please add literature about interaction compost and mycorizal 2. The reference style we use is IEEE style (numbering). Please follow the template and adjust it 3. Every table and figure must be explained in the text					
RECOMMENDATION					
OPTIONS					DECISION
ACCEPT AS IS <i>with or without only minor changes to be made by editorial staff</i>					
ACCEPT WITH MINOR REVISIONS <i>With only minor changes to be made by the author</i>					√
RETURN TO AUTHOR(S) FOR IMPORTANT MODIFICATIONS <i>Author to revise & resubmit for another round of reviews</i>					
REJECT					



International Conference on Engineering, Technology, and Social Sciences
Research and Community Development Center
Universitas Islam Negeri Maulana Malik Ibrahim Malang
Jalan Gajayana 50 Malang Indonesia
Iconetos.uin-malang.ac.id

ACCEPTANCE LETTER

Dear **St. Subaedah**,

We are pleased to inform you that your paper:

Paper ID : **90**
Title : **Application of Arbuscular Mycorrhizae and Compost on Growth and Yield of Soybean**
Author(s) : **St. Subaedah, Andi Ralle, Suraedah Alimuddin**
Departement Agrotechnology Universitas Muslim Indonesia

Submitted to The 2nd International Conferences on Engineering, Technology and Social Science (ICONETOS 2020) is **ACCEPTED** for presentation. As for the publication, there will be another review round after the conference. So Please keep up the update even after the conference is over.

The conference will be held online on October 31st, 2020. The link will be sent to you after the registration form is completed. Thank you for joining the conference.

Best regards,



Prof. Dr. Tutik Hamidah, M.Ag
ICONETOS Chair