

Type of Delivery and Length of Stay in Hypertension Patients

by Dr. Dr. Nasrudin Sp.og(k).,mars.

Submission date: 18-Jun-2020 04:29PM (UTC+0700)

Submission ID: 1345873655

File name: Type_of_Delivery_and_Length_of_Stay_in_Hypertension_Patients.pdf (171.99K)

Word count: 3511

Character count: 19112

Type of Delivery and Length of Stay in Hypertension Patients

¹ Syahriana^{1,2}, Nasrudin Andi Mappaware^{1,3}, Yusrawati Hasibuan⁴, Elizawarda^{4,5} & Andi Nilawati Usman^{6,7}

¹ Postgraduate Program, Department of Midwifery, Hasanuddin University, Makassar, South Sulawesi, Indonesia

² Regional Public Hospital of Makassar, Makassar, South Sulawesi, Indonesia

³ Faculty of Medicine, Indonesia Muslim University, Makassar, South Sulawesi, Indonesia

⁴ Department of Midwifery, Medan Polytechnic of Health Sciences, Medan, North Sumatera, Indonesia

⁵ Human Resource Education and Training Unit, Department of Midwifery, Medan Polytechnic of Health Sciences, Medan, North Sumatera, Indonesia

⁶ Hallal Center for Public Health Research, Faculty of Public Health, Hasanuddin University, Makassar, South Sulawesi, Indonesia

⁷ Faculty of Public Health, Mandala Waluya College, Kendari, Southeast Sulawesi, Indonesia

Correspondence: Syahriana, Postgraduate Program, Department of Midwifery, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia; Tel: 62-821-9166-4594. E-mail: syahriana789@gmail.com

Received: April 23, 2018 Accepted: May 11, 2018 Online Published: May 16, 2018

doi:10.5539/gjhs.v10n6p158

URL: <https://doi.org/10.5539/gjhs.v10n6p158>

Abstract

Introduction: The length of stay (LOS) for pregnant mothers with vaginal delivery is different from that of caesarean delivery. Hypertension during pregnancy may be an indication to conduct caesarean delivery that affects length of stay in hospitals.

Material and Methods: The total collected samples were 108 pregnant mothers and they were divided into 2 groups, 54 pregnant mothers with normal blood pressure and 54 pregnant mothers with hypertension. Maternal characteristics, routine hematology, blood pressure (BP) status, type of delivery and length of stay were analyzed.

Results: Type of delivery, maternal age, height, weight, body mass index (BMI), systolic blood pressure, diastolic blood pressure, length of stay, and routine hematologic profile except leukocytes showed statistically significant correlations with hypertension for the total 108 pregnant mothers. Pregnant mothers with caesarean delivery due to pregnancy induced hypertension (PIH) was threefold than those of normal blood pressure (95% confidence interval, 1.343-6.563). Statistical analyses indicated significant correlations between type of delivery and length of stay for pregnant mothers with hypertension after delivery (p -value = 0.000).

Conclusion: On average, pregnant mothers who had hypertension with caesarean delivery spent at least 49-72 hours in the hospital after delivery. Indeed, the Royal College of Obstetricians and Gynecologists recommended that pregnant mothers with vaginal delivery should stay in a hospital for 1 or 2 days and those with caesarean section were suggested to stay 3 or 4 days after delivery. However, some pregnant mothers in this study spent shorter time to stay at the hospital after delivery. Therefore, it is suggested for pregnant mothers to stay at a hospital in a more sufficient period of time after delivery to give an adequate time for physicians to classify, to diagnose, or to treat complications after delivery.

Keywords: pregnancy induced hypertension, type of delivery, length of stay

1. Introduction

Pregnancy induced hypertension is a complication that contributes significantly to maternal and perinatal morbidity and mortality (Uzan et al., 2011; MacDonald-Wallis et al., 2012; Kazemian et al., 2014; Kattah & Garovic, 2014; McCarthy et al., 2015). Pregnancy induced hypertension, preeclampsia and complications occur at gestational age more than 20 weeks with maternal blood pressure at $\geq 140 / 90$ mmHg (American College of Obstetricians and Gynecologists, 2013; Kattah & Garovic, 2014; Melamed et al., 2014; Magee et al., 2016; Zhang et al., 2017). Preeclampsia and eclampsia contribute to maternal death for every 3 minutes worldwide (Mammaro et al., 2009; Meshram et al., 2014). Pregnancy induced hypertension (PIH) is one of the most common pregnancy complications (Durst et al., 2016) that calls for consideration to conduct a certain type of delivery

(Haroush et al., 2005). Fetal distress caused by preeclampsia and eclampsia requires doctors physicians to perform cesarean section (Alanis et al., 2008). Length of stay after delivery at a health facility is vary among pregnant mothers (Campbell et al., 2016). Some pregnant mothers spend either shorter time or longer time compared with the actual needs. Specifically, length of stay (LOS) in a shorter period of time leads to insufficient time for physicians to detect, to diagnose, or treat complications of delivery that will in turn increase the possibility of morbidity and mortality (MD et al., 2016; Plough et al., 2017). Length of stay at a hospital for pregnant mothers with vaginal delivery is different from that of caesarean section (SC), hypertension in pregnancy may be an indication to conduct caesarean section that affects length of stay in hospitals. Therefore, this study aims to determine correlations between type of delivery and length of stay in pregnant mothers with PIH and to compare length of stay in the hospital between pregnant mothers with vaginal delivery and those with caesarean section with PIH.

2. Materials and Methods

2.1 Study Design and Setting

A prospective cross-sectional study was conducted at the Sitti Khadijah I Mother and Child Hospital Makassar, South Sulawesi, Indonesia during June until December 2017. The total 108 samples of pregnant mothers were collected in the study which divided into two groups, 54 pregnant mothers who gave birth with hypertension and 54 pregnant mothers who gave birth with normal tension. Purposive sampling technique was used in accordance with the inclusion and exclusion criteria.

2.2 Variables and Data Collection

Data of samples were screened by medical officers for enrollment with maternal inclusion criteria both with hypertension and normal tension together with independent variables that include age, duration of stay, type of delivery, systolic blood pressure, diastolic blood pressure, body mass index (BMI), and routine hematology. The exclusion criteria in this study were mothers who had multiple gestation, history of heart disease, diabetes and renal disease.

2.3 Ethical Considerations

This study received ethical clearance from the Research and Ethics Committee, Faculty of Medicine of Hasanuddin University and was registered in 1073/H4.8.4.5.31/PP36-KOMETIK/2017.

2.4 Data Processing and Analysis

All collected data were encoded, and then they were entered and analyzed. We performed the statistical analyses by comparing maternal characteristics with blood pressure status which presented as mean $\pm$ standard deviation and frequencies for categorical variables, correlations between type of delivery with blood pressure status, and correlations between type of delivery and length of stay in hypertension patients. The variables were compared by using chi-square test, T-test and Mann-Whitney test.

3. Results

3.1 Socio-Demographic, Maternal Anthropometric Profile, and Routine Hematologic

In this study, maternal characteristic data with the total 108 mothers showed that type of delivery, maternal age, height, weight, BMI, systolic blood pressure (BP), diastolic blood pressure (BP), length of stay, and routine hematologic profile except leukocytes showed statistically significant correlations with maternal hypertension. Level of education, employment status, gravidity and gestational age showed no statistically significant differences between the two groups.

Table 1. Comparison of socio-demographic and anthropometric profile

Variable	Normal Tension Group (n=54)	PIH Group (n = 54)	p- value
Level of Education, n (%)			
High	25	30	0.441
Low	29	24	
Status of Employment, n (%)			
Employed	29	26	0.700
Unemployed	25	28	
Gravidity, n (%)			
Primiparous	20	22	0.844
Multiparous	34	32	
Age (mean ± SD)	28.87(1.2)	32.04(5.7)	0.007
Height (mean ± SD)	153.87(5.9)	156.31(5.6)	0.030
Weight (mean ± SD)	61.89(1.5)	73.52(1.4)	0.000
BMI (mean ± SD)	26.22(3.7)	29.97(5.2)	0.000
GA (mean ± SD)	38.44(1.7)	38.06 (2.0)	0.088
Systolic BP (mean ± SD)	114.81(9.1)	156.48 (18.5)	0.000
Diastolic BP (mean ± SD)	74.48 (6.5)	101.67 (11.3)	0.000
LOS (mean ± SD)	1.94 (1.0)	2.44 (1.0)	0.019
Routine Hematology (mean ± SD)			
Hemoglobin	10.95 (1.5)	11.51 (1.4)	0.050
Leukocytes	10.73 (2.6)	11.65 (3.5)	0.124
Erythrocytes	4.13 (0.9)	4.35 (0.5)	0.000
Hematocrit	31.58 (4.0)	35.55(4.4)	0.000
Platelets	294.41(67.3)	247.46 (64.2)	0.000
Glucose	95.81 (14.1)	103.81 (17.3)	0.007

3.2 Type Of Delivery

Table 2 shows that hypertension significantly affects the type of delivery (p-value=0.011). Pregnant mothers with caesarean delivery due to pregnancy induced hypertension (PIH) was threefold than those of normal blood pressure (95% confidence interval, 1.343-6.563).

Table 2. Types of delivery with different blood pressures

Delivery	Normal Tension Group (n=54)	PIH Group (n = 54)	OR (95% CI)	p- value
Vaginal	30	16	2.969	0.011
Caesarean	24	38	(1.343-6.563)	

3.3 Length of Stay: Correlation And Association

The percentage of normal blood pressure with vaginal delivery was higher than that of caesarean delivery of 55.6% with LOS ranging from 24-47 hours (56.7%) for vaginal and 48-72 hours (54.2%) for caesarean. The percentage of

pregnant mothers who had hypertension with cesarean delivery was higher than that of vaginal delivery (70.4%) with the length of stay (LOS) ranging from 48 to 72 hours (50%), whereas normal delivery was <24 hours (50%). These data showed that there were correlations between type of delivery and blood pressure status with length of stay in the hospital.

Table 3. Correlations between types of delivery and lengths of stay by four categories

Category	Percentage woman by category	Length Of Stay				P Value
		< 24 h	24-47 h	48-72 h	> 72 h	
		Count (%)	Count (%)	Count (%)	Count (%)	
Normal Tension + Vaginal Delivery	55.6%	6 (20%)	17 (56.7%)	6 (20%)	1 (3.3%)	0.000
Normal Tension + Secarean	44.4%	0 (0%)	2 (8.3%)	13 (54.2%)	9 (37.5%)	
PIH + Vaginal Delivery	29.6%	8 (50%)	5 (3.2%)	3 (18.8%)	0 (0%)	0.000
PIH + secarean	70.4%	0 (0%)	13 (34.2%)	19 (50%)	6 (15.8%)	

4. Discucion

Our findings indicated that the average length of stay in the hospital for mothers who had normal blood pressure with vaginal delivery was 24-47 h. This is considerably longer than those with hypertension who have shorter treatment (<24 h) with a proportion of 55.6% versus 29.6%. However, length of stay was statistically predicted longer for pregnant mothers who underwent caesarean section (NICE, 2011). Referring to risk factors, there are differences in terms of age, height, weight, body mass index of pregnant mothers. Blood pressure status (diastolic blood pressure and systolic blood pressure) also affect both groups and routine hematology profiles were different for both groups except leukocytes. Pregnant women with hypertension had the average age of 32.02 years old with the body mass index of 29.97, whereas pregnant women with normal blood pressure had the average age of 28.87 years old with the body mass index of 26.22. Overweight or obesity is a risk factor for the occurrence of preeclampsia (Sohlberg et al., 2009; Knight, Kurinczuk, & Spark, 2010; Kazemian et al., 2014) and reveals an ascending pattern with the increase of BMI (Reyes et al., 2012; Bilano et al., 2014; Gudnadóttir et al., 2016; Li et al., 2016). In terms of maternal age, mothers within the range of 30-34 years old are 2 times more likely experience preeclampsia (Choudhury et al., 2014; Chan et al., 2015). Our findings showed that women with PIH treated by caesarean delivery up to threefold (Table 2). In this study, most pregnant mothers with hypertension had caesarean delivery with the length of stay at 48-72 hours. This is in line with pregnant mothers who have preeclampsia and eclampsia in which they should be treated with caesarean delivery (NICE, 2010; NICE, 2011; Mylonas and Friese, 2015). with the length of stay is 48-72 hours (Moroy et al., 2007), as recommended by the Royal College of Obstetricians and Gynaecologists that pregnant mothers with vaginal delivery should spend 1 or 2 days at hospitals and those with caesarean section ought to spend 3 or 4 days at hospitals (NICE, 2011). Although some pregnant women in this study with hypertension treated by vaginal delivery had shorter time to stay in hospitals with the average of length of stay was <24 h, this could lead to potential hemorrhage (WHO, 2013), since the first 24 hours of postpartum is the highest risk period for the mother and newborn. This study bear some limitations that include samples were collected in small number.

5. Conclusion

This study showed that women with PIH mostly gave birth to caesarean section with the average length of stay was 48-72 hours. This is in accordance with the recommendations of NICE that pregnant mothers with hypertension complications such as preeclampsia and eclampsia should be treated with caesarean section. Although pregnant mothers with caesarean section should stay for 48-72 hours after delivery as stated in the recommendations, some pregnant mothers in this study spent shorter time to stay at the hospital after delivery that could lead hemorrhage. Therefore, it is suggested for pregnant mothers to stay at a hospital in a more sufficient period of time after delivery to give an adequate time for physicians to classify, to diagnose, or to treat complications.

Acknowledgements

The authors express our sincere thanks to all the participants in this study. The authors also thanks director and staff of the Sitti Khadijah I Makassar Mother and Child Hospital for their support and assistance.

Competing Interests Statement

The authors declare that there are no competing or potential conflicts of interest.

References

- Alanis, M. C., Robinson, C. J., Hulsey, T. C., Ebeling, M., & Johnson, D. D. (2008). Early-onset severe preeclampsia: induction of labor vs elective cesarean delivery and neonatal outcomes. *American Journal of Obstetrics & Gynecology*, 199(3), 262-e1. <https://doi.org/10.1016/j.ajog.2008.06.076>
- American College of Obstetricians and Gynecologists. (2013). Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' task force on hypertension in pregnancy. *Obstetrics and gynecology*, 122(5), 1122. <https://doi.org/10.1097/01.AOG.0000437382.03963.88>
- Ben - Haroush, A., Yogeve, Y., Glickman, H., Kaplan, B., Hod, M., & Bar, J. (2005). Mode of delivery in pregnant women with hypertensive disorders and unfavorable cervix following induction of labor with vaginal application of prostaglandin E2. *Acta obstetrica et gynecologica Scandinavica*, 84(7), 665-671. <https://doi.org/10.1111/j.0001-6349.2005.00681.x>
- Campbell, O. M., Cegolon, L., Macleod, D., & Benova, L. (2016). Length of stay after childbirth in 92 countries and associated factors in 30 low-and middle-income countries: compilation of reported data and a cross-sectional analysis from nationally representative surveys. *PLoS medicine*, 13(3), e1001972. <https://doi.org/10.1371/journal.pmed.1001972>
- Chan, T. F., Tung, Y. C., Wang, S. H., Lee, C. H., Lin, C. L., & Lu, P. Y. (2015). Trends in the incidence of pre-eclampsia and eclampsia in Taiwan between 1998 and 2010. *Taiwanese Journal of Obstetrics and Gynecology*, 54(3), 270-274. <https://doi.org/10.1016/j.tjog.2013.06.021>
- Choudhury, K. N., Mainuddin, A. K. M., Wahiduzzaman, M., & Islam, S. M. S. (2014). Serum lipid profile and its association with hypertension in Bangladesh. *Vascular health and risk management*, 10, 327. <https://doi.org/10.2147/VHRM.S61019>
- Durst, J. K., Subramaniam, A., Tang, Y., Szychowski, J. M., Campbell, S. B., Biggio, J. R., & Harper, L. M. (2017). Mode of delivery in nulliparous women with gestational hypertension undergoing early term induction of labor. *The Journal of Maternal-Fetal & Neonatal Medicine*, 30(19), 2291-2296. <https://doi.org/10.1080/14767058.2016.1247153>
- Gudnadóttir, T. A., Bateman, B. T., Hernández-Díaz, S., Luque-Fernandez, M. A., Valdimarsdóttir, U., & Zoega, H. (2016). Body mass index, smoking and hypertensive disorders during pregnancy: a population based case-control study. *PloS one*, 11(3), e0152187. <https://doi.org/10.1371/journal.pone.0152187>
- Kattah, A. G., & Garovic, V. D. (2013). The management of hypertension in pregnancy. *Advances in chronic kidney disease*, 20(3), 229-239. <https://doi.org/10.1053/j.ackd.2013.01.014>
- Kazemian, E., Sotoudeh, G., Dorosty-Motlagh, A. R., Eshraghian, M. R., & Bagheri, M. (2014). Maternal obesity and energy intake as risk factors of pregnancy-induced hypertension among Iranian women. *Journal of health, population, and nutrition*, 32(3), 486. <https://doi.org/10.4172/2167-0420.1000116>
- Knight, M., Kurinczuk, J. J., Spark, P., & Brocklehurst, P. (2010). Extreme obesity in pregnancy in the United Kingdom. *Obstetrics & Gynecology*, 115(5), 989-997. <https://doi.org/10.1097/AOG.0b013e3181da8f09>
- Li, K., Yao, C., Yang, X., Di, X., Li, N., Dong, L., ... & Zheng, M. (2016). Body mass index and the risk of cardiovascular and all-cause mortality among patients with hypertension: A population-based prospective cohort study among adults in Beijing, China. *Journal of epidemiology*, 26(12), 654-660. <https://doi.org/10.2188/jea.JE20150323>
- Macdonald-Wallis, C., Lawlor, D. A., Fraser, A., May, M., Nelson, S. M., & Tilling, K. (2012). Blood pressure change in normotensive, gestational hypertensive, preeclamptic, and essential hypertensive pregnancies. *Hypertension*, HYPERTENSIONAHA-111. <https://doi.org/10.1161/HYPERTENSIONAHA.111.187039>
- Magee, L. A., von Dadelszen, P., Singer, J., Lee, T., Rey, E., Ross, S., ... & Gafni, A. (2016). The CHIPS Randomized Controlled Trial (Control of Hypertension in Pregnancy Study): is severe hypertension just an elevated blood pressure?. *Hypertension*, HYPERTENSIONAHA-116. <https://doi.org/10.1161/HYPERTENSIONAHA.116.07862>
- Mammaro, A. et al. (2009). *Hypertensive Disorders in Pregnancy* (3(1), pp. 1-5).
- McCarthy, E. A., Carins, T. A., Hannigan, Y., Bardien, N., Shub, A., & Walker, S. P. (2015). Effectiveness and

- safety of 1 vs 4 h blood pressure profile with clinical and laboratory assessment for the exclusion of gestational hypertension and pre-eclampsia: a retrospective study in a university affiliated maternity hospital. *BMJ open*, 5(11), e009492. <https://doi.org/10.1136/bmjopen-2015-009492>
- Melamed, N., Ray, J. G., Hladunewich, M., & Cox, B. (2014). Gestational hypertension and preeclampsia: are they the same disease?. *Journal of Obstetrics and Gynaecology Canada*, 36(7), 642-647. [https://doi.org/10.1016/S1701-2163\(15\)30545-4](https://doi.org/10.1016/S1701-2163(15)30545-4)
- Meshram, D. P. et al. (2014). *Maternal and foetal outcomes in Pregnancy Induced Hypertension -A hospital based study* (3(4), pp. 23-26).
- Moroy, P., Ustunyurt, E., Tapisiz, O. L., Tuncay, G., Aytan, H., Ustunyurt, O. B., ... & Mollamahmutoglu, L. (2007). Factors influencing the postpartum length of hospital stay in eclamptic women. *Taiwanese Journal of Obstetrics and Gynecology*, 46(4), 410-413. [https://doi.org/10.1016/S1028-4559\(08\)60012-1](https://doi.org/10.1016/S1028-4559(08)60012-1)
- Mylonas, I., & Friese, K. (2015). Indications for and risks of elective cesarean section. *Deutsches Ärzteblatt International*, 112(29-30), 489. <https://doi.org/10.3238/arztebl.2015.0489>
- National Collaborating Centre for Women's and Children's Health (UK). (2010). Hypertension in pregnancy: the management of hypertensive disorders during pregnancy. <https://doi.org/10.1136/bmj.c2207>
- NICE. (2011). Caesarean Section Guidelines', NICE Clinical Guideline (p. 282). <https://doi.org/10.1007/s00404-011-2201-6>
- Oberdörster, G., Castranova, V., Asgharian, B., & Sayre, P. (2015). Inhalation exposure to carbon nanotubes (CNT) and carbon nanofibers (CNF): Methodology and dosimetry. *Journal of Toxicology and Environmental Health, Part B*, 18(3-4), 121-212.
- Ota, E., Ganchimeg, T., Mori, R., & Souza, J. P. (2014). Risk factors of pre-eclampsia/eclampsia and its adverse outcomes in low-and middle-income countries: a WHO secondary analysis. *PloS one*, 9(3), e91198. <https://doi.org/10.1371/journal.pone.0091198>
- Plough, A. C., Galvin, G., Li, Z., Lipsitz, S. R., Alidina, S., Henrich, N. J., ... & McDonald, R. (2017). Relationship Between Labor and Delivery Unit Management Practices and Maternal Outcomes. *Obstetrics & Gynecology*, 130(2), 358-365. <https://doi.org/10.1097/AOG.0000000000002128>
- Reyes, L. M., García, R. G., Ruiz, S. L., Camacho, P. A., Ospina, M. B., Aroca, G., ... & López-Jaramillo, P. (2012). Risk factors for preeclampsia in women from Colombia: a case-control study. *PloS one*, 7(7), e41622. <https://doi.org/10.1371/journal.pone.0041622>
- Sohlberg, S., Stephansson, O., Cnattingius, S., & Wikström, A. K. (2012). Maternal body mass index, height, and risks of preeclampsia. *American journal of hypertension*, 25(1), 120-125. <https://doi.org/10.1038/ajh.2011.175>
- Uzan, J., Carbonnel, M., Piconne, O., Asmar, R., & Ayoubi, J. M. (2011). Pre-eclampsia: pathophysiology, diagnosis, and management. *Vascular health and risk management*, 7, 467. <https://doi.org/10.2147/VHRM.S20181>
- World Health Organization [WHO]. (2013). *Postnatal care of the mother and newborn 2013* (pp. 1-72). <https://doi.org/9789241506649>
- Zhang, S., Wang, L., Leng, J., Liu, H., Li, W., Zhang, T., ... & Hu, G. (2017). Hypertensive disorders of pregnancy in women with gestational diabetes mellitus on overweight status of their children. *Journal of human hypertension*, 31(11), 731. <https://doi.org/10.1038/jhh.2017.17>

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).

Type of Delivery and Length of Stay in Hypertension Patients

ORIGINALITY REPORT

11%

SIMILARITY INDEX

7%

INTERNET SOURCES

12%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Devianti Tandiallo, Mardiana Ahmad, Syafruddin Syarif, Nasrudin A. Mappaware, Prihantono Prihantono, Burhanuddin Bahar. "Web-Based Partograph on Early Detection of Emergency Cases and Referral Processes", Global Journal of Health Science, 2019
Publication

4%
- 2

Submitted to Universidad Nacional de Colombia
Student Paper

2%
- 3

Submitted to Universitas Hasanuddin
Student Paper

1%
- 4

Amal Almalki, Mohammed Khaled Al-Hanawi. "Public Private Partnerships and Collaboration in the Health Sector in the Kingdom of Saudi Arabia: A Qualitative Study", Global Journal of Health Science, 2018
Publication

1%
- 5

www.j3.jstage.jst.go.jp
Internet Source

1%

Exclude quotes On
Exclude bibliography On

Exclude matches < 25 words