

Recording Data Labour With Documentation Midwifery Based On Word Electric Browser (WEB)

Juwita¹, Mardiana Ahmad¹, Syafruddin Syarif^{1,2}, Nasruddin A.Mappaware^{1,3},
Prihantono^{1,4} & Burhanuddin Bahar^{1,5}

¹ Postgraduate Program, Departement of Midwifery, Hasanuddin University, Makassar, South Sulawesi, Indonesia.

² Departement of Electrical Engineering, Hasanuddin University, Makassar, South Sulawesi, Indonesia

³ Medical Faculty, Indonesia Muslim University, Makassar, South Sulawesi, Indonesia

⁴ Faculty of Medicine, Departement of Surgery, Hasanuddin University, Makassar, South Sulawesi, Indonesia

⁵ Faculty of Public Health, Department of Nutrition, Hasanuddin University, Makassar, South Sulawesi, Indonesia

Correspondence: Juwita, Postgraduate Program, Department of Midwifery, Hasanuddin University, Makassar, South Sulawesi 90245, Indonesia: Tel: 62-853-4230-2358. E-mail: juwitarahman8@gmail.com

Received: March 23, 2019 Accepted: April 29, 2019 Online Published: May 1, 2019

Abstract

Introduction: Midwifery documentation is a proof of recording and reporting possessed by midwives in carrying out records in documenting care provided.

Material and Methods: The research method used was the comparative study with the comparative descriptive design. The total subjects of 30 inpartu mothers were chosen using the purposive sampling technique. The subjects were divided into two groups: 15 of the mothers used the conventional midwifery documentation and 15 of them used the WEB-based. The data were analyzed using the univariate analysis in order to look at the median value, and the bivariate analysis with Mann-Whitney test.

Results: The study results indicated the the highest frequency was found in the WEB-based partograph, namely he filling speed was 26 (86.7%), the highest frequency of partograph filling truth was that of the WEB-based partograph

= 25 (83.3%), while the highest Relevance of the data frequency was found in the WEB-based partograph with the data Relevance = 27 (90%). The result of the statistical test had the p value of 0.000, 0.000, and 0.000 ($<\alpha=0.05$) meaning there was a difference of the WEB-based midwifery documentation compared to the conventional midwifery documentation

Conclusion: Thus it can be concluded that web-based midwifery documentation is more effective than conventional midwifery documentation. There are differences in the aspects of ease, speed, security and relevance of data to recording labor history data so that it needs to be applied in documenting midwifery care

Keywords: WEB-based midwifery documentation, conventional

1. Introduction

Partograph is a tool to monitor labor delivery, assess and manage labor, detect problems early and complicate labor, and record and report on their labor data. In connection with patient referral and documentation of partograph, complete and systematic recorded data are needed starting from the first stage of labor to the fourth stage in a data storage system in the computer. Data Storage System used is the data storage system My SQL (Structured Query Language) as an RDBMS (Relational Database Management System) which is distributed free under the GPL (General Public License) license which can be used in terms of data storage, data and information management. (Baillie, Chadwick, Mann, & Brooke-Read, 2013; Kitila, Gmariam, Molla, & Nemera, 2014; Asibong et al., 2014; Okokon, Oku, Agan, Asibong, Essien, & Monjok, 2014; Anita, 2016; Andrianto, 2017; Balikuddembe,

Tumwesigye, Wakholi, & Tylleskär, 2017; Bedwell, Levin, Pett, & Lavender, 2017; Erawantini et al., 2017; Mandiwa & Zamawe, 2017; Fatusi, Makinde, Adeyemi, Orji, & Onwudiegwu, 2008; Nugroho, Fitriasi, & Widada, 2015; Ramdhani, Isnanto, & Windasari, 2015; Sovia & Febio, 2011; Sundari, 2016; Widyaningsih & Astutingsih, 2016).

The study obtained a total of 1070 partograph reviewed, 58.6% of the partograph had no record of maternal blood pressure, 65.3% had no documentation of temperature, 25.4% Molding was not recorded, 14.9% of fetal heart rate was not recorded and 12.0% of the decrease in fetal head is not recorded so that there are still many shortcomings in systematic recording and documentation (Balikuddembe et al., 2017; Mandiwa & Zamawe, 2017; Nurmiyati, 2011). According to WHO, the use of partograph will reduce long labor 6.4% to 3.4% of cases of Caesarea 44% to 21%. At present partograph recording and reporting has not been implemented properly as a result of abnormalities and complications that occur in the labor process often not detected so that it slows in clinical decision making. During this time reporting and documentation by midwives was based on recording in partograph on paper, however, based on several studies said that there were still many midwives who did not complete the data in partograph properly and systematically, therefore a recording system, faster reporting and documenting partograph in a computer and web-based data storage to facilitate the work of midwives in storing and re-accessing current and past labor history so as to prevent partograph file accumulation (Andrea, Dodd, 2014; Fahdhy & Chongsuvivatwong, 2005; Yisma, Dessalegn, Astatkie, & Fesseha, 2013a, 2013b; Baillie et al., 2013; Gans-Lartey, O'Brien, Gyekye, & Schopflocher, 2013; Ollerhead and Osrin, 2014; Aliona Masika, Peter Katongole, & Govule, 2015; JNPK-KR, 2012; Bailey, Wilson, & Yoong, 2015; Opoku & Nguah, 2015; Fatouh, 2015; Vidyashri & Nagarathna, 2015; Nugroho, Fitriasi, & Widada, 2015; Byukusenge et al., 2016; Kwast, 1994; Yisma et al., 2013a; Opoku & Nguah, 2015; Sinaga, 2016; Susilowati & Sulistiyaningsih, 2011; Tayade & Jadhao, 2012; WHO, 2011; 2014). Based on the description above, the author was motivated to develop a web-based partograph midwifery documentation database research system. This is intended to streamline the process of recording, reporting and documenting partograph in aspects of speed, convenience, security, and convenience by using the Subjective, Objective, Analysis and Management (SOAP) methods. And in accessing again and seeing the recording of labor history and actions taken by the midwife so that it can accelerate clinical decisions, plan actions taken and streamline reporting, documentation in midwifery and prevent the accumulation of partograph files (Bhatt, Kar, Shashank, & Somarajan, 2004; Chaturvedi, S. et al. 2015; Erawantini, F. et al. 2017; Hadi et al., 2017; Hasan, Hossain, Arafat, Khan, Ahmed, & Shabnam, 2017; Underwood, Sterling, & Bennett, 2012; Schweers, Khalid, Underwood, Bishnoi, & Chhugani, 2016; Jennifer, 2012; Souza et al., 2015; Lavender, Hart, & Smyth, 2013; Mathibe-Neke, Lebeko, & Motupa, 2013; Kitila et al., 2014; Kamath, Nagarathna and Sharanya, 2015; Mandiwa and Zamawe, 2017; Lew & Ghassemzadeh, 2018; Nursing and Midwifery Council 2010a; 2010b; Sama, C. B. et al. 2017; Seo, J.-H. et al. 2016; Sinha, S. Shrivastava, & S. Shrivastava, 2016; Sudarti, Fuziah. 2011; Varney, 2008)

2. Materials and Methods

2.1 Design of the Study and Sample Population

The research design used was quasy experiment using a treatment group on web-based partograph midwifery documentation and control groups in conventional partograph obstetric documentation. This research was conducted at Bara-Barayya Health Center, Antang Public Health Center and Makassar City Health Service from November 2018 to January 2019. The research tool here is a hardware and software component that is used as a tool in supporting the research carried out, this tool will be used as a data processing tool and the making of the system to be designed. The research method used was the comparative study with the comparative descriptive design. The total subjects of 30 mothers in labor were chosen using the purposive sampling technique. The subjects were divided into two groups: 15 of the mothers used the conventional midwifery documentation and 15 of them used the WEB-based midwifery documentation. The data were analyzed using the univariate analysis in order to look at the median value and the bivariate analysis with Mann-Whitney test. Sampling from the study population was carried out by purposive sampling, where the population that met the inclusion criteria was included as sample members. The number of subjects was 30 patient data, 30 childbirth patients were documented using conventional partograph midwifery documentation and web-based partograph midwifery documentation. Samples were divided into 2 groups of data, 15 patient data filled using conventional partograph (manual) and 15 patient data filled using computer-based partograph.

Furthermore, 30 patient data will be filled by 9 midwives in the maternity room of Antang Perumnas Public Health Center, 8 midwives in the delivery room of Bara-Baraya Makassar Public Health Center and 1 Midwife at the Makassar City Health Office will see the partograph documentation as data reporting, 17 of these midwives will

fill in 30 patient data using conventional partograph midwifery documentation (manual) and patient data is filled using web-based partograph midwifery documentation. Midwives of Public Health Center who were used as subjects were categorized by age, length of work, training that had been attended. The results of the filling will be assessed using an observation sheet. The primary data obtained directly on the data source is when conducting interviews with midwives and delivery mothers, making observations on filling partograph documentation carried out by midwives on the progress of labor, care provided and results obtained.




Figure 1. The Flowchart of the Study

2.2 Data Collection

The primary data obtained directly on the data source is when observing the actions taken by the midwife and the results recorded, namely at the Antang Perumnas Health Center, Makassar Bara-Barayya Health Center and Makassar City Health Office. Secondary data obtained indirectly to support writing in this study through documents or records of medical records of mothers who have given birth at the research site. The total subjects of 30 inpartu mothers were chosen using the purposive sampling technique. The subjects were divided into two groups: 15 of the mothers used the conventional midwifery documentation and 15 of them used the WEB-based midwifery documentation namely at the Antang Perumnas Public Health Center, Makassar Bara-Barayya Public Health Center and Makassar City Health Office.

2.3 Ethical Considerations

This study received ethical clearance from the Research and Ethics Committee of Faculty of Medicine, Hasanuddin University and was registered in 972/h4.8.4.5.31/PP36-KOMETIK/2018. And protocol number: UH18100691.

2.4 Data Management and Analysis

Descriptive analyses were presented as mean $\pm$ standard deviation and frequencies for categorical variables. Bivariate analyses were analyzed and processed using independent t-test in pairs, Mann-Whitney to determine the differences between all groups at 5% level of significance. Spearman's correlation was used to determine further correlations. Data were analyzed using the Statistical Package for Social Science (SPSS) version 24 for Windows.

3. Results

Sampling from the study population was carried out by purposive sampling, where the population that met the inclusion criteria was included as sample members. The number of subjects was 30 patient data, 30 deliveries were documented using conventional partograph midwifery documentation and web-based partograph midwifery documentation. Samples are divided into 2 groups of data, 15 patient data filled using conventional partograph (manual) and 15 patient data filled using computer-based partograph.

3.1 Univariate Analysis

Midwifery Documentation Comparison From Time Speed Aspects, Truths Charging, Data Security and Data Relevance Midwifery Documentation Website-Based Partographs

Table 1. Characteristics of filling in website system partograph midwifery documentation and conventional partograph midwifery documentation from aspects of ease of filling, time speed, data security, data relevance

Characteristics	Midwifery Documentation	
	Website Based	Conventional
	F %	F %
Time Speed		
Hurry up	26 (86.7)	
Long	4 (13.3)	30 (100)
Truth of Charging		
Right	25 (83.3)	9 (30)
Not exactly	5 (16.7)	21 (70)
Data Relevance		
Right	27 (90)	9 (30)
Not right	3 (10)	21 (70)
Total	100	100

Source : Primary Data, Paired t test.

Table 2. Average scores for filling website-based partograph documentation and conventional partographs from the speed aspect of charging time

Characteristics	Mean	SD	Min – Maks
Charging time			
Website-based partograph documentation	21.7	1	19-24
Conventional Partographical Documentation	31.1	1	29-34

Source: Primary Data, Paired t test.

3.2 Bivariate Analysis

Analysis of differences in website-based partograph midwifery documentation and conventional partograph midwifery documentation against.

Table 3. Differences in midwifery partographies of websites based on websites and midwifery documentation of conventional partographs on recording of childbirth history data

Variable	N	P value	α
Website and conventional time speed aspects are	30	0.000	0.05
Truthful aspects of charging websites and conventional	30	0.000	
Relevance aspects of website data and conventional	30	0.000	

Source: Primary Data, Mann-Withney Test.

4. Discussion

Table 1 The results obtained the speed of time needed for filling Website-based partograph and conventional partograph, the highest frequency was found on web-based partograph, namely fast charging 26 (86.7%), while the truth of the highest frequency partograph filling was website-based partograph, 25 (83.3%) . Data security obtained the highest frequency, namely on website-based partograph, which is 30 (100%), while the highest frequency data relevance is website-based partograph with data relevance of 27 (90%). The results of this study indicate that the use of website-based partograph midwifery documentation will make it easier for midwives to process the recording, reporting and recording of labor history data, website-based partograph midwifery documentation will minimize errors and have a good level of truth. This is in line with the research conducted by Heather et al., 2012, Heather et al., 2014 which found that the use of electronic partograph was more effective and efficient with the use of hardware and software as a documentation and recording of labor history that could be used as a tool to train labor health about how to use the correct partograph with limited human resources so that it has an impact on the completeness of partograph filling, reducing the error rate (Qureshi, Sekadde-Kigundu, & Mutiso, 2010; Pirie, 2011; Ontario, 2013; Muslihatun et al., 2013; Ollerhead & Osrin, 2014; Sharma, Deka, & Das, 2015).

Table 2 gaining from the speed of filling the website-based partograph midwifery documentation and conventional midwifery documentation, the average time required for filling in website-based partograph midwifery documentation was 21.7 minutes, with a standard deviation of ± 1 second, while documentation of conventional partograph midwifery time needed to fill more that is 31.1 minutes with a standard deviation of ± 1 second.

The results of research conducted by Yulianti et al. (2018) got the use of electronic partograms in recording and reporting partograph documentation with incoming data speeds of 82% with an error rate of 18%, in the process of recording and reporting the time used for charging is relatively shorter 8.6 hours, so it can be concluded that the use of electronic partograms allows data to enter faster with a lower error rate than conventional partograms

Table 3 shows the analysis of data using Test, obtaining a p value of 0.000 (<0.05) means that there are differences in website-based partograph midwifery documentation and conventional partograph midwifery documentation on recording delivery history data from aspects of filling filling truth, data security and data relevance (Sugiyono, 2014).

The statistical test results require an average time of 21.7 minutes, ease of filling is also seen in the correct level of charging that is true 83.3%. The ease of filling will have an effect on the time needed to be shorter, the human resources needed are more simple and the quality of midwifery services is more optimal. This is in accordance with the research conducted stating that for facilitation aspects in the process of documenting midwifery and care provided, it gives information to officers on what has been recorded so that it makes it easier in the process of managing subsequent actions (Lavender, Hart and Smyth, 2013; Fatouh & Ramadan, 2015; Ontario, C. of M. of 2013; Toemandoek, Wagey, & Loho, 2015).

5. Conclusion

Website-based partograph midwifery documentation is better than conventional midwifery documentation seen from the aspect of a short time velocity of 21.7 minutes using web-based partograph midwifery documentation and obtaining a 31.1 minute filling time speed in using conventional partograph midwifery documentation, the correct aspect of filling with columns filled with complete, systematic and appropriate, and relevant data relevance because it is stored in digital form. A web-based partograph midwifery documentation database effectively re-accesses labor history records and actions that have been given.

Acknowledgements

The authors would like to express their special thanks to the Antang Perumnas Public Health Center, Bara-Baraya Public Health Center, Makassar City Health Office and staff for their support and assistance.

Competing Interests Statement

The authors declare that there are no competing or potential conflicts of interest.

References

- Kitila, S. B., Gmariam, A., Molla, A., & Nemera, G. (2014). Utilization of partograph during labour and birth outcomes at Jimma University. *Journal of Pregnancy and Child Health*, 1(101), 2. <https://doi.org/10.4172/2376-127X.1000101>

- Fatusi, A. O., Makinde, O. N., Adeyemi, A. B., Orji, E. O., & Onwudiegwu, U. (2008). Evaluation of health workers' training in use of the partogram. *International Journal of Gynecology & Obstetrics*, 100(1), 41-44. <https://doi.org/10.1016/j.ijgo.2007.07.020>
- Ashish KC, Johan Wrammert, Robert B.Clark, Uwe Ewalt, Mats Malqvist (2016). Inadequate Fetal Heart Rate Monitoring And Poor Use Of Partogram Associated With Intrapartum Stillbirth: a Case-Referent Study In Nepal. *BMC Pregnancy and Childbirth*. doi: 10.1186/s12884-016-1034-5. <https://doi.org/10.1186/s12884-016-1034-5>
- Aliona Masika, M., Peter Katongole, S., & Govule, P. (2015). Improving Partograph Documentation and Use by Health Workers of Bwera Hospital: A Process Improvement Research. *International Journal of Nursing and Health Science*. Retrieved from <http://www.openscienceonline.com/journal/ijnhs>
- Andrianto, P. (2017). Web Web Based Health Service Information System at Public Health Center. *Proceedings of the National Computer and Informatics Seminar (SENASKI)*.
- Anita, W. (2016) Learning Methods in Partographic Documentation Skills in Midwifery Care in Childbirth: Literature Review. *Jurnal Endurance*. <https://doi.org/10.22216/jen.v1i3.1073>
- Asibong, U., Okokon, I. B., Agan, T. U., Oku, A., Opiah, M., Essien, E. J., & Monjok, E. (2014). The use of the partograph in labor monitoring: A cross-sectional study among obstetric caregivers in General Hospital, Calabar, Cross River State, Nigeria. *International journal of women's health*, 6, 873. <https://doi.org/10.2147/IJWH.S49188>
- Bailey, S., Wilson, G., & Yoong, W. (2015). What factors affect documentation by midwives? A prospective study assessing relationship between length of shift, workload and quality of note keeping. *Midwifery*, 31(8), 787-792. <https://doi.org/10.1016/j.midw.2015.04.001>
- Baillie, L., Chadwick, S., Mann, R., & Brooke-Read, M. (2013). A survey of student nurses' and midwives' experiences of learning to use electronic health record systems in practice. *Nurse Education in Practice*, 13(5), 437-441. <https://doi.org/10.1016/j.nepr.2012.10.003>
- Balikuddembe, M. S., Tumwesigye, N. M., Wakholi, P. K., & Tylleskär, T. (2017). Computerized childbirth monitoring tools for health care providers managing labor: a scoping review. *JMIR medical informatics*, 5(2), e14. <https://doi.org/10.2196/medinform.6959>
- Bedwell, C., Levin, K., Pett, C., & Lavender, D. T. (2017). A realist review of the partograph: when and how does it work for labour monitoring?. *BMC pregnancy and childbirth*, 17(1), 31. <https://doi.org/10.1186/s12884-016-1213-4>
- Bhatt, B. M., Kar, G., Shashank, S., & Somarajan, S. (2013, September). Designing interfaces for healthcare workers: a case study of the electronic partogram. In *Proceedings of the 11th Asia Pacific Conference on Computer Human Interaction* (pp. 187-191). ACM. <https://doi.org/10.1145/2525194.2525268>
- Byukusenge, J. B., Adomako, E., Lukas, S., Mugarura, C., Umucyo, J., Mukagatare, S., ... & Wong, R. (2016). Implementing a labor monitoring guideline and midwives responsibilities to increase the completion rate of partograph in Muhima Hospital, Rwanda. *On the Horizon*, 24(4), 335-340. <https://doi.org/10.1108/OTH-07-2016-0036>
- Erawantini, F. et al. (2017). Design Input Data Determining Risk Factors for Early Type 2 Diabetes Mellitus Based on the Web. *Health Journal*, 5(1).
- Fahdhy, M., & Chongsuvivatwong, V. (2005) Evaluation of World Health Organization partograph implementation by midwives for maternity home birth in Medan, Indonesia. *Midwifery*. <https://doi.org/10.1016/j.midw.2004.12.010>
- Fatouh, E., & Ramadan, S. (2015). Effect of Using Paperless Partogram on the Management and Outcome of Labour and the Nurses' Opinion. *Journal of Education and Practice*, 6(8), 17-23. Retrieved from <https://eric.ed.gov/?id=EJ1082718>.
- Gans-Lartey, F., O'Brien, B. A., Gyekye, F. O., & Schopfloch, D. (2013). The relationship between the use of the partograph and birth outcomes at Korle-Bu teaching hospital. *Midwifery*, 29(5), 461-467. <https://doi.org/10.1016/j.midw.2012.03.002>

- Hadi, S. P. I., Kuntjoro, T., Sumarni, S., Anwar, M. C., Widyawati, M. N., & Pujiastuti, R. S. E. (2017). The Development of E-partograph Module as a Learning Platform for Midwifery Students: The ADDIE Model. *Belitung Nursing Journal*, 3(2), 148-156. <https://doi.org/10.33546/bnj.77>
- Hasan, M. J., Hossain, M. S., Arafat, S. Y., Khan, A. S., Ahmed, Z., & Shabnam, F. (2017). Comparative Analysis of Life Curve Mobile Application: an Easier Alternative to Existing Traditional Paper Partograph. *IOSR-JMCA*. <https://doi.org/10.9790/0050-04015561>
- Underwood, H., Sterling, S. R., & Bennett, J. K. (2012, October). Improving maternal labor monitoring in Kenya using digital pen technology: a user evaluation. In *2012 IEEE Global Humanitarian Technology Conference* (pp. 282-287). IEEE. <https://doi.org/10.1109/GHTC.2012.77>
- Schweers, J., Khalid, M., Underwood, H., Bishnoi, S., & Chhugani, M. (2016). mLabour: Design and evaluation of a mobile partograph and labor ward management application. *Procedia engineering*, 159, 35-43. <https://doi.org/10.1016/j.proeng.2016.08.061>
- Souza, J. P., Oladapo, O. T., Bohren, M. A., Mugerwa, K., Fawole, B., Moscovici, L., ... & Tunçalp, Ö. (2015). The development of a simplified, effective, labour monitoring-to-action (SELMA) tool for better outcomes in labour difficulty (BOLD): study protocol. *Reproductive health*, 12(1), 49. <https://doi.org/10.1186/s12978-015-0029-4>
- JNPK-KR. (2012). *Normal Delivery Care*. Jakarta: Indonesian Ministry of Health.
- Vidyashri, K. C., & Nagarathna, G. (2015). Documentation of the modified WHO partograph during labour in a South Indian tertiary care hospital. *Journal of Evolution of Medical and Dental Sciences*, 4(82), 14415-14422. <https://doi.org/10.14260/jemds/2015/2050>
- Kennedy, H. P., & Shannon, M. T. (2004). Keeping birth normal: research findings on midwifery care during childbirth. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 33(5), 554-560. <https://doi.org/10.1177/0884217504268971>
- Kitila, S. B., Gmariam, A., Molla, A., & Nemera, G. (2014). Utilization of partograph during labour and birth outcomes at Jimma University. *Journal of Pregnancy and Child Health*, 1(101), 2. <https://doi.org/10.4172/jpch.1000101>.
- Kwast, B. E. (1994). World Health Organization partograph in management of labour. World Health Organization Maternal Health and Safe Motherhood Programme. *Lancet* (London, England). [https://doi.org/10.1016/S0140-6736\(94\)92528-3](https://doi.org/10.1016/S0140-6736(94)92528-3)
- Landry, E., Pett, C., Fiorentino, R., Ruminjo, J., & Mattison, C. (2014). Assessing the quality of record keeping for cesarean deliveries: results from a multicenter retrospective record review in five low-income countries. *BMC pregnancy and childbirth*, 14(1), 139. <https://doi.org/10.1186/1471-2393-14-139>
- Lavender, T., Hart, A., & Smyth, R. M. (2013). Effect of partogram use on outcomes for women in spontaneous labour at term. *Cochrane database of systematic reviews*, (7). <https://doi.org/10.1002/14651858.CD005461.pub4>
- Lew, V. and Ghassemzadeh, S. (2018) SOAP Notes. *StatPearls*, 1. <https://doi.org/10.1016/B978-0-323-07759-0.00020-1>.
- Mathibe-Neke, J. M., Lebeko, F. L., & Motupa, B. (2013). The partograph: A labour management tool or a midwifery record?. *International Journal of Nursing and Midwifery*, 5(8), 145-153. <https://doi.org/10.5897/IJNM2013.0115>
- Mandiwa, C., & Zamawe, C. (2017). Documentation of the partograph in assessing the progress of labour by health care providers in Malawi's South-West zone. *Reproductive health*, 14(1), 134. <https://doi.org/10.1186/s12978-017-0401-7>
- Markos, D., & Bogale, D. (2015). Documentation status of the modified World Health Organization partograph in public health institutions of Bale zone, Ethiopia. *Reproductive health*, 12(1), 81. <https://doi.org/10.1186/s12978-015-0074-z>
- Maureen D, et al. (2015). 'Documentation in labour among midwives in Madonna University Teaching Hospital elele, river state, Nigeria. *Internastional Journal of Refroduction, Contraception, Obstetrics and Gyneekology*. <https://doi.org/10.18203/2320-1770.ijrcog20150719>

- Mathews, J. E., Rajaratnam, A., George, A., & Mathai, M. (2007). Comparison of two World Health Organization partographs. *International Journal of Gynecology & Obstetrics*, 96(2), 147-150. <https://doi.org/10.1016/j.ijgo.2006.08.016>
- Muslihatun, et al. (2013). *Midwifery Documentation*. Fitramaya : Yogyakarta.
- Nugroho, B., Fitriasih, S. H., & Widada, B. (2015). Medical Record Information System at Masaran I Sragen Public Health Center. *Jurnal TIKomSiN*. <https://doi.org/10.30646/TIKOMSIN.V3I2.203>.
- Nurmiyati, T. (2011). Relationship between Knowledge Level and Midwife Compliance with Partograph Use in Madukara District and Pagentan Subdistrict, Banjarnegara Regency. *Permata Medika Scientific Journal*, 2(1). Retrieved from <http://www.journal.stikes-bpm.ac.id/index.php/arsip/volume-2-no-1-juli-2013/17>.
- Nursing and Midwifery Council. (2010a). Standards for medicines management Record keeping. *Standards for Medicine Management*. Retrieved from <https://www.nmc.org.uk/standards/standards-for-post-registration>.
- Nursing and Midwifery Council. (2010b). Standards Record of keeping for proficiency nurse and midwife prescribers. *Education And Training*. Retrieved from <https://www.nmc.org.uk/standards/standards-for-post-registration>
- Okokon, I. B., Oku, A. O., Agan, T. U., Asibong, U. E., Essien, E. J., & Monjok, E. (2014). An evaluation of the knowledge and utilization of the partograph in primary, secondary, and tertiary care settings in Calabar, South-South Nigeria. *International journal of family medicine*, 2014. <https://doi.org/10.1155/2014/105853>
- Ollerhead, E., & Osrin, D. (2014). Barriers to and incentives for achieving partograph use in obstetric practice in low-and middle-income countries: a systematic review. *BMC pregnancy and childbirth*, 14(1), 281. <https://doi.org/10.1186/1471-2393-14-281>
- Ontario, C. of M. of. (2013). *Record Keeping Standard for Midwives, CMO Policy Suite*.
- Opoku, B. K., & Nguah, S. B. (2015). Utilization of the modified WHO partograph in assessing the progress of labour in a metropolitan area in Ghana. *Research Journal of Women's Health*, 2(1), 2. <https://doi.org/10.7243/2054-9865-2-2>
- Oxford. (2013). *Oxford Handbook of Midwifery*. Kebedanan Oxford: From Midwives to Midwives.
- Perinatal Health Program. (2010). *Perinatal Forms Guideline 4 A Guide For Completion of the British Columbia Labour Partogram* (BCPHP 1583).
- Pirie, S. (2011) Documentation and record keeping. *Journal of perioperative practice*. <https://doi.org/10.1177/175045891102100103>
- Qureshi, Z. P., Sekadde-Kigundu, C., & Mutiso, S. M. (2010). Rapid assessment of partograph utilisation in selected maternity units in Kenya. *East African medical journal*, 87(6), 235-241. <https://doi.org/10.4314/eamj.v87i6.63081>
- Ramdhani, A. et al. (2015). Development of Expert Systems for Diagnosis of Web-Based Hepatitis Using Certainty Factor. *JTsiskom*.
- Sama, C. B., Takah, N. F., Danwe, V. K., Melo, U. F., Dingana, T. N., & Angwafo III, F. F. (2017). Knowledge and utilization of the partograph: A cross-sectional survey among obstetric care providers in urban referral public health institutions in northwest and southwest Cameroon. *PloS one*, 12(2), e0172860. <https://doi.org/10.1371/journal.pone.0172860>
- Seo, J. H., Kong, H. H., Im, S. J., Roh, H., Kim, D. K., Bae, H. O., & Oh, Y. R. (2016). A pilot study on the evaluation of medical student documentation: assessment of SOAP notes. *Korean journal of medical education*, 28(2), 237. <https://doi.org/10.3946/kjme.2016.26>
- Sharma, R., Deka, G., & Das, G. C. (2015). The Paperless partograph - The new user-friendly and simpler tool for monitoring labour. *IOSR Journal of Dental and Medical Sciences*. <https://doi:10.9790/0853-141193943>.
- Sinaga, R. N. (2016). Analysis and Design of a Medical Record Information System at the Akper Clinic in the Middle/ Srivijaya White Garuda. *Journal of Information Systems Management*, 1(1), 2540-8011.
- Sinha, D., Shrivastava, S., & Shrivastava, S. (2016). Management of Labour in Primigravida with Who Modified Partograph. *Scholars Journal of Applied Medical Sciences*, 4, 3457-3461. <https://doi:10.21276/sjams.2016.4.9.59>

- Sovia, R., & Febio, J. (2011). Build e-library applications using html, php script, and mysql database. *Journal of Information and Education Technology*, 3(1), 2011. Retrieved from <http://ejournal.stikom-db.ac.id/index.php/processor/article/view/28>
- Sudarti, F. (2011). *Textbook for Midwifery Documentation*. Nuha Medika: Jakarta.
- Sugiyono. (2014). *Educational Research Methods Quantitative, Qualitative Approach and R & D*. Bandung: Alfabeta.
- Sundari, J. (2016). Web-Based Puskesmas Service Information System. *Indonesian Journal on Software Engineering. IJSE - Indonesian Journal on Software Engineering*, 2(1).
- Susilowati, & Sulistyaningsih, S. H. (2011). Analysis of Factors Associated with Making Partographs by Midwives at Dukuhseti Health Center and Tayu Ii Health Center, Pati Regency. *Midwifery and Health Journal*, 3(2).
- Tayade, S., & Jadhao, P. (2012). The impact of use of modified WHO partograph on maternal and perinatal outcome. *International Journal of Biomedical and Advance Research*, 3(4), 256-262. <https://doi.org/10.7439/ijbar.v3i4.398>
- Toemandoek, J. P., Wagey, F., & Loho, M. (2015). Knowledge and Attitudes of Midwives Regarding the Use of Partographies in Childbirth at Poned Public Health Center, Manado City. *Journal e-Clinic (eCI)*, 3(2). Retrieved from <https://ejournal.unsrat.ac.id/index.php/eclinic/article/view/9344/8921>
- Varney, H. (2008). *Varney's Midwifery*. EGC : Jakarta.
- Widyaningsih, P., & Astutingsih, A. (2016). Web-Based Expert System Application for Consultation on Pregnancy Problems Using Forward Chaining and Production Rule. *INITI MEDICAL APPLICATION Journal, IMAGE MEDIKA SURAKARTA*.
- Wildan dan Hidayat. (2009). *Midwifery Documentation*. Salemba Medika: Jakarta.
- World Health Organization [WHO]. (2011). *WHO recommendations for Induction of labour*. World Health Organization. [https://doi.org/10.1016/0140-6736\(93\)92727-B](https://doi.org/10.1016/0140-6736(93)92727-B).
- World Health Organization [WHO]. (2014). *WHO recommendations for augmentation of labour*. World Health Organization. [https://doi.org/10.1016/0140-6736\(93\)92727-B](https://doi.org/10.1016/0140-6736(93)92727-B).
- Yisma, E., Dessalegn, B., Astatkie, A., & Fesseha, N. (2013a). Completion of the modified World Health Organization (WHO) partograph during labour in public health institutions of Addis Ababa, Ethiopia. *Reproductive health*, 10(1), 23. <https://doi.org/10.1186/1742-4755-10-23>
- Yisma, E., Dessalegn, B., Astatkie, A., & Fesseha, N. (2013b). Knowledge and utilization of partograph among obstetric care givers in public health institutions of Addis Ababa, Ethiopia. *BMC pregnancy and childbirth*, 13(1), 17. <https://doi.org/10.1186/1471-2393-13-17>
- Yulianti, N. T., Ahmad, M., Syarif, S., Budu, & Sunarno, I. (2017). Comparison Partograph Based on Computer System and Partograph Conventional on Monitoring Record Report Progress of Labor. *International Journal of ChemTech Research*, 11(05). 363-368. <https://doi.org/10.20902/IJCTR.2018.110540>

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).