

ALUR PROSES KORESPONDENSI

“Impact of iso-protein feed with different carbohydrate and lipid ratios on growth and feed efficiency of mud crab, *Scylla serrata*”

Hasnidar Hasnidar, Andi Tamsil, Nursyahrani

**NAMA JURNAL : Egyptian Journal of Aquatic Biology & Fisheries
Volume 28(2): 67-81 (2024)**

No.	Kegiatan	Waktu	Keterangan
A.	Article profile (article review procedur)		By OJS
	1. Submission summary	9 Juli 2024	
B.	Article profile (article review procedur)		By email
	1. Editorial system registration	31 Agustus 2022	
	2. “Submitted Manuscript to the Egyption Journal journal ”	22 Desember 2023	
	3. Under review		
	a. Manuskrip awal		
	b. Proses review	29 Desember 2023- 2 Maret 2024	
	c. Ketengan sudah publish		
C	Publish	11 Maret 2024	

- Editor-in-Chief Comments
- ✓ Select Manuscript Type
- ✓ Enter Manuscript Title
- ✓ Add/Remove Authors
- ✓ Enter Abstract
- ✓ Enter Keywords
- ✓ Select Related Subjects
- ✓ Additional Comments
- Attach Files
- ✓ Cover Letter & Checklist
- ➔ Finish Submission

Finish Submission

You can not finish your submission until completing the:
Attach Files (Manuscript Main File)

Your Submission Summary

Manuscript ID	EJABF-2312-3892 (R1)
Manuscript Title	Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, Scylla serrata
Manuscript Type	Original Article
Main Subjects	Aquaculture of Aquatic Fauna
Abstract	The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aims to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, Scylla serrata. Mud crabs with a weight of 70±4.7 g/head, and a carapace length of 5±0.5 cm were reared using crab boxes and placed in ponds, for 60 days. Pterygoplichthys pardalis meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios, i.e. treatments A (31 and 10%); B (33 and 9%); C (34 and 7%); and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage/MP; growth (absolute growth/WG, specific growth rate/SGR); feed efficiency (protein retention/PR, lipid retention/LR, feed conversion ratio/FCR, and survival/SR); and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with a ratio of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment, i.e. 31 and 10%, is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.
Keywords	mud crab, protein, carbohydrate, lipid ratio
Suggested Reviewers	Tahya, Akbar Marzuki
Comments	good

Authors

#	Name	Email Address	Degree	Position	Phone	Country	Affiliation
1	Hasnidar, Hasnidar* 	hasnidar.yasin@umi.ac.id	PhD	Associate Professor	085259595996	Indonesia	Faculty of Fisheries and Marine Science, Universitas Muslim Indonesia
2	Andi Tamsil, Andi Tamsil	andi.tamsil@umi.ac.id	PhD	Associate Professor	0821-8838-8332	Indonesia	Muslim University Indonesia
3	syahran, Nursyahran 	nursyahran00@gmail.com	MSc	Other	085255233596	Indonesia	Institut Teknologi dan Bisnis Maritim Balikpapan Makassar

Submit Date	2023-12-22 15:53:32
Current Status	Manuscript Needs Revision (Minor Revision)
Modify Date	2024-09-07 11:06:09

Related Files

#	File Type	File Name	Size	File Description	Upload Date
1	Title Page	Title Page.docx	12.75 KB	Autor Name	2023-12-22
2	Manuscript Main File	Manuskrip Impact of iso-protein.docx	99.76 KB	Manuscrip	2023-12-22
3	Table	Table 1 (1).docx	27.93 KB		2023-12-22

4	Archived Main File	Manuskrip Impact of iso-protein.docx	99.76 KB	EJABF-2312-3892	2023-12-22
---	--------------------	--------------------------------------	----------	-----------------	------------

[Decline Revision](#)[Previous](#)



Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Editorial system registration

1 pesan

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
Balas Ke: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
Kepada: hasnidar.yasin@umi.ac.id

31 Agustus 2022 pukul 11.57

Dear Hasnidar Hasnidar,

Welcome to **Egyptian Journal of Aquatic Biology and Fisheries** website.

Thank you for your registration in the Editorial System Online Submission and peer review tracking system.
Below please find your username and confidential password, which you need to access the system at:

<https://ejabf.journals.ekb.eg/>

Your username is: **En131164**

Your password is: **iy9mr8rq84**

Please save this information in a safe place.

Once you login , you can change your password on your profile.

Editorial Office

Egyptian Journal of Aquatic Biology and Fisheries

<https://ejabf.journals.ekb.eg/>



Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Acknowledgement of Submission (Manuscript #EJABF-2312-3892)

1 pesan

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
Balas Ke: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
Kepada: hasnidar.yasin@umi.ac.id
Cc: ejabf2017@gmail.com

22 Desember 2023 pukul 20.53

Manuscript ID: EJABF-2312-3892

Manuscript Title: **Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, Scylla serrata**

Authors: Hasnidar Hasnidar, Andi Tamsil Andi Tamsil, Nursyahran syahran

Dear **Dr. Hasnidar Hasnidar**

I wish to acknowledge receiving the of the above mentioned manuscript.

It should be noted that the manuscript will be reviewed for possible publication in the Scientific Journals Management System.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Our editorial decision will be brought to your attention once the paper has been reviewed due the referees consideration.

I wish to take this opportunity to thank you for sharing your work with us.

Truly yours,

Executive managing Editor of **Egyptian Journal of Aquatic Biology and Fisheries**

Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, *Scylla serrata*

Hasnidar¹, Andi Tamsil²

^{1,2}Department of Aquaculture, Faculty of Fisheries and Marine Sciences. Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km. 05 Makassar 90231, South Sulawesi, Indonesia.

*Corresponding Author: hasnidar.yasin@umi.ac.id

ARTICLE INFO

Article History:

Received:

Accepted:

Online:

Keywords:

mud crab
protein
carbohydrate
lipid
ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aims to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70 ± 4.7 g/head, and a carapace length of 5 ± 0.5 cm were reared using crab boxes and placed in ponds, for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios, i.e. treatments A (31 and 10%); B (33 and 9%); C (34 and 7%); and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage/MP; growth (absolute growth/WG, specific growth rate/SGR); feed efficiency (protein retention/PR, lipid retention/LR, feed conversion ratio/FCR, and survival/SR); and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with a ratio of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment, i.e. 31 and 10%, is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

INTRODUCTION

Mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for

crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if carried out with low production costs (Apine *et al.*, 2023). Feed contributes more than half of production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with appropriate nutrient content can result in good growth and survival (Chepkirui-Boit *et al.*, 2011). The nutritional content of the feed is always related to the materials used (Manam, 2023).

Efforts to obtain cheap and quality feed ingredients continue. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* (Castelnaud, 1855) meal (Cano-Salgado *et al.*, 2021; Hasnidar *et al.*, 2022) and maggot meal (Herawati *et al.*, 2020; Ender Herawati *et al.*, 2019). Both ingredients are expected to replace commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (Panase *et al.*, 2018; Hasnidar *et al.*, 2022). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (Panase *et al.*, 2018); catfish, *Clarias gariepinus* (Mukhlis *et al.* 2020); vaname shrimp, *Litopenaeus vannamei* (Hasnidar *et al.*, 2022). Apart from *P. pardalis* meal; maggot, *Chrysomya megacephala* meal can also be used as a protein source (Siva Raman *et al.*, 2022). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, has a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (Fasakin *et al.*, 2003). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of milkfish, *Chanos chanos* (Herawati *et al.*, 2020); *Salmo salar* (Belghit *et al.*, 2019); Atlantic salmon (Bruni *et al.*, 2020); *Oreochromis niloticus* (Yildirim-Aksoy *et al.*, 2020); *Monopterus albus* (Hu *et al.*, 2020); *Micropterus salmoides* (Peng *et al.*, 2021); *Clarias gariepinus* (Fawole *et al.* 2020).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 46.90% to 47.03% (Unnikrishnan and Paulraj, 2010); 32 to 40% (Catacutan, 2002). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on the source from which it is obtained (Day *et al.*, 2022; Gorissen *et al.*, 2018). In addition to using cheap and quality protein sources, for efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (Kabir *et al.* 2020).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to Suhenda and Samsudin (2008), lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, so that the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to Zainuddin *et al.* (2014), nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth as well as liver degeneration. According to

Catacutan (2002), mud crabs grow well when given feed with a lipid content of 6% or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32% and 8% (Suhenda and Samsudin, 2008), 55% and 7% in golden mandarin fish, *Siniperca scherzei* (Sankian *et al.*, 2018)(Sankian *at al.*, 2007). Furthermore, in white shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth and carbohydrate digestibility (Zainuddin *et al.*, 2014). Information is needed on whether crabs can utilize carbohydrates and lipid in the right ratio as a sparing effect on protein use. This study aims to analyze the use of iso-protein feed with different carbohydrate-lipid ratios on molting, growth and feed efficiency of mud crabs. The results of this study are expected to be useful to encourage the availability of independent and cheap feed and appropriate feed formulations among crab culture.

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023), testing feed chemistry and crabs (May, 2023). The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted, dried for five days. To reduce moisture during drying, the fish was then baked at 70°C for 24 hours (Panase *et al.*, 2018). The dried *P. pardalis* was then ground using a blender, sieved using a 425 µm mesh size sieve. Then the *P. pardalis* meal is ready for use. Maggots were obtained from maggot farming in Makassar. Maggot is cleaned, and dried in the sun until dry. After drying the maggot is oven at 60°C for 12 hours (Alpionita, 2020), then mashed using a blender and sieved using a size 425 µm sieve. Furthermore, maggot meal is ready for use.

Preparation of feed.

Each feed ingredient used is weighed as needed. The ingredients and the amount needed for each treatment are presented in Table 1. The weighed feed ingredients are mixed evenly by stirring manually until the ingredients are homogeneous. Furthermore, a small amount of ingredients such as fish oil, vitamin premix, CMC was added and stirred again. After all the ingredients are evenly mixed, warm water is added to the mixture of feed ingredients until it forms a dough. The feed dough is stirred until it does not stick to the hands, then the dough is molded into pellets using a feed printer. The pelleted feed is then spread evenly on a tray and dried in the sun, then the feed is ready to be applied. The proximate test results of the test feed are presented in Table 2.

Mud crabs, *S. serrata* with a weight of 70±4.7 g/crab, and a carapace length of 5±0.5 cm. Mud crabs were obtained from fishermen collecting in the location of the study site. The culture

container used is a crab box measuring length x width x height, that is 20x15x15 cm as many as 60 pieces, each crab box is filled with 1 crab. At the bottom and sides of the crab box, plastic nets with a mesh size of 1 inch were installed to prevent feed and feed residue from falling through the holes in the crab box.

Furthermore, crab boxes are placed in a row on a maintenance raft made of bamboo, with floats from used plastic bottles. Water changes in the pond were made once a day. Feed with iso-protein composition with different ratio of carbohydrate and lipid (feed formulation for each treatment is listed in Table 1). The crabs were fed 5% of the biomass weight per day with the frequency of feeding 2 a day at 7 am and 6 pm, during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain using digital scales (O-haus) with an accuracy of 0.1 g; carapace width using a sliding mister with an accuracy of 0.1 mm. Measurement of physical and chemical properties of water, i.e. temperature, pH, salinity, dissolved oxygen, ammonia and nitrite. Temperature, pH and salinity were measured daily, dissolved oxygen, ammonia and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on AOAC (Helrich, 1990). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours, crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (Molting Percentage/MP), growth (Weight gain/WG, and Specific Growth Rate/SGR), feed efficiency (Protein Retention/PR, Lipid Retention/LR, Feed Conversion Ratio/FCR and Survival Rate/SR). The formulation used the Boonyaratpalin (1989) equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = \frac{\ln \text{final weight} - \ln \text{initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab weight}) - \text{initial weight}}$$

$$SR = \frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Tables 1 and 2). To determine the effect of treatment on the test parameters, an Analysis of Variance (ANOVA) was performed, followed by Duncan at a significance level of 95% ($p < 0.05$). All data are presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA). The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment with different carbohydrate and lipid ratio showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3). Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased but the lipid and carbohydrate content decreased (Table 4). The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation (Table 5).

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3). The chemical (proximate) test results of the four feed treatments contained 34.49 - 34.97% protein, 30.77 to 35.29% carbohydrate, 6.31 to 9.94% fat. The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters. The protein content of the feed in this study was lower than the needs of mangrove crabs according to (Unnikrishnan and Paulraj, 2010), i.e. 46.90 to 47.03% and 32 to 40% (Catacutan, 2002), so it is thought to cause molting, growth and feed efficiency are not optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to Furuichi and Yone (1982), carbohydrate requirements vary among species, carnivorous species require lower carbohydrates than herbivorous species. Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to Mente (2006), protein functions as a source of energy for growth and tissue repair. Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (Radhakrishnan *et al.*, 2020).

The carbohydrate content of the feed in this study was 30.77 to 35.29%, higher than the crustacean requirement level which is only about 20 to 30% (Wang and Chen, 2016), 18 to 20% in mud crab (Zhang *et al.*, 2022); *Litopenaeus vannamei* (Wang *et al.*, 2014); *Scylla serrata*

(Truong *et al.*, 2008); *Penaeus monodon* (Fabricius) (Alava and Pascual, 1987). The high carbohydrate content in this study aims to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the test parameters. Crab growth is a function of molting (Tamsil and Hasnidar, 2018). The more often a crab molts, the faster its growth. The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to metabolize carbohydrates properly so that molting and growth are less than optimal. The same thing was also stated by (Zhang *et al.*, 2022), that carbohydrate levels of 24.0 to 30.0% in feed negatively affect growth, prolong the molting cycle; and affect crab health (Li *et al.*, 2013). The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. According to Stevens and Jewett (2014), crabs begin physiological preparations for molting about 2 months before releasing their shells. Meanwhile, mangrove crabs grow rapidly through molting occurring every 5-30 days depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth. However, feed with quantity and quality below the optimum will negatively affect growth (Nguyen *et al.*, 2014).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely premolting, molting (ecdysis), postmolting, intermolting, the longest stage is intermolting. If the energy from food is sufficient for growth, the intermolting phase will be faster, and vice versa. Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. When the old carapace is detached (ecdysis), the new carapace will replace it and have a larger size. Therefore, the increase in crab size only occurs during molting. According to Stevens and Jewett (2014), the increase in size of crab every time it molts is around 20% to 25%. Therefore, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions, but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (Tocher, 2003); increase metabolism and respiration in mitochondria (Yuan *et al.*, 2019). Feed lipid levels in this study of 6.31 to 9.94% are levels that can support good crab growth, which is 2 to 10% (Sheen and D'Abramo, 1991; Zhao *et al.*, 2014) or 2 to 12% of the dry weight of food (Catacutan, 2002). The lipid requirements of organisms vary widely depending on species and environmental conditions (Holme *et al.*, 2007). According to Bowser and Rosemark (1981), if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*; inhibit the development of *Marsupenaeus japonicus* larvae (Teshima *et al.*, 1986). Furthermore, high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (Sun *et al.*, 2020). Lipid metabolism in crabs still needs to be studied more deeply because there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells and daily metabolic processes. The results showed that the treatment had no effect on protein retention. Factors that affect protein retention include size, water temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (Suhenda *et al.*, 2004; Jobling, 2012). Because of the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components including sex hormones, and molting activity (Sheen and Wu, 2003); growth and survival (Aaqillah-amr *et al.*, 2021); reproduction and egg quality (Islam and Yahya, 2016). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.13; lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5 (Hastuti *et al.*, 2016); 2.57 to 3.60 (Hudita *et al.*, 2020). The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals. The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth. The feed given must have a compact texture and not easily crushed. The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, this result is higher than the research reported by Hastuti *et al.* (2016) which is 0 to 83%; 63 to 80% (Yulianto *et al.*, 2019). This difference is thought to be due to several factors including treatment, internal and external factors of the test animals. The high survival rate of the results of this study is thought to be because the feed provided produces enough energy to sustain the life of the crab. Environmental conditions, especially salinity, are at a less than optimal level, but mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5), showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. This is thought to be because crabs are less able to metabolize carbohydrates compared to fats. Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. This is thought to be because the energy from carbohydrates and lipids is used to adapt to the non-optimal physical-chemical

environment of water, especially salinity (Table 6). According to Tseng and Hwang (2008) and Wang et al. (2012), carbohydrates can fulfill the high energy needs of aquatic animals during stressful conditions. Furthermore Wang et al. (2014), optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vanname shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth and feed efficiency of mud crabs. However, there is a trend that the treatment with a carbohydrate and lipid ratio of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, 35 and 6%. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

REFERENCES

- Aaqillah-amr M.A., Hidir A., Azra M.N., Noordiyana M.N.** 2021. Interactions between food , feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava V.R., Pascual F.P.** 1987. Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita P.** 2020. Performance of oven shelf type drying on BSF larva. Sriwijaya University.
- Apine E., Ramappa P., Bhatta R., et al.** 2023. Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit I., Liland N.S., Gjesdal P., et al.** 2019. Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December): 609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin M.** 1989. Methodologies for vitamin requirement studies. Pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser P.R., Rosemark R.** 1981. Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.
- Bruni L., Belghit I., Lock E.J., et al.** 2020. Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado M.P., Álvarez-González C.A., Martínez-García R., et al.** 2021. *Pterygoplichthys* spp. (siluriformes: Loricariidae) meal is suitable for the culture of nile tilapia *Oreochromis niloticus* (cichlidae) juveniles. *Ciencia Rural* **52** (1): 1–10. DOI: 10.1590/0103-8478cr20200801
- Catacutan M.R.** 2002. Growth and body composition of juvenile mud crab, *Scylla serrata*, fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8

- Cerbule K., Grimsno L., Herrmann B., Grimaldo E.** 2023. Increasing sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820
- Chepkirui-Boit V., Ngugi C.C., Bowman J., et al.** 2011. Growth performance, survival, feed utilization and nutrient utilization of African catfish (*Clarias gariepinus*) larvae co-fed Artemia and a micro-diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning. *Aquaculture Nutrition* **17** (2). DOI: 10.1111/j.1365-2095.2009.00737.x
- Day L., Cakebread J.A., Loveday S.M.** 2022. Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Endar Herawati V., Susilo A., Hutabarat J., et al.** 2019. Optimization of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) for Growth and Feed Utilization Efficiency of Juvenile *Litopenaeus Vannamei*. *Biotech. Env. Sc* **21** (2): 284–297.
- Fasakin E.A., Balogun A.M., Ajayi O.O.** 2003. Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole F.J., Adeoye A.A., Tiamiyu L.O., et al.** 2020. Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518** (October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi M., Yone Y.** 1982. Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi (Japanese Edition)* **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen S.H.M., Crombag J.J.R., Senden J.M.G., et al.** 2018. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* **50** (12): 1685–1695. DOI: 10.1007/s00726-018-2640-5
- Hasnidar H., Tamsil A., Saenong M., et al.** 2022. Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti Y.P., Nadeak H., Affandi R., Faturrohman K.** 2016. Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati V.E., Pinandoyo, Windarto S., et al.** 2020. Maggot meal (*Hermetia illucens*) substitution on fish meal to growth performance, and nutrient content of milkfish (*Chanos Chanos*). *HAYATI Journal of Biosciences* **27** (2): 154–165. DOI: 10.4308/hjb.27.2.154
- Holme M.-H., Southgate P.C., Zeng C.** 2007. Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme M.H., Zeng C., Southgate P.C.** 2009. A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu Y., Huang Y., Tang T., et al.** 2020. Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345

- Hudita K., Agustoni, Lokapirnasari W.P.** 2020. Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) addition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam M.L., Yahya K.** 2016. Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam T., Saha D., Bhowmik S., et al.** 2022. Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling M.** 2012. National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir K.A., Verdegem M.C.J., Verreth J.A.J., et al.** 2020. Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan C.P.** 1999. Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. *ACIAR Proceedings*.
- Li X.F., Wang Y., Liu W.B., et al.** 2013. Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell R.T., Cho C.Y., Cowey C.B., et al.** 1993. Nutrient Requirements of Fish. DOI: 10.17226/2115
- Manam V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente E.** 2006. Protein nutrition in crustaceans. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **1** (043): 1–7. DOI: 10.1079/PAVSNR20061043
- Mukhlis, Lausu, Majid M.** 2020. Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. *Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3*.
- Nguyen N.T.B., Chim L., Lemaire P., Wantiez L.** 2014. Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI: 10.1016/j.aquaculture.2014.09.014
- Panase P., Uppapong S., Tuncharoen S., et al.** 2018. Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekong giant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng K., Mo W., Xiao H., et al.** 2021. Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan G., Shivkumar, Mannur V.S., et al.** 2020. Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.

- Sankian Z., Khosravi S., Kim Y.O., Lee S.M.** 2018. Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen S.S., D'Abramo L.R.** 1991. Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen S.S., Wu S.W.** 2003. Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forsk., 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva Raman S., Stringer L.C., Bruce N.C., Chong C.S.** 2022. Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens B.G., Jewett S.C.** 2014. Growth, Molting, and Feeding of King Crabs. DOI: 10.1201/b16664-12
- Suhenda N., Samsudin R.** 2008. Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.
- Suhenda N., Tahapari E., Slembrouck J., Moreau Y.** 2004. Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun P., Jin M., Jiao L., et al.** 2020. Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil A., Hasnidar.** 2018. The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima S.I., Kanazawa A., Yamashita M.** 1986. Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher D.R.** 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong P.H., Anderson A.J., Mather P.B., et al.** 2008. Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng Y.C., Hwang P.P.** 2008. Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148** (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan U., Paulraj R.** 2010. Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang X., Li E., Chen L.** 2016. A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang X., Li E., Qin J.G., et al.** 2014. Growth, body composition, and ammonia tolerance of

- juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220
- Wang Y., Li E., Yu N., et al.** 2012. Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. *PLoS ONE* **7** (5): 1–10. DOI: 10.1371/journal.pone.0037316
- Yildirim-Aksoy M., Eljack R., Schrimsher C., Beck B.H.** 2020. Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. *Aquaculture Reports* **17** (May): 100373. DOI: 10.1016/j.aqrep.2020.100373
- Yuan Y., Sun P., Jin M., et al.** 2019. Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Frontiers in Physiology* **10** (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto B., Sunaryo, Taufiq N.S.P., et al.** 2019. Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. *Ecology Environment and Conservation Journal* **25** (july suppl.): 92–101.
- Zainuddin, Haryati, Aslamyah S.** 2014. Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. *Journal of Aquaculture Research & Development* **05** (06). DOI: 10.4172/2155-9546.1000274
- Zhang X., Jin M., Luo J., et al.** 2022. Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Aquaculture Nutrition* **2022**: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao J., Wen X., Li S., et al.** 2014. Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). *Aquaculture Research* **47** (1–13): 887–899. DOI: 10.1111/are.12547

Table 1. Feed ingredients and quantities required for each treatment

Feed ingredients	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix	3	3	3	3
CMC	0.5	0.5	0.5	0.5
Total	100	100	100	100

- 1) Vitamin premix (in 1 kg of diet): vit. A 90.000 IU; vit. D3 30.000 IU; vit. K3 36 mg; vit. E 225 mg; vit. B1 90 mg; vit. B2 135 mg; vit. B6 90 mg; vit. B12 90 mg; vit. C 240 mg; calcium D-pentathenate 120 mg; folic acid 45 mg, biotin 300 mg, inositol 375 mg, nicotinamide 600 mg, cholin chloride 450 mg;
- 2) Mineral premix (in 1 kg of diet): FeCl₃ .4H₂O 1660 mg; ZnSO₄ 100 mg; MnSO₄ 67,5 mg; CuSO₄ 20 mg; KI 1,5 mg, CoSO₄ .7H₂O 1,0 mg; Ca(H₂PO₄) 11.000 mg; MgSO₄ .7H₂O 13.000 mg; K₂HPO₄ .8.000 mg;
- 3) BETN (Extract material without nitrogen) NFE (Nitrogen free extract)

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775)

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test results				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. Polytechnic Tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber. Treatments consisted of

A1 = carbohydrate : lipid ratio of 31% : 10%

A2 = carbohydrate : lipid ratio of 33% : 9%

A3 = carbohydrate : lipid ratio of 34% : 7%

A4 = carbohydrate : lipid ratio of 35% : 6%

MANUSKRIP AWAL

Table 3. Molting, growth and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratio

Parameters	Feed with different carbohydrate and lipid ratio			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	31.67±19.90 ^a	31.33±15.37 ^a	29.67±11.23 ^a	29.0±12.49 ^a
WG (g)	634.40±220.25 ^a	632.13±143.12 ^a	617.00±139.23 ^a	584.67±161.09 ^a
SGR (%)	1.78±0.64 ^a	1.79±0.52 ^a	1.72±0.45 ^a	1.18±0.27 ^a
PR (%)	94±28 ^a	74±29 ^a	56±25 ^a	55±23 ^a
LR (%)	14.59±6.882 ^a	9.78±3.91 ^a	11.01±5.39 ^a	8.17±6.50 ^a
FCR	1.94±0.65 ^a	2.03±0.74 ^a	2.13±0.74 ^a	2.09±1.04 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	73.33±23.09 ^a	73.33±11.55 ^a

Table 4. Chemical/proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) before and after the research.

	Feed with different carbohydrate and lipid ratios	Proximate composition		
		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
Final meat	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

Table 4. Physico-chemical variable, measurement result, optimal level and measuring tools

Physico-chemical vareable	Measurement Results	Optimal Level	Measuring
Temperatur	10 to 30 ⁰ C	29 ⁰ C (Hastuti et al. 2019); 20 to 30 ⁰ C ((Alberts-Hubatsch et al. 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti et al. 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti et al. 2015); 20 ppt ((Suyono 2021)	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman et al. 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti et al. 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti et al. 2019)	Spectrofotometer



Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
23.05
Balas Ke: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
Kepada: hasnidar.yasin@umi.ac.id

29 Desember 2023 pukul

Manuscript ID: EJABF-2312-3892

Manuscript Title: **Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, *Scylla serrata***

Authors: Hasnidar Hasnidar, Andi Tamsil Andi Tamsil, Nursyahrani syahrani

Dear **Dr. Hasnidar Hasnidar**

Evaluation process of the above mentioned manuscript has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s) comments and revise your manuscript within the period of defined time.

Because we are trying to facilitate timely publication of manuscripts submitted to journal, your revised manuscript should be uploaded as soon as possible. If it is not possible for you to submit your revision in a reasonable amount of time, we may have to consider your paper as a new submission.

Once again, thank you for submitting your manuscript to this journal and I look forward to receiving your revision.

Truly yours,

Editorial Office of **Egyptian Journal of Aquatic Biology and Fisheries**

Dear Author,

Please use the attached format template of the Journal (that is present in the Guide for Authors section) after correcting your manuscript and stick to all instructions to save time on publishing, and send it to mtkhalil52@hotmail.com, and put the number of the article.

please save it in word 97/2003 Regards,

EJABF

Reviewers Recommendation:

Reviewer 1:

File Sent by Reviewer:

https://ejabf.journals.ekb.eg/jufile?__file=XDkBIz63hpLvAeevQR3GwhlZ8xqM6_4U4RrIJAXHZGI7e2CAARcCp2BOttdvFGPLy2f40NLoK_k_JQ5dNyT3vtp7IC_St_KA4OhqqJC1sMouHXOy8nSYAThasm5vvZd8oiN4NqLoZA2Of.EAlksfvgmyX5samYZRxBXx0xdoQ0ni3A20kqKyp1kQqkqNryztH7KPs_alqaydugV5K8Dw--https://ejabf.journals.ekb.eg/jufile?__file=YTQhw67AxX4BHm.wGdtMYSkTaFWHaVUj8gDjJ0X2QTLE8BrL6B6FUvatYjAwG2c6.GggXocSPwBV7.DyzNH0YTT6QY0SxdnZKuej8hITyHn7Tb4rE9pEmbsZ_FZiF2EBE4hrH8b00DPMHJX80BxqvsWerc1LMXPY0rxesNUOObjb2Dhgszu6LI5YnBaEagMXCSOKwqsucRJkoiQc.xsbwTQ—

Reviewer Comment For Author:

Dear Editor-in-Chief,

Manuscript ID: EJABF-2312-3892

Manuscript Title: **Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, *Scylla serrata***

In general, this article is acceptable for publication after minor revision (please see the attached file containing the reviewer comments and modifications that aimed to improve the quality of the manuscript).

Manuscript Evaluation Form:

https://ejabf.journals.ekb.eg/author?__au=xnGb9JzZftL95xt9NMlaBLKBgpPm4xwesPp6E7MEITXmAEssotZG7Xli.Jb8xRhU

TABLE REVIEWER COMMENTS

Table 1. Feed ingredients and quantities required for each treatment.

Feed ingredients	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix ¹	3	3	3	3
CMC ²	0.5	0.5	0.5	0.5
Total	100	100	100	100

1) Vitamin premix (in one kg of diet): vit. A 90.000 IU; vit. D3 30.000 IU; vit. K3 36 mg; vit. E 225 mg; vit. B1 90 mg; vit. B2 135 mg; vit. B6 90 mg; vit. B12 90 mg; vit. C 240 mg; calcium D-pentathenate 120 mg; folic acid 45 mg, biotin 300 mg, inositol 375 mg, nicotinamide 600 mg, cholin chloride 450 mg;

2) Carboxymethylcellulose.

Commented [H1]: Any abbreviation in table should be identified under table.

Commented [n2R1]: already fixed

TABLE REVIEWER COMMENTS

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775).

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test results				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. Polytechnic Tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber.

TABLE REVIEWER COMMENTS

Table 3. Molting, growth and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratio.

Parameters	Feed with different carbohydrate and lipid ratio			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	33.33±11.55 ^a	33.33±11.55 ^a	26.67±11.55 ^a	26.67±11.55 ^a
WG (g)	158.18±2.94 ^a	158.03±3.63 ^a	155.25±2.65 ^a	153.54±4.05 ^a
SGR (%)	1.79±0.14 ^a	1.78±0.05 ^a	1.72±0.41 ^a	1.71±0.04 ^a
PR (%)	17.98±0.49 ^a	16.74±0.89 ^a	16.43±0.89 ^a	16.31±0.38 ^a
LR (%)	9.19±0.10 ^a	8.97±0.06 ^a	8.55±0.71 ^a	8.65±0.49 ^a
FCR	1.94±0.16 ^a	2.03±0.14 ^a	2.09±0.12 ^a	2.10±1.09 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	86.67±11.55 ^a	73.33±11.55 ^a

- MP (molting percentage), WG (weight gain), SGR (specific growth rate), PR (protein retention), LR (lipid retention), FCR (feed conversion ratio), SR (survival rate)
- The results of the analysis of variance showed that the treatments did not significantly affect (P>0.05) the test parameters.
- Mean values followed by the same letter in the same row indicate not significantly different (P>0.01) based on Duncan Multiple Range Test (DMRT)

Commented [H3]: Any abbreviation in table should be identified under table.

Commented [n4R3]: has been included below the table

Commented [H5]: Authors should identify under tables, what does the letter "a" mean.

Commented [n6R5]: The results of the analysis of variance showed that the treatments did not significantly affect (P>0.05) the test parameters. Mean values followed by the same letter in the same row indicate not significantly different (P>0.01) based on Duncan Multiple Range Test (DMRT).

TABLE REVIEWER COMMENTS

Table 4. Chemical/proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) before and after the research.

		Proximate composition		
Feed with different carbohydrate and lipid ratios		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
Final meat				
	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

TABLE REVIEWER COMMENTS

Table 5. Physico-chemical variable, measurement result, optimal level and measuring tools.

Physico-chemical variable	Measurement Results	Optimal Level	Measuring
Temperature	10 to 30°C	29°C (Hastuti et al. 2019); 20 to 30°C ((Alberts-Hubatsch et al. 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti et al. 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti et al. 2015); 20 ppt ((Suyono 2021)	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman et al. 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti et al. 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti et al. 2019)	Spectrofotometer

TEXT REVIEWER COMMENTS

Impact of iso-protein feed with ~~different ratio~~ carbohydrate and lipid ~~ratios~~ on growth and feed efficiency of mud crab, *Scylla serrata*

^{1,2}Department of Aquaculture, Faculty of Fisheries and Marine Sciences. Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km. 05 Makassar 90231, South Sulawesi, Indonesia.

ARTICLE INFO

Article History:

Received:
Accepted:
Online:

Keywords:

mud crab
protein
carbohydrate
lipid
ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aimed to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70 ± 4.7 g/head, and a carapace length of 5 ± 0.5 cm were reared using crab boxes and placed in ponds, for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios, i.e.: treatments A (31 and 10%); B (33 and 9%); C (34 and 7%); and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage (/MP); growth (absolute weight growth “WG”, and specific growth rate “SGR”); feed efficiency (protein retention “PR”, lipid retention “LR”, feed conversion ratio “FCR”, and survival “SR”); and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with a different ratios of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment “i.e. 31 and 10%”, is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

Formatted: Justified, Indent: First line: 0 cm

INTRODUCTION

Mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if carried out with low production costs (Apine *et al.*, 2023). Feed contributes more than half of production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with appropriate nutrient content can result in good growth and survival (Chepkirui-Boit

TEXT REVIEWER COMMENTS

et al., 2011). The nutritional content of the feed is always related to the materials used (Manam, 2023).

Efforts to obtain cheap and quality feed ingredients continue. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* (Castelnau, 1855) meal (Cano-Salgado *et al.*, 2021; Hasnidar *et al.*, 2022) and maggot meal (Herawati *et al.*, 2020; Endar Herawati *et al.*, 2019). Both ingredients are expected to replace commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (Panase *et al.*, 2018; Hasnidar *et al.*, 2022). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (Panase *et al.*, 2018); catfish, *Clarias gariepinus* (Mukhlis *et al.* 2020); vaname shrimp, *Litopenaeus vannamei* (Hasnidar *et al.*, 2022). Apart from *P. pardalis* meal; maggot, *Chrysomya megacephala* meal can also be used as a protein source (Siva Raman *et al.*, 2022). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, has a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (Fasakin *et al.*, 2003). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of milkfish, *Chanos chanos* (Herawati *et al.*, 2020); *Salmo salar* (Belghit *et al.*, 2019); Atlantic salmon (Bruni *et al.*, 2020); *Oreochromis niloticus* (Yildirim-Aksoy *et al.*, 2020); *Monopterus albus* (Hu *et al.*, 2020); *Micropterus salmoides* (Peng *et al.*, 2021); *Clarias gariepinus* (Fawole *et al.* 2020).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 46.90% to 47.03% (Unnikrishnan and Paulraj, 2010); 32 to 40% (Catacutan, 2002). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on ~~the source from which it is obtained~~ its source (Day *et al.*, 2022; Gorissen *et al.*, 2018). In addition to using cheap and quality protein sources, for efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (Kabir *et al.* 2020).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to Suhenda and Samsudin (2008), lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, so that the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to Zainuddin *et al.* (2014), nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth as well as liver degeneration. According to Catacutan (2002), mud crabs grow well when given feed with a lipid content of 6% or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32% and 8% (Suhenda and Samsudin, 2008), 55% and 7% in golden

TEXT REVIEWER COMMENTS

mandarin fish, *Siniperca scherzei* (Sankian ~~et al.~~, 2018)(Sankian *et al.*, 2018; (Sankian *at al.*, 2007).-Furthermore, in white shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth and carbohydrate digestibility (Zainuddin *et al.*, 2014). Information is needed on whether crabs can utilize carbohydrates and lipid in the right ratio as a sparing effect on protein use. This study aims to analyze the use of iso-protein feed with different carbohydrate-lipid ratios on molting, growth, and feed efficiency of mud crabs.-The results of this study are expected to be useful to encourage the availability of independent and cheap feed and appropriate feed formulations among crab culture.

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023), testing feed chemistry and crabs (May, 2023).-The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted, dried for five days. -To reduce moisture during drying, the fish was then baked at 70°C for 24 hours (Panase *et al.*, 2018). The dried *P. pardalis* was then ground using a blender, sieved using a 425 µm mesh size sieve. -Then the *P. pardalis* meal is ready for use. -Maggots were obtained from maggot farming in Makassar. Maggot is cleaned, and dried in the sun until dry. After drying the maggot is oven at 60°C for 12 hours (Alpionita, 2020), then mashed using a blender and sieved using a size 425 µm sieve. Furthermore, maggot meal is ready for use.

Preparation of feed.

Each feed ingredient used is weighed as needed. The ingredients and the amount needed for each treatment are presented in Table "1". The weighed feed ingredients are mixed evenly by stirring manually until the ingredients are homogeneous. Furthermore, a small amounts of ingredients such as fish oil, vitamin premix, and CMC was were added and stirred again. After all the ingredients are were evenly mixed, warm water is was added to the mixture of feed ingredients until it forms formed a dough. The feed dough is was stirred until it does did not stick to the hands, then the dough is was molded into pellets using a feed printer. -The pelleted feed is was then spread evenly on a tray and dried in the sun, then the feed is-was ready to be applied.-The proximate test results of the test feed are-were presented in Table "2".

Mud crabs, *S. serrata* with a weight of 70±4.7 g/crab and a carapace length of 5±0.5 cm. Mud crabs were obtained from fishermen collecting in the location of the study site.-The culture container used is was a crab box measuring length × width × height, that is was 20 × 15 × 15 cm as many as 60 pieces, each crab box is-was filled with one+ crab.-At the bottom and sides of the crab box, plastic nets with a mesh size of 4-one inch were installed to prevent feed and feed

Formatted: Space After: 0 pt

Commented [H1]: Any abbreviation should be identified as the 1st appearance.

TEXT REVIEWER COMMENTS

residue from falling through the holes in the crab box.

Furthermore, crab boxes are placed in a row on a maintenance raft made of bamboo, with floats from used plastic bottles. Water changes in the pond were made once a day. Feed with iso-protein composition with different ratios of carbohydrate and lipid (feed formulation for each treatment is listed in Table 1). The crabs were fed 5% of the biomass weight per day with the frequency of feeding 2 a day at 7 am and 6 pm, during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain using digital scales (Ohaus) with an accuracy of 0.1 g; carapace width using a sliding mister with an accuracy of 0.1 mm. Measurement of physical and chemical properties of water, i.e. temperature, pH, salinity, dissolved oxygen, ammonia, and nitrite. Temperature, pH, and salinity were measured daily, dissolved oxygen, ammonia, and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on AOAC (Helrich, 1990). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours, crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (Molting Percentage (%MP)), growth (Weight gain (%WG), and Specific Growth Rate (%SGR)), feed efficiency (Protein Retention (%PR), Lipid Retention (%LR), Feed Conversion Ratio (%FCR) and Survival Rate (%SR)). The formulation used the Boonyaratpalin (1989) equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = -\frac{\ln \text{final weight} - \ln \text{initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab of weight}) - \text{initial weight}}$$

$$SR = -\frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Tables 1 and 2). To

TEXT REVIEWER COMMENTS

determine the effect of treatment on the test parameters, an ~~Analysis~~-analysis of Variance (ANOVA) was performed, followed by Duncan at a significance level of 95% ($p < 0.05$). All data ~~are~~-were presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA).-The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment with different carbohydrate and lipid ratio showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3). Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased, but the lipid and carbohydrate content decreased (Table 4). The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation were shown in-(Table "5")-.

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3).-The chemical (proximate) test results of the four feed treatments contained 34.49 - 34.97% protein, 30.77 to 35.29% carbohydrate, 6.31 to 9.94% fat. -The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters.-The protein content of the feed in this study was lower than the needs of mangrove crabs according to (Unnikrishnan and Paulraj, 2010), i.e. 46.90 to 47.03% and 32 to 40% (Catacutan, 2002), so it is thought to cause molting, growth and feed efficiency are not optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to Furuichi and Yone (1982), carbohydrate requirements vary among species, carnivorous species require lower carbohydrates than herbivorous species.-Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to Mente (2006), protein functions as a source of energy for growth and tissue repair.-Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (Radhakrishnan *et al.*, 2020).

The carbohydrate content of the feed in this study was 30.77 to 35.29%, higher than the crustacean requirement level, which is only about 20 to 30% (Wang and Chen, 2016), 18 to 20% in mud crab (Zhang *et al.*, 2022); *Litopenaeus vannamei* (Wang *et al.*, 2014); *Scylla serrata* (Truong *et al.*, 2008); and *Penaeus monodon* (Fabricius) (Alava and Pascual, 1987). The high carbohydrate content in this study aims to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

TEXT REVIEWER COMMENTS

test parameters.-Crab growth is a function of molting (Tamsil and Hasnidar, 2018). -The more often a crab molts, the faster its growth. -The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to metabolize carbohydrates properly so that molting and growth are less than optimal. The same thing was also stated by (Zhang *et al.*, 2022), that carbohydrate levels of 24.0 to 30.0% in feed negatively affect growth, prolong the molting cycle; and affect crab health (Li *et al.*, 2013).-The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. -According to Stevens and Jewett (2014), crabs begin physiological preparations for molting about 2 months before releasing their shells.-Meanwhile, mangrove crabs grow rapidly through molting occurring every 5-30 days depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth.-However, feed with quantity and quality below the optimum will negatively affect growth (Nguyen *et al.*, 2014).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely premolting, molting (ecdysis), postmolting, intermolting, the longest stage is intermolting. -If the energy from food is sufficient for growth, the intermolting phase will be faster, and vice versa.-Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. -When the old carapace is detached (ecdysis), the new carapace will replace it and have a larger size.-Therefore, the increase in crab size only occurs during molting.-According to Stevens and Jewett (2014), the increase in size of crab every time it molts is around 20% to 25%. -Therefore, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions, but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (Tocher, 2003); increase metabolism and respiration in mitochondria (Yuan *et al.*, 2019). Feed lipid levels in this study of 6.31 to 9.94% are levels that can support good crab growth, which is 2 to 10% (Sheen and D'Abramo, 1991; Zhao *et al.*, 2014) or 2 to 12% of the dry weight of food (Catacutan, 2002). The lipid requirements of organisms vary widely depending on species and environmental conditions (Holme *et al.*, 2007). According to Bowser and Rosemark (1981), if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*; inhibit the development of *Marsupenaeus japonicus* larvae (Teshima *et al.*, 1986).-Furthermore, high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (Sun *et al.*, 2020). -Lipid metabolism in crabs still needs to be studied more deeply because there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells and daily metabolic processes. -The results showed that the treatment had no effect on protein retention. -Factors that affect protein retention include size, water

Formatted: Font: Italic

TEXT REVIEWER COMMENTS

temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (Suhenda *et al.*, 2004; Jobling, 2012). Because of the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components including sex hormones, and molting activity (Sheen and Wu, 2003); growth and survival (Aaqillah-amr *et al.*, 2021); reproduction and egg quality (Islam and Yahya, 2016). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.10; lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5 (Hastuti *et al.*, 2016); 2.57 to 3.60 (Hudita *et al.*, 2020).-The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals.-The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth.

The feed given must have a compact texture and not easily crushed.-The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, this result is higher than the research reported by Hastuti *et al.* (2016) which is 0 to 83%; 63 to 80% (Yulianto *et al.*, 2019). This difference is thought to be due to several factors including treatment, internal and external factors of the test animals.-The high survival rate of the results of this study is thought to be because the feed provided produces enough energy to sustain the life of the crab.-Environmental conditions, especially salinity, are at a less than optimal level, but mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5), showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. -This is thought to be because crabs are less able to metabolize carbohydrates compared to fats. -Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. -This is thought to be because the energy from carbohydrates and lipids is used to adapt to the non-optimal physical-chemical environment of water, especially salinity (Table 65).-According to Tseng and Hwang (2008) and Wang *et al.* (2012), carbohydrates can fulfill the high energy needs of aquatic animals during

TEXT REVIEWER COMMENTS

stressful conditions. Furthermore Wang et al. (2014), optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vanname shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth, and feed efficiency of mud crabs. However, there is a trend that the treatment with a carbohydrate and lipid ratio of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, 35 and 6%. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

REFERENCES

- Aaqillah-amr M.A., Hidir A., Azra M.N., Noordiyana M.N.** 2021. Interactions between food , feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava V.R., Pascual F.P.** 1987. Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita P.** 2020. Performance of oven shelf type drying on BSF larva. Sriwijaya University.
- Apine E., Ramappa P., Bhatta R., et al.** 2023. Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit I., Liland N.S., Gjesdal P., et al.** 2019. Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December): 609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin M.** 1989. Methodologies for vitamin requirement studies. Pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser P.R., Rosemark R.** 1981. Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.
- Bruni L., Belghit I., Lock E.J., et al.** 2020. Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado M.P., Álvarez-González C.A., Martínez-García R., et al.** 2021. *Pterygoplichthys* spp. (siluriformes: Loricariidae) meal is suitable for the culture of Nile tilapia *Oreochromis niloticus* (Cichlidae) juveniles. *Ciencia Rural* **52** (1): 1–10. DOI: 10.1590/0103-8478cr20200801
- Catcutan M.R.** 2002. Growth and body composition of juvenile mud crab, *Scylla serrata*, fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8
- Cerbule K., Grimsø L., Herrmann B., Grimaldo E.** 2023. Increasing sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the Barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820

TEXT REVIEWER COMMENTS

- Chepkirui-Boit V., Ngugi C.C., Bowman J., et al.** 2011. Growth performance, survival, feed utilization and nutrient utilization of African catfish (*Clarias gariepinus*) larvae co-fed Artemia and a micro-diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning. *Aquaculture Nutrition* **17** (2). DOI: 10.1111/j.1365-2095.2009.00737.x
- Day L., Cakebread J.A., Loveday S.M.** 2022. Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Endar Herawati V., Susilo A., Hutabarat J., et al.** 2019. Optimization of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) for Growth and Feed Utilization Efficiency of Juvenile *Litopenaeus Vannamei*. *Biotech. Env. Sc* **21** (2): 284–297.
- Fasakin E.A., Balogun A.M., Ajayi O.O.** 2003. Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole F.J., Adeoye A.A., Tiarniyu L.O., et al.** 2020. Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518** (October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi M., Yone Y.** 1982. Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi (Japanese Edition)* **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen S.H.M., Crombag J.J.R., Senden J.M.G., et al.** 2018. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* **50** (12): 1685–1695. DOI: 10.1007/s00726-018-2640-5
- Hasnidar H., Tamsil A., Saenong M., et al.** 2022. Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti Y.P., Nadeak H., Affandi R., Faturrohman K.** 2016. Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati V.E., Pinandoyo, Windarto S., et al.** 2020. Maggot meal (*Hermetia illucens*) substitution on fish meal to growth performance, and nutrient content of milkfish (*Chanos Chanos*). *HAYATI Journal of Biosciences* **27** (2): 154–165. DOI: 10.4308/hjb.27.2.154
- Holme M.-H., Southgate P.C., Zeng C.** 2007. Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme M.H., Zeng C., Southgate P.C.** 2009. A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu Y., Huang Y., Tang T., et al.** 2020. Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345
- Hudita K., Agustoni, Lokapirnasari W.P.** 2020. Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) addition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab

TEXT REVIEWER COMMENTS

- (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam M.L., Yahya K.** 2016. Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam T., Saha D., Bhowmik S., et al.** 2022. Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling M.** 2012. National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir K.A., Verdegem M.C.J., Verreth J.A.J., et al.** 2020. Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan C.P.** 1999. Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. *ACIAR Proceedings*.
- Li X.F., Wang Y., Liu W.B., et al.** 2013. Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell R.T., Cho C.Y., Cowey C.B., et al.** 1993. Nutrient Requirements of Fish. DOI: 10.17226/2115
- Manam V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente E.** 2006. Protein nutrition in crustaceans. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **1** (043): 1–7. DOI: 10.1079/PAVSNNR20061043
- Mukhlis, Lausu, Majid M.** 2020. Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. *Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3*.
- Nguyen N.T.B., Chim L., Lemaire P., Wantiez L.** 2014. Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI: 10.1016/j.aquaculture.2014.09.014
- Panase P., Uppapong S., Tuncharoen S., et al.** 2018. Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekong giant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng K., Mo W., Xiao H., et al.** 2021. Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan G., Shivkumar, Mannur V.S., et al.** 2020. Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.
- Sankian Z., Khosravi S., Kim Y.O., Lee S.M.** 2018. Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant

TEXT REVIEWER COMMENTS

- enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen S.S., D'Abramo L.R.** 1991. Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen S.S., Wu S.W.** 2003. Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forsk., 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva Raman S., Stringer L.C., Bruce N.C., Chong C.S.** 2022. Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens B.G., Jewett S.C.** 2014. Growth, Molting, and Feeding of King Crabs. DOI: 10.1201/b16664-12
- Suhenda N., Samsudin R.** 2008. Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.
- Suhenda N., Tahapari E., Slembrouck J., Moreau Y.** 2004. Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun P., Jin M., Jiao L., et al.** 2020. Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil A., Hasnidar.** 2018. The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima S.I., Kanazawa A., Yamashita M.** 1986. Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher D.R.** 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong P.H., Anderson A.J., Mather P.B., et al.** 2008. Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng Y.C., Hwang P.P.** 2008. Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148** (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan U., Paulraj R.** 2010. Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang X., Li E., Chen L.** 2016. A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang X., Li E., Qin J.G., et al.** 2014. Growth, body composition, and ammonia tolerance of juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220

TEXT REVIEWER COMMENTS

- Wang Y., Li E., Yu N., et al.** 2012. Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. PLoS ONE 7 (5): 1–10. DOI: 10.1371/journal.pone.0037316
- Yildirim-Aksoy M., Eljack R., Schrimsher C., Beck B.H.** 2020. Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. Aquaculture Reports 17 (May): 100373. DOI: 10.1016/j.aqrep.2020.100373
- Yuan Y., Sun P., Jin M., et al.** 2019. Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. Frontiers in Physiology 10 (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto B., Sunaryo, Taufiq N.S.P., et al.** 2019. Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. Ecology Environment and Conservation Journal 25 (july suppl.): 92–101.
- Zainuddin, Haryati, Aslamyah S.** 2014. Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. Journal of Aquaculture Research & Development 05 (06). DOI: 10.4172/2155-9546.1000274
- Zhang X., Jin M., Luo J., et al.** 2022. Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. Aquaculture Nutrition 2022: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao J., Wen X., Li S., et al.** 2014. Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). Aquaculture Research 47 (1–13): 887–899. DOI: 10.1111/are.12547

Editor-in-Chief Comments

Select Manuscript Type

Enter Manuscript Title

→ Add/Remove Authors

✓ Enter Abstract

✓ Enter Keywords

✓ Select Related Subjects

✓ Additional Comments

✓ Attach Files

✓ Cover Letter & Checklist

✓ Finish Submission



Editor-in-Chief Comment for Author

Reviewers Recommendation

Reviewer 1 EJABF-2312-3892 Minor Revision

Reviewer Comment For Author:

Dear Editor-in-Chief,
Manuscript ID: EJABF-2312-3892
Manuscript Title: Impact of iso-protein feed with ratio carbohydrate and lipid on growth and feed efficiency of mud crab, Scylla serrate

In general, this article is acceptable for publication after minor revision (please see the attached file containing the reviewer comments and modifications that aimed to improve the quality of the manuscript).

File Sent by Reviewer:

Manuscript Evaluation Form

-  EJABF-2312-3892-Text-Reviewer Comments.docx
-  EJABF-2312-3892-Tabels -Reviewer Comments.docx

Original Manuscript Manuscript Needs Revision (Minor Revision)
Status

Revision Due Date 2024-01-13

Editor-in-Chief Comment

Next

Impact of iso-protein feed with [different ratio](#) carbohydrate and lipid [ratios](#) on growth and feed efficiency of mud crab, *Scylla serrata*

^{1,2}Department of Aquaculture, Faculty of Fisheries and Marine Sciences. Universitas Muslim Indonesia, Jl. Urip Sumoharjo Km. 05 Makassar 90231, South Sulawesi, Indonesia.

ARTICLE INFO

Article History:

Received:
Accepted:
Online:

Keywords:

mud crab
protein
carbohydrate
lipid
ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aimed to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70±4.7 g/head, and a carapace length of 5±0.5 cm were reared using crab boxes and placed in ponds, for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios, i.e.: treatments A (31 and 10%); B (33 and 9%); C (34 and 7%); and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage (MP); growth (absolute weight growth “WG”, and specific growth rate “SGR”); feed efficiency (protein retention “PR”, lipid retention “LR”, feed conversion ratio “FCR”, and survival “SR”); and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while waterphysico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with a [different ratios](#) of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment “, i.e. 31 and 10%”, is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

Formatted: Justified, Indent: First line: 0 cm

INTRODUCTION

Mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if carried out with low production costs (Apine *et al.*, 2023). Feed contributes more than half of production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with appropriate nutrient content can result in good growth and survival (Chepkirui-Boit

et al., 2011). The nutritional content of the feed is always related to the materials used (Manam, 2023).

Efforts to obtain cheap and quality feed ingredients continue. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* (Castelnau, 1855) meal (Cano-Salgado *et al.*, 2021; Hasnidar *et al.*, 2022) and maggot meal (Herawati *et al.*, 2020; Endar Herawati *et al.*, 2019). Both ingredients are expected to replace commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (Panase *et al.*, 2018; Hasnidar *et al.*, 2022). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (Panase *et al.*, 2018); catfish, *Clarias gariepinus* (Mukhlis *et al.* 2020); vaname shrimp, *Litopenaeus vannamei* (Hasnidar *et al.*, 2022). Apart from *P. pardalis* meal; maggot, *Chrysomya megacephala* meal can also be used as a protein source (Siva Raman *et al.*, 2022). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, has a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (Fasakin *et al.*, 2003). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of milkfish, *Chanos chanos* (Herawati *et al.*, 2020); *Salmo salar* (Belghit *et al.*, 2019); Atlantic salmon (Bruni *et al.*, 2020); *Oreochromis niloticus* (Yildirim-Aksoy *et al.*, 2020); *Monopterus albus* (Hu *et al.*, 2020); *Micropterus salmoides* (Peng *et al.*, 2021); *Clarias gariepinus* (Fawole *et al.* 2020).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 46.90% to 47.03% (Unnikrishnan and Paulraj, 2010); 32 to 40% (Catacutan, 2002). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on ~~the source from which it is obtained~~ [its source](#) (Day *et al.*, 2022; Gorissen *et al.*, 2018). In addition to using cheap and quality protein sources, for efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (Kabir *et al.* 2020).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to Suhenda and Samsudin (2008), lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, so that the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to Zainuddin *et al.* (2014), nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth as well as liver degeneration. According to Catacutan (2002), mud crabs grow well when given feed with a lipid content of 6% or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32% and 8% (Suhenda and Samsudin, 2008), 55% and 7% in golden

mandarin fish, *Siniperca scherzeri* (Sankian *et al.*, 2018)(Sankian *et al.*, 2018; (Sankian *et al.*, 2007). Furthermore, in white shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth and carbohydrate digestibility (Zainuddin *et al.*, 2014). Information is needed on whether crabs can utilize carbohydrates and lipid in the right ratio as a sparing effect on protein use. This study aims to analyze the use of iso-protein feed with different carbohydrate-lipid ratios on molting, growth, and feed efficiency of mud crabs. The results of this study are expected to be useful to encourage the availability of independent and cheap feed and appropriate feed formulations among crab culture.

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023), testing feed chemistry and crabs (May, 2023). The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted, dried for five days. To reduce moisture during drying, the fish was then baked at 70°C for 24 hours (Panase *et al.*, 2018). The dried *P. pardalis* was then ground using a blender, sieved using a 425 µm mesh size sieve. Then the *P. pardalis* meal is ready for use. Maggots were obtained from maggot farming in Makassar. Maggot is cleaned, and dried in the sun until dry. After drying the maggot is oven at 60°C for 12 hours (Alpionita, 2020), then mashed using a blender and sieved using a size 425 µm sieve. Furthermore, maggot meal is ready for use.

Preparation of feed.

Each feed ingredient used is weighed as needed. The ingredients and the amount needed for each treatment are presented in Table "1". The weighed feed ingredients are mixed evenly by stirring manually until the ingredients are homogeneous. Furthermore, a small amounts of ingredients such as fish oil, vitamin premix, and CMC was were added and stirred again. After all the ingredients are were evenly mixed, warm water is was added to the mixture of feed ingredients until it forms formed a dough. The feed dough is was stirred until it does did not stick to the hands, then the dough is was molded into pellets using a feed printer. The pelleted feed is was then spread evenly on a tray and dried in the sun, then the feed is-was ready to be applied. The proximate test results of the test feed are-were presented in Table "2".

Mud crabs, *S. serrata* with a weight of 70±4.7 g/crab and a carapace length of 5±0.5 cm. Mud crabs were obtained from fishermen collecting in the location of the study site. The culture container used is was a crab box measuring length × width × height, that is was 20 × 15 × 15 cm as many as 60 pieces, each crab box is-was filled with one+ crab. At the bottom and sides of the crab box, plastic nets with a mesh size of 1-one inch were installed to prevent feed and feed

Formatted: Space After: 0 pt

Commented [H1]: Any abbreviation should be identified as the 1st appearance.

residue from falling through the holes in the crab box.

Furthermore, crab boxes are placed in a row on a maintenance raft made of bamboo, with floats from used plastic bottles. Water changes in the pond were made once a day. Feed with iso-protein composition with different ratios of carbohydrate and lipid (feed formulation for each treatment is listed in Table 1). The crabs were fed 5% of the biomass weight per day with the frequency of feeding 2 a day at 7 am and 6 pm, during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain using digital scales (Ohaus) with an accuracy of 0.1 g; carapace width using a sliding mister with an accuracy of 0.1 mm. Measurement of physical and chemical properties of water, i.e. temperature, pH, salinity, dissolved oxygen, ammonia, and nitrite. Temperature, pH, and salinity were measured daily, dissolved oxygen, ammonia, and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on AOAC (Helrich, 1990). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours, crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (Molting Percentage (%MP)), growth (Weight gain (%WG), and Specific Growth Rate (%SGR)), feed efficiency (Protein Retention (%PR), Lipid Retention (%LR), Feed Conversion Ratio (FCR) and Survival Rate (%SR)). The formulation used the Boonyaratpalin (1989) equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = \frac{\ln \text{final weight} - \ln \text{initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab of weight}) - \text{initial weight}}$$

$$SR = \frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Tables 1 and 2). To

determine the effect of treatment on the test parameters, an ~~Analysis-analysis~~ of Variance (ANOVA) was performed, followed by Duncan at a significance level of 95% ($p < 0.05$). All data ~~are-were~~ presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA). The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment with different carbohydrate and lipid ratio showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3). Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased, but the lipid and carbohydrate content decreased (Table 4). The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation ~~were shown in~~ (Table "5"). -

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3). ~~The chemical (proximate) test results of the four feed treatments contained 34.49 - 34.97% protein, 30.77 to 35.29% carbohydrate, 6.31 to 9.94% fat. The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters. The protein content of the feed in this study was lower than the needs of mangrove crabs according to (Unnikrishnan and Paulraj, 2010), i.e. 46.90 to 47.03% and 32 to 40% (Catacutan, 2002), so it is thought to cause molting, growth and feed efficiency are not optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to Furuichi and Yone (1982), carbohydrate requirements vary among species, carnivorous species require lower carbohydrates than herbivorous species. Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to Mente (2006), protein functions as a source of energy for growth and tissue repair. Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (Radhakrishnan et al., 2020).~~

The carbohydrate content of the feed in this study was 30.77 to 35.29%, higher than the crustacean requirement level, which is only about 20 to 30% (Wang and Chen, 2016), 18 to 20% in mud crab (Zhang et al., 2022); *Litopenaeus vannamei* (Wang et al., 2014); *Scylla serrata* (Truong et al., 2008); ~~and~~ *Penaeus monodon* (Fabricius) (Alava and Pascual, 1987). The high carbohydrate content in this study aims to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

test parameters. Crab growth is a function of molting (Tamsil and Hasnidar, 2018). The more often a crab molts, the faster its growth. The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to metabolize carbohydrates properly so that molting and growth are less than optimal. The same thing was also stated by (Zhang *et al.*, 2022), that carbohydrate levels of 24.0 to 30.0% in feed negatively affect growth, prolong the molting cycle; and affect crab health (Li *et al.*, 2013). The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. According to Stevens and Jewett (2014), crabs begin physiological preparations for molting about 2 months before releasing their shells. Meanwhile, mangrove crabs grow rapidly through molting occurring every 5-30 days depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth. However, feed with quantity and quality below the optimum will negatively affect growth (Nguyen *et al.*, 2014).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely premolting, molting (ecdysis), postmolting, intermolting, the longest stage is intermolting. If the energy from food is sufficient for growth, the intermolting phase will be faster, and vice versa. Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. When the old carapace is detached (ecdysis), the new carapace will replace it and have a larger size. Therefore, the increase in crab size only occurs during molting. According to Stevens and Jewett (2014), the increase in size of crab every time it molts is around 20% to 25%. Therefore, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions, but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (Tocher, 2003); increase metabolism and respiration in mitochondria (Yuan *et al.*, 2019). Feed lipid levels in this study of 6.31 to 9.94% are levels that can support good crab growth, which is 2 to 10% (Sheen and D'Abramo, 1991; Zhao *et al.*, 2014) or 2 to 12% of the dry weight of food (Catacutan, 2002). The lipid requirements of organisms vary widely depending on species and environmental conditions (Holme *et al.*, 2007). According to Bowser and Rosemark (1981), if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*; inhibit the development of *Marsupenaeus japonicus* larvae (Teshima *et al.*, 1986). Furthermore, high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (Sun *et al.*, 2020). Lipid metabolism in crabs still needs to be studied more deeply because there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells and daily metabolic processes. The results showed that the treatment had no effect on protein retention. Factors that affect protein retention include size, water

Formatted: Font: Italic

temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (Suhenda *et al.*, 2004; Jobling, 2012). Because of the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components including sex hormones, and molting activity (Sheen and Wu, 2003); growth and survival (Aaqillah-amr *et al.*, 2021); reproduction and egg quality (Islam and Yahya, 2016). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.10; lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5 (Hastuti *et al.*, 2016); 2.57 to 3.60 (Hudita *et al.*, 2020). The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals. The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth.

The feed given must have a compact texture and not easily crushed. The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, this result is higher than the research reported by Hastuti *et al.* (2016) which is 0 to 83%; 63 to 80% (Yulianto *et al.*, 2019). This difference is thought to be due to several factors including treatment, internal and external factors of the test animals. The high survival rate of the results of this study is thought to be because the feed provided produces enough energy to sustain the life of the crab. Environmental conditions, especially salinity, are at a less than optimal level, but mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5), showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. This is thought to be because crabs are less able to metabolize carbohydrates compared to fats. Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. This is thought to be because the energy from carbohydrates and lipids is used to adapt to the non-optimal physical-chemical environment of water, especially salinity (Table 65). According to Tseng and Hwang (2008) and Wang *et al.* (2012), carbohydrates can fulfill the high energy needs of aquatic animals during

stressful conditions. Furthermore Wang et al. (2014), optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vanname shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth, and feed efficiency of mud crabs. However, there is a trend that the treatment with a carbohydrate and lipid ratio of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, 35 and 6%. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

REFERENCES

- Aaqillah-amr M.A., Hidir A., Azra M.N., Noordiyana M.N.** 2021. Interactions between food , feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava V.R., Pascual F.P.** 1987. Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita P.** 2020. Performance of oven shelf type drying on BSF larva. Sriwijaya University.
- Apine E., Ramappa P., Bhatta R., et al.** 2023. Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit I., Liland N.S., Gjesdal P., et al.** 2019. Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December):609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin M.** 1989. Methodologies for vitamin requirement studies. Pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser P.R., Rosemark R.** 1981. Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.
- Bruni L., Belghit I., Lock E.J., et al.** 2020. Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado M.P., Álvarez-González C.A., Martínez-García R., et al.** 2021. *Pterygoplichthys* spp. (siluriformes: Loricariidae) meal is suitable for the culture of nile tilapia *Oreochromis niloticus* (cichlidae) juveniles. *Ciencia Rural* **52** (1): 1–10. DOI: 10.1590/0103-8478cr20200801
- Catcutan M.R.** 2002. Growth and body composition of juvenile mud crab, *Scylla serrata*, fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8
- Cerbule K., Grimsø L., Herrmann B., Grimaldo E.** 2023. Increasing sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820

- Chepkirui-Boit V., Ngugi C.C., Bowman J., et al.** 2011. Growth performance, survival, feed utilization and nutrient utilization of African catfish (*Clarias gariepinus*) larvae co-fed *Artemia* and a micro-diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning. *Aquaculture Nutrition* **17** (2). DOI: 10.1111/j.1365-2095.2009.00737.x
- Day L., Cakebread J.A., Loveday S.M.** 2022. Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Endar Herawati V., Susilo A., Hutabarat J., et al.** 2019. Optimization of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) for Growth and Feed Utilization Efficiency of Juvenile *Litopenaeus Vannamei*. *Biotech. Env. Sc* **21** (2): 284–297.
- Fasakin E.A., Balogun A.M., Ajayi O.O.** 2003. Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole F.J., Adeoye A.A., Tihamiyu L.O., et al.** 2020. Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518**(October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi M., Yone Y.** 1982. Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi* (Japanese Edition) **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen S.H.M., Crombag J.J.R., Senden J.M.G., et al.** 2018. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* **50** (12): 1685–1695. DOI: 10.1007/s00726-018-2640-5
- Hasnidar H., Tamsil A., Saenong M., et al.** 2022. Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti Y.P., Nadeak H., Affandi R., Faturrohman K.** 2016. Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati V.E., Pinandoyo, Windarto S., et al.** 2020. Maggot meal (*Hermetia illucens*) substitution on fish meal to growth performance, and nutrient content of milkfish (*Chanos Chanos*). *HAYATI Journal of Biosciences* **27** (2): 154–165. DOI: 10.4308/hjb.27.2.154
- Holme M.-H., Southgate P.C., Zeng C.** 2007. Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme M.H., Zeng C., Southgate P.C.** 2009. A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu Y., Huang Y., Tang T., et al.** 2020. Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345
- Hudita K., Agustoni, Lokapirnasari W.P.** 2020. Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) addition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab

- (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam M.L., Yahya K.** 2016. Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam T., Saha D., Bhowmik S., et al.** 2022. Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling M.** 2012. National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir K.A., Verdegem M.C.J., Verreth J.A.J., et al.** 2020. Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan C.P.** 1999. Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. *ACIAR Proceedings*.
- Li X.F., Wang Y., Liu W.B., et al.** 2013. Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell R.T., Cho C.Y., Cowey C.B., et al.** 1993. Nutrient Requirements of Fish. DOI: 10.17226/2115
- Manam V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente E.** 2006. Protein nutrition in crustaceans. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **1** (043): 1–7. DOI: 10.1079/PAVSNNR20061043
- Mukhlis, Lausu, Majid M.** 2020. Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. *Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3*.
- Nguyen N.T.B., Chim L., Lemaire P., Wantiez L.** 2014. Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI: 10.1016/j.aquaculture.2014.09.014
- Panase P., Uppapong S., Tuncharoen S., et al.** 2018. Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekonggiant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng K., Mo W., Xiao H., et al.** 2021. Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan G., Shivkumar, Mannur V.S., et al.** 2020. Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.
- Sankian Z., Khosravi S., Kim Y.O., Lee S.M.** 2018. Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant

- enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen S.S., D'Abramo L.R.** 1991. Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen S.S., Wu S.W.** 2003. Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forsk., 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva Raman S., Stringer L.C., Bruce N.C., Chong C.S.** 2022. Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens B.G., Jewett S.C.** 2014. Growth, Molting, and Feeding of King Crabs. DOI: 10.1201/b16664-12
- Suhenda N., Samsudin R.** 2008. Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.
- Suhenda N., Tahapari E., Slembrouck J., Moreau Y.** 2004. Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun P., Jin M., Jiao L., et al.** 2020. Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil A., Hasnidar.** 2018. The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima S.I., Kanazawa A., Yamashita M.** 1986. Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher D.R.** 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong P.H., Anderson A.J., Mather P.B., et al.** 2008. Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng Y.C., Hwang P.P.** 2008. Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148** (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan U., Paulraj R.** 2010. Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang X., Li E., Chen L.** 2016. A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang X., Li E., Qin J.G., et al.** 2014. Growth, body composition, and ammonia tolerance of juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220

- Wang Y., Li E., Yu N., et al.** 2012. Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. PLoS ONE 7 (5): 1–10. DOI: 10.1371/journal.pone.0037316
- Yildirim-Aksoy M., Eljack R., Schrimsher C., Beck B.H.** 2020. Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. Aquaculture Reports 17 (May): 100373. DOI: 10.1016/j.aqrep.2020.100373
- Yuan Y., Sun P., Jin M., et al.** 2019. Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. Frontiers in Physiology 10 (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto B., Sunaryo, Taufiq N.S.P., et al.** 2019. Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. Ecology Environment and Conservation Journal 25 (july suppl.): 92–101.
- Zainuddin, Haryati, Aslamyah S.** 2014. Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. Journal of Aquaculture Research & Development 05 (06). DOI: 10.4172/2155-9546.1000274
- Zhang X., Jin M., Luo J., et al.** 2022. Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. Aquaculture Nutrition 2022: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao J., Wen X., Li S., et al.** 2014. Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). Aquaculture Research 47 (1–13): 887–899. DOI: 10.1111/are.12547

Table 1. Feed ingredients and quantities required for each treatment.

Feed ingredients	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix	3	3	3	3
CMC	0.5	0.5	0.5	0.5
Total	100	100	100	100

1) Vitamin premix (in one kg of diet): vit. A 90.000 IU; vit. D3 30.000 IU; vit. K3 36 mg; vit. E 225 mg; vit. B1 90 mg; vit. B2 135 mg; vit. B6 90 mg; vit. B12 90 mg; vit. C 240 mg; calcium D-pentathenate 120 mg; folic acid 45 mg, biotin 300 mg, inositol 375 mg, nicotinamide 600 mg, cholin chloride 450 mg;

2) Mineral premix (in one kg of diet): FeCl₃ .4H₂O 1660 mg; ZnSO₄ 100 mg; MnSO₄ 67,5 mg; CuSO₄ 20 mg; KI 1,5 mg, CoSO₄ .7H₂O 1,0 mg; Ca (H₂PO₄) 11.000 mg; MgSO₄.7H₂O 13.000 mg; K₂HPO₄ 8.000 mg.

3) BETN (extract material without nitrogen), NFE (nitrogen free extract)

Commented [H1]: Any abbreviation in table should be identified under table.

Commented [H2]: These parameters are not found in the table.

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775).

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test results				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. Polytechnic Tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber.

Table 3. Molting, growth and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratio.

Parameters	Feed with different carbohydrate and lipid ratio			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	33.33±11.55 ^a	33.33±11.55 ^a	26.67±11.55 ^a	26.67±11.55 ^a
WG (g)	158.18±2.94 ^a	158.03±3.63 ^a	155.25±2.65 ^a	153.54±4.05 ^a
SGR (%)	1.79±0.14 ^a	1.78±0.05 ^a	1.72±0.41 ^a	1.71±0.04 ^a
PR (%)	17.98±0.49 ^a	16.74±0.89 ^a	16.43±0.89 ^a	16.31±0.38 ^a
LR (%)	9.19±0.10 ^a	8.97±0.06 ^a	8.55±0.71 ^a	8.65±0.49 ^a
FCR	1.94±0.16 ^a	2.03±0.14 ^a	2.09±0.12 ^a	2.10±1.09 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	86.67±11.55 ^a	73.33±11.55 ^a

Commented [H3]: Any abbreviation in table should be identified under table.

Commented [H4]: Authors should identify under tables, what does the letter “a” mean.

Table 4. Chemical/proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) before and after the research.

Feed with different carbohydrate and lipid ratios		Proximate composition		
		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
Final meat	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

Table 5. Physico-chemical variable, measurement result, optimal level and measuring tools.

Physico-chemical variable	Measurement Results	Optimal Level	Measuring
Temperature	10 to 30°C	29°C (Hastuti et al. 2019); 20 to 30°C ((Alberts-Hubatsch et al. 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti et al. 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti et al. 2015); 20 ppt ((Suyono 2021)	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman et al. 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti et al. 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti et al. 2019)	Spectrofotometer

Egyptian Journal of Aquatic Biology & Fisheries
Zoology Department, Faculty of Science,
Ain Shams University, Cairo, Egypt.
ISSN 1110 – 6131
Vol. (–):2312 – 3892 (2023)
www.ejabf.journals.ekb.eg



Impact of iso-protein feed with different carbohydrate and lipid ratios on growth and feed efficiency of mud crab, *Scylla serrata*

Hasnidar^{1*} Andi Tamsil¹, Nursyahran²

1- Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90231

2- Institute of Maritime Technology and Business Balikpapan, Makassar 90245

*Corresponding Author: hasnidar.yasin@umi.ac.id

ARTICLE INFO

Article History:

Received:

Accepted:

Online:

Keywords:

Mud crab, protein
carbohydrate, lipid
ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aims to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70 ± 4.7 g/head, and a carapace length of 5 ± 0.5 cm were reared using crab boxes and placed in ponds, for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios: treatments A (31 and 10%), B (33 and 9%), C (34 and 7%), and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage (MP); growth (absolute weight growth “WG” and specific growth rate “SGR”; feed efficiency (protein retention “PR”, lipid retention “LR”, feed conversion ratio “FCR”, and survival “SR”; and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with different ratios of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment “31 and 10%” is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

INTRODUCTION

Mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if carried out with low production costs (Apine *et al.*, 2023). Feed contributes more than half of production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with appropriate nutrient content can result in good

growth and survival (Chepkirui-Boit *et al.*, 2011). The nutritional content of the feed is always related to the materials used (Manam, 2023).

Efforts to obtain cheap and quality feed ingredients continue. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* (Castelnau, 1855) meal (Cano-Salgado *et al.*, 2021; Hasnidar *et al.*, 2022) and maggot meal (Herawati *et al.*, 2020; Endar Herawati *et al.*, 2019). Both ingredients are expected to replace commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (Panase *et al.*, 2018; Hasnidar *et al.*, 2022). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (Panase *et al.*, 2018); catfish, *Clarias gariepinus* (Mukhlis *et al.* 2020); vaname shrimp, *Litopenaeus vannamei* (Hasnidar *et al.*, 2022). Apart from *P. pardalis* meal; maggot, *Chrysomya megacephala* meal can also be used as a protein source (Siva Raman *et al.*, 2022). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, has a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (Fasakin *et al.*, 2003). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of milkfish, *Chanos chanos* (Herawati *et al.*, 2020); *Salmo salar* (Belghit *et al.*, 2019); Atlantic salmon (Bruni *et al.*, 2020); *Oreochromis niloticus* (Yildirim-Aksoy *et al.*, 2020); *Monopterus albus* (Hu *et al.*, 2020); *Micropterus salmoides* (Peng *et al.*, 2021); *Clarias gariepinus* (Fawole *et al.* 2020).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 46.90% to 47.03% (Unnikrishnan and Paulraj, 2010); 32 to 40% (Catacutan, 2002). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on its source (Day *et al.*, 2022; Gorissen *et al.*, 2018). In addition to using cheap and quality protein sources, for efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (Kabir *et al.* 2020).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to Suhenda and Samsudin (2008), lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, so that the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to Zainuddin *et al.* (2014), nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth as well as liver degeneration. According to Catacutan (2002), mud crabs grow well when given feed with a lipid content of 6% or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32% and 8% (Suhenda and Samsudin, 2008), 55% and 7% in golden mandarin fish, *Siniperca scherzei* (Sankian *et al.*, 2018; Furthermore, in white

shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023), testing feed chemistry and crabs (May, 2023). The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted, dried for five days. To reduce moisture during drying, the fish was then baked at 70°C for 24 hours (Panase *et al.*, 2018). The dried *P. pardalis* was then ground using a blender, sieved using a 425 µm mesh size sieve. Then the *P. pardalis* meal is ready for use. Maggots were obtained from maggot farming in Makassar. Maggot is cleaned, and dried in the sun until dry. After drying the maggot is oven at 60°C for 12 hours (Alpionita, 2020), then mashed using a blender and sieved using a size 425 µm sieve. Furthermore, maggot meal is ready for use.

Preparation of feed.

Each feed ingredient used is weighed as needed. The ingredients and the amount needed for each treatment are presented in Table “1”. The weighed feed ingredients are mixed evenly by stirring manually until the ingredients are homogeneous. Furthermore, small amounts of ingredients such as fish oil, vitamin premix, and Carboxy metil cellulose (CMC) were added and stirred again. After all the ingredients were evenly mixed, warm water was added to the mixture of feed ingredients until it formed a dough. The feed dough was stirred until it did not stick to the hands, then the dough was molded into pellets using a feed printer. The pelleted feed was then spread evenly on a tray and dried in the sun, then the feed was ready to be applied.

Table 1. Feed ingredients and quantities required for each treatment.

Feed ingredients	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix ¹	3	3	3	3
CMC ²	0.5	0.5	0.5	0.5
Total	100	100	100	100

- 1) Vitamin premix (in one kg of diet): vit. A 90.000 IU; vit. D3 30.000 IU; vit. K3 36 mg; vit. E 225 mg; vit. B1 90 mg; vit. B2 135 mg; vit. B6 90 mg; vit. B12 90 mg; vit. C 240 mg; calcium D-pentathenate 120 mg; folic acid 45 mg, biotin 300 mg, inositol 375 mg, nicotinamide 600 mg, cholin chloride 450 mg;
- 2) Carboxymethylcellulose.

The proximate test results of the test feed were presented in Table “2”.

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775).

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test results				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. Polytechnic Tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber.

Mud crabs, *S. serrata* with a weight of 70 ± 4.7 g/crab and a carapace length of 5 ± 0.5 cm, were obtained from fishermen collecting in the location of the study site. The culture container used was a crab box measuring length $\times$ width $\times$ height that was $20 \times 15 \times 15$ cm as many as 60 pieces, each crab box was filled with one crab. At the bottom and sides of the crab box, plastic nets with a mesh size of one inch were installed to prevent feed and feed residue from falling through the holes in the crab box.

Furthermore, crab boxes are placed in a row on a maintenance raft made of bamboo, with floats from used plastic bottles. Water changes in the pond were made once a day. Feed with iso-protein composition with different ratios of carbohydrate and lipid (feed formulation for each treatment is listed in Table “1”). The crabs were fed 5% of the biomass weight per day with the frequency of feeding 2 a day at 7 am and 6 pm, during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain using digital scales (O-haus) with an accuracy of 0.1 g; carapace width using a sliding mister with an accuracy of 0.1 mm. Measurement of physical and chemical properties of water, i.e. temperature, pH, salinity, dissolved oxygen, ammonia, and nitrite. Temperature, pH, and salinity were measured daily, dissolved oxygen, ammonia, and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on AOAC (Helrich, 1990). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours, crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (molting percentage “MP”), growth (weight gain “WG”, and specific growth rate “SGR”), feed efficiency (protein retention “PR”, lipid retention “LR”, feed conversion ratio “FCR”, and survival rate “SR”). The formulation used the Boonyaratpalin (1989) equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = \frac{\ln \text{final weight} - \ln \text{initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab of weight}) - \text{initial weight}}$$

$$SR = \frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Table 1). To determine the effect of treatment on the test parameters, an analysis of Variance (ANOVA) was performed, followed by Duncan Multiple Range Test (DMRT) at a significance level of 95% ($p < 0.05$). All data were presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA). The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment with different carbohydrate and lipid ratio showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3).

Table 3. Molting, growth and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratio.

Parameters	Feed with different carbohydrate and lipid ratio			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	33.33±11.55 ^a	33.33±11.55 ^a	26.67±11.55 ^a	26.67±11.55 ^a
WG (g)	158.18±2.94 ^a	158.03±3.63 ^a	155.25±2.65 ^a	153.54±4.05 ^a
SGR (%)	1.79±0.14 ^a	1.78±0.05 ^a	1.72±0.41 ^a	1.71±0.04 ^a
PR (%)	17.98±0.49 ^a	16.74±0.89 ^a	16.43±0.89 ^a	16.31±0.38 ^a
LR (%)	9.19±0.10 ^a	8.97±0.06 ^a	8.55±0.71 ^a	8.65±0.49 ^a
FCR	1.94±0.16 ^a	2.03±0.14 ^a	2.09±0.12 ^a	2.10±1.09 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	86.67±11.55 ^a	73.33±11.55 ^a

- MP (molting percentage), WG (weight gain), SGR (specific growth rate), PR (protein retention), LR (lipid retention), FCR (feed conversion ratio), SR (survival rate)
- The results of the analysis of variance showed that the treatments did not significantly affect ($P>0.05$) the test parameters.
- Mean values followed by the same letter in the same row indicate not significantly different ($P>0.01$) based on Duncan Multiple Range Test (DMRT)

Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased, but the lipid and carbohydrate content decreased (Table 4).

Table 4. Chemical/proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) initial and final the research.

	Feed with different carbohydrate and lipid ratios	Proximate composition		
		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
Final meat	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation were shown in Table “5”.

Table 5. Physico-chemical variable, measurement result, optimal level and measuring tools.

Physico-chemical variable	Measurement Results	Optimal Level	Measuring
Temperature	10 to 30°C	29°C (Hastuti <i>et al.</i> 2019); 20 to 30°C (Alberts-Hubatsch <i>et al.</i> 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti <i>et al.</i> 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti <i>et al.</i> 2015); 20 ppt ((Suyono 2021)	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman <i>et al.</i> 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti <i>et al.</i> 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti <i>et al.</i> 2019)	Spectrofotometer

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3). The chemical (proximate) test results of the four feed treatments contained 34.49 - 34.97% protein, 30.77 to 35.29% carbohydrate, 6.31 to 9.94% fat. The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters. The protein content of the feed in this study was lower than the needs of

mangrove crabs according to (Unnikrishnan and Paulraj, 2010), i.e. 46.90 to 47.03% and 32 to 40% (Catacutan, 2002), so it is thought to cause molting, growth and feed efficiency are not optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to Furuichi and Yone (1982), carbohydrate requirements vary among species, carnivorous species require lower carbohydrates than herbivorous species. Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to Mente (2006), protein functions as a source of energy for growth and tissue repair. Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (Radhakrishnan *et al.*, 2020).

The carbohydrate content of the feed in this study was 30.77 to 35.29%, higher than the crustacean requirement level, which is only about 20 to 30% (Wang and Chen, 2016), 18 to 20% in mud crab (Zhang *et al.*, 2022); *Litopenaeus vannamei* (Wang *et al.*, 2014); *Scylla serrata* (Truong *et al.*, 2008); and *Penaeus monodon* (Fabricius) (Alava and Pascual, 1987). The high carbohydrate content in this study aims to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the test parameters. Crab growth is a function of molting (Tamsil and Hasnidar, 2018). The more often a crab molts, the faster its growth. The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to metabolize carbohydrates properly so that molting and growth are less than optimal. The same thing was also stated by (Zhang *et al.*, 2022), that carbohydrate levels of 24.0 to 30.0% in feed negatively affect growth, prolong the molting cycle; and affect crab health (Li *et al.*, 2013). The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. According to Stevens and Jewett (2014), crabs begin physiological preparations for molting about 2 months before releasing their shells. Meanwhile, mangrove crabs grow rapidly through molting occurring every 5-30 days depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth. However, feed with quantity and quality below the optimum will negatively affect growth (Nguyen *et al.*, 2014).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely premolting, molting (ecdysis), postmolting, intermolting, the longest stage is intermolting. If the energy from food is sufficient for growth, the intermolting phase will be faster, and vice versa. Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. When the old carapace is detached (ecdysis), the new carapace will replace it and have a larger size. Therefore, the increase in crab size only occurs during molting. According to Stevens and Jewett (2014), the increase in size of crab every time it molts is around 20% to 25%. Therefore, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions, but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (Tocher, 2003); increase metabolism and respiration in mitochondria (Yuan *et al.*, 2019). Feed lipid levels in this study of 6.31 to 9.94% are levels that can support good crab growth, which is 2 to 10% (Sheen and D'Abramo, 1991; Zhao *et al.*, 2014) or 2 to 12% of the dry weight of food (Catacutan, 2002). The lipid requirements of organisms vary widely depending on species and environmental conditions (Holme *et al.*, 2007). According to Bowser and Rosemark (1981), if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*; inhibit the development of *Marsupenaeus japonicus* larvae (Teshima *et al.*, 1986). Furthermore, high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (Sun *et al.*, 2020). Lipid metabolism in crabs still needs to be studied more deeply because there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells and daily metabolic processes. The results showed that the treatment had no effect on protein retention. Factors that affect protein retention include size, water temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (Suhenda *et al.*, 2004; Jobling, 2012). Because of the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components including sex hormones, and molting activity (Sheen and Wu, 2003); growth and survival (Aaqillah-amr *et al.*, 2021); reproduction and egg quality (Islam and Yahya, 2016). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.10; lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5 (Hastuti *et al.*, 2016); 2.57 to 3.60 (Hudita *et al.*, 2020). The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals. The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth. The feed given must have a compact texture and not easily crushed. The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, this result is higher than the research reported by Hastuti *et al.* (2016) which is 0 to 83%; 63 to 80% (Yulianto *et al.*, 2019). This difference is thought to be due to several factors including treatment, internal and external factors of the test animals. The high survival rate of the results of this study is thought to be because the feed provided produces enough energy to sustain the life of the crab. Environmental conditions, especially salinity, are at a less than optimal level, but

mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5), showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. This is thought to be because crabs are less able to metabolize carbohydrates compared to fats. Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. This is thought to be because the energy from carbohydrates and lipids is used to adapt to the non-optimal physical-chemical environment of water, especially salinity (Table 5). According to Tseng and Hwang (2008) and Wang *et al.* (2012), carbohydrates can fulfill the high energy needs of aquatic animals during stressful conditions. Furthermore Wang *et al.* (2014), optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vanname shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth, and feed efficiency of mud crabs. However, there is a trend that the treatment with a carbohydrate and lipid ratio of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, 35 and 6%. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

REFERENCES

- Aaqillah-amr M.A., Hidir A., Azra M.N., Noordiyana M.N.** 2021. Interactions between food , feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava V.R., Pascual F.P.** 1987. Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita P.** 2020. Performance of oven shelf type drying on BSF larva. Sriwijaya Univercity.
- Apine E., Ramappa P., Bhatta R., et al.** 2023. Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit I., Liland N.S., Gjesdal P., et al.** 2019. Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December): 609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin M.** 1989. Methodologies for vitamin requirement studies. Pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser P.R., Rosemark R.** 1981. Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.

- Bruni L., Belghit I., Lock E.J., et al.** 2020. Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado M.P., Álvarez-González C.A., Martínez-García R., et al.** 2021. *Pterygoplichthys* spp. (siluriformes: Loricariidae) meal is suitable for the culture of nile tilapia *oreochromis niloticus* (cichlidae) juveniles. *Ciencia Rural* **52** (1): 1–10. DOI: 10.1590/0103-8478cr20200801
- Catacutan M.R.** 2002. Growth and body composition of juvenile mud crab, *Scylla serrata*, fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8
- Cerbule K., Grimsmo L., Herrmann B., Grimaldo E.** 2023. Increasing sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820
- Chepkirui-Boit V., Ngugi C.C., Bowman J., et al.** 2011. Growth performance, survival, feed utilization and nutrient utilization of African catfish (*Clarias gariepinus*) larvae co-fed *Artemia* and a micro-diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning. *Aquaculture Nutrition* **17** (2). DOI: 10.1111/j.1365-2095.2009.00737.x
- Day L., Cakebread J.A., Loveday S.M.** 2022. Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Endar Herawati V., Susilo A., Hutabarat J., et al.** 2019. Optimization of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) for Growth and Feed Utilization Efficiency of Juvenile *Litopenaeus Vannamei*. *Biotech. Env. Sc* **21** (2): 284–297.
- Fasakin E.A., Balogun A.M., Ajayi O.O.** 2003. Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole F.J., Adeoye A.A., Tiamiyu L.O., et al.** 2020. Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518** (October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi M., Yone Y.** 1982. Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi (Japanese Edition)* **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen S.H.M., Crombag J.J.R., Senden J.M.G., et al.** 2018. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* **50** (12): 1685–1695. DOI: 10.1007/s00726-018-2640-5
- Hasnidar H., Tamsil A., Saenong M., et al.** 2022. Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti Y.P., Nadeak H., Affandi R., Faturrohman K.** 2016. Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati V.E., Pinandoyo, Windarto S., et al.** 2020. Maggot meal (*Hermetia illucens*) substitution on fish meal to growth performance, and nutrient content of milkfish (*Chanos Chanos*). *HAYATI Journal of Biosciences* **27** (2): 154–165. DOI: 10.4308/hjb.27.2.154

- Holme M.-H., Southgate P.C., Zeng C.** 2007. Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme M.H., Zeng C., Southgate P.C.** 2009. A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu Y., Huang Y., Tang T., et al.** 2020. Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345
- Hudita K., Agustoni, Lokapirnasari W.P.** 2020. Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) addition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam M.L., Yahya K.** 2016. Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam T., Saha D., Bhowmik S., et al.** 2022. Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling M.** 2012. National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir K.A., Verdegem M.C.J., Verreth J.A.J., et al.** 2020. Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan C.P.** 1999. Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. ACIAR Proceedings.
- Li X.F., Wang Y., Liu W.B., et al.** 2013. Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell R.T., Cho C.Y., Cowey C.B., et al.** 1993. Nutrient Requirements of Fish. DOI: 10.17226/2115
- Manam V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente E.** 2006. Protein nutrition in crustaceans. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **1** (043): 1–7. DOI: 10.1079/PAVSNNR20061043
- Mukhlis, Lausu, Majid M.** 2020. Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3.
- Nguyen N.T.B., Chim L., Lemaire P., Wantiez L.** 2014. Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI:

10.1016/j.aquaculture.2014.09.014

- Panase P., Uppapong S., Tuncharoen S., et al.** 2018. Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekong giant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng K., Mo W., Xiao H., et al.** 2021. Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan G., Shivkumar, Mannur V.S., et al.** 2020. Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.
- Sankian Z., Khosravi S., Kim Y.O., Lee S.M.** 2018. Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen S.S., D'Abramo L.R.** 1991. Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen S.S., Wu S.W.** 2003. Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forsk., 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva Raman S., Stringer L.C., Bruce N.C., Chong C.S.** 2022. Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens B.G., Jewett S.C.** 2014. Growth, Molting, and Feeding of King Crabs. DOI: 10.1201/b16664-12
- Suhenda N., Samsudin R.** 2008. Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.
- Suhenda N., Tahapari E., Slembrouck J., Moreau Y.** 2004. Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun P., Jin M., Jiao L., et al.** 2020. Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil A., Hasnidar.** 2018. The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima S.I., Kanazawa A., Yamashita M.** 1986. Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher D.R.** 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong P.H., Anderson A.J., Mather P.B., et al.** 2008. Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng Y.C., Hwang P.P.** 2008. Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148**

- (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan U., Paulraj R.** 2010. Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang X., Li E., Chen L.** 2016. A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang X., Li E., Qin J.G., et al.** 2014. Growth, body composition, and ammonia tolerance of juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220
- Wang Y., Li E., Yu N., et al.** 2012. Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. *PLoS ONE* **7** (5): 1–10. DOI: 10.1371/journal.pone.0037316
- Yildirim-Aksoy M., Eljack R., Schrimsher C., Beck B.H.** 2020. Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. *Aquaculture Reports* **17** (May): 100373. DOI: 10.1016/j.aqrep.2020.100373
- Yuan Y., Sun P., Jin M., et al.** 2019. Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Frontiers in Physiology* **10** (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto B., Sunaryo, Taufiq N.S.P., et al.** 2019. Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. *Ecology Environment and Conservation Journal* **25** (july suppl.): 92–101.
- Zainuddin, Haryati, Aslamyah S.** 2014. Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. *Journal of Aquaculture Research & Development* **05** (06). DOI: 10.4172/2155-9546.1000274
- Zhang X., Jin M., Luo J., et al.** 2022. Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Aquaculture Nutrition* **2022**: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao J., Wen X., Li S., et al.** 2014. Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). *Aquaculture Research* **47** (1–13): 887–899. DOI: 10.1111/are.12547



Pada Sen, 26 Feb 2024 pukul 21.30 Magdy Khalil <mtkhalil52@hotmail.com> menulis:

Dear Authors,

Please revise carefully the attached version of your article before Language editing and publishing, especially the references, to match that of the Journal;

Example

Van der Geer, J.; Hanraads, J.A.J. ; Khalil, M. and Lupton, R.A. (2010). The art of writing a scientific article. J. Sci. Commun., 163: 51–59.

et al. should be written italic

Line space of the text is 1.15

Correct the chronological order of the references inside the text;

Please Send the cost of publication (\$160) to Magdy Tawfik Khalil Ibrahim

Commercial International Bank (CIB)

Address: Thawra Branch- Cairo, Egypt

Account No. 100016289684

SWIFT CODE: CIBEEGXX081

IBAN: EG450010000600000100016289684

Beneficial address: Faculty of Science, Ain Shams University, Cairo, Egypt

Please write your full name in the receipt of the Bank

Egyptian Journal of Aquatic Biology & Fisheries
 Zoology Department, Faculty of Science,
 Ain Shams University, Cairo, Egypt.
 ISSN 1110 – 6131
 Vol. 28 (1): (2024)
www.ejabf.journals.ekb.eg



Impact of iso-protein feed with different carbohydrate and lipid ratios on growth and feed efficiency of mud crab, *Scylla serrata*

Hasnidar^{1*}, Andi Tamsil¹, Nursyahrani²

1- Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90231

2- Institute of Maritime Technology and Business Balikpapan, Makassar 90245

*Corresponding Author: hasnidar.yasin@umi.ac.id

ARTICLE INFO

Article History:

Received: Dec. 22, 2023

Accepted: Feb. 25, 2024

Online:

Keywords:

Mud crab, protein
 carbohydrate, lipid
 ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aims to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70 ± 4.7 g/head, and a carapace length of 5 ± 0.5 cm were reared using crab boxes and placed in ponds, for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios: treatments A (31 and 10%), B (33 and 9%), C (34 and 7%), and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage (MP); growth (absolute weight growth "WG" and specific growth rate "SGR"; feed efficiency (protein retention "PR", lipid retention "LR", feed conversion ratio "FCR", and survival "SR"; and physical-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with different ratios of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment "31 and 10%" is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

INTRODUCTION

The Mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if carried out with low production costs (Apine *et al.*, 2023). Feed contributes more than half of production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with appropriate nutrient content can result in good

growth and survival (**Chepkirui-Boit *et al.*, 2011**). The nutritional content of the feed is always related to the materials used (**Manam, 2023**).

Efforts to obtain cheap and quality feed ingredients continue. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* (**Castelnau, 1855**) meal (**Cano-Salgado *et al.*, 2022**; **Hasnidar *et al.*, 2022**) and maggot meal (**Herawati *et al.*, 2019**; **Herawati *et al.*, 2020**). Both ingredients are expected to replace commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (**Panase *et al.*, 2018**; **Hasnidar *et al.*, 2022**). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (**Panase *et al.*, 2018**). catfish, *Clarias gariepinus* (**Mukhlis *et al.*, 2020**) vaname shrimp, *Litopenaeus vannamei* (**Hasnidar *et al.*, 2022**). Apart from *P. pardalis* meal; maggot, *Chrysomya megacephala* meal can also be used as a protein source (**Siva Raman *et al.*, 2022**). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, has a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (**Fasakin *et al.*, 2003**). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of milkfish, *Chanos chanos* (**Herawati *et al.*, 2020**); *Salmo salar* (**Belghit *et al.*, 2019**); Atlantic salmon (**Bruni *et al.*, 2020**); *Oreochromis niloticus* (**Yildirim-Aksoy *et al.*, 2020**); *Monopterus albus* (**Hu *et al.*, 2020**); *Micropterus salmoides* (**Peng *et al.*, 2021**); *Clarias gariepinus* (**Fawole *et al.*, 2020**).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 46.90% to 47.03% (**Unnikrishnan and Paulraj, 2010**); 32 to 40% (**Catacutan, 2002**). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on its source (**Gorissen *et al.*, 2018**; **Day *et al.*, 2022**). In addition to using cheap and quality protein sources, for efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (**Kabir *et al.*, 2020**).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to **Suhenda and Samsudin (2008)**, lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, so that the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to **Zainuddin *et al.*, (2014)**, nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth as well as liver degeneration. According to **Catacutan (2002)**, mud crabs grow well when given feed with a lipid content of 6% or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32% and 8% (**Suhenda and Samsudin, 2008**), 55% and 7% in golden mandarin fish, *Siniperca scherzei* (**Sankian *et al.*, 2018**), Furthermore, in white

shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023), testing feed chemistry and crabs (May, 2023). The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted, dried for five days. To reduce moisture during drying, the fish was then baked at 70°C for 24 hours (**Panase et al., 2018**). The dried *P. pardalis* was then ground using a blender, sieved using a 425 µm mesh size sieve. Then the *P. pardalis* meal is ready for use. Maggots were obtained from maggot farming in Makassar. Maggot is cleaned, and dried in the sun until dry. After drying the maggot is oven at 60°C for 12 hours (**Alpionita, 2020**), then mashed using a blender and sieved using a size 425 µm sieve. Furthermore, maggot meal is ready for use.

Preparation of feed.

Each feed ingredient used is weighed as needed. The ingredients and the amount needed for each treatment are presented in Table “1”. The weighed feed ingredients are mixed evenly by stirring manually until the ingredients are homogeneous. Furthermore, small amounts of ingredients such as fish oil, vitamin premix, and Carboxy metil cellulose (CMC) were added and stirred again. After all the ingredients were evenly mixed, warm water was added to the mixture of feed ingredients until it formed a dough. The feed dough was stirred until it did not stick to the hands, then the dough was molded into pellets using a feed printer. The pelleted feed was then spread evenly on a tray and dried in the sun, then the feed was ready to be applied. The proximate test results of the test feed were presented in Table “2”.

Table 1. Feed ingredients and quantities required for each treatment.

Feed ingredients	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix ¹	3	3	3	3
CMC ²	0.5	0.5	0.5	0.5
Total	100	100	100	100

1) Vitamin premix (in one kg of diet): vit. A 90.000 IU; vit. D3 30.000 IU; vit. K3 36 mg; vit. E 225 mg; vit. B1 90 mg; vit. B2 135 mg; vit. B6 90 mg; vit. B12 90 mg; vit. C 240 mg; calcium D-pentathenate 120 mg; folic acid 45 mg, biotin 300 mg, inositol 375 mg, nicotinamide 600 mg, cholin chloride 450 mg;

2) Carboxymethylcellulose.

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775)

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test results				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. Polytechnic Tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber.

Mud crabs, *S. serrata* with a weight of 70 ± 4.7 g/crab and a carapace length of 5 ± 0.5 cm, were obtained from fishermen collecting in the location of the study site. The culture container used was a crab box measuring length $\times$ width $\times$ height that was $20 \times 15 \times 15$ cm as many as 60 pieces, each crab box was filled with one crab. At the bottom and sides of the crab box, plastic nets with a mesh size of one inch were installed to prevent feed and feed residue from falling through the holes in the crab box.

Furthermore, crab boxes are placed in a row on a maintenance raft made of bamboo, with floats from used plastic bottles. Water changes in the pond were made once a day. Feed with iso-protein composition with different ratios of carbohydrate and lipid (feed formulation for each treatment is listed in Table “1”). The crabs were fed 5% of the biomass weight per day with the frequency of feeding 2 a day at 7 am and 6 pm, during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth

sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain using digital scales (O-haus) with an accuracy of 0.1 g; carapace width using a sliding mister with an accuracy of 0.1 mm. Measurement of physical and chemical properties of water, i.e. temperature, pH, salinity, dissolved oxygen, ammonia, and nitrite. Temperature, pH, and salinity were measured daily, dissolved oxygen, ammonia, and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on AOAC (Helrich, 1990). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours, crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (molting percentage “MP”), growth (weight gain “WG”, and specific growth rate “SGR”), feed efficiency (protein retention “PR”, lipid retention “LR”, feed conversion ratio “FCR”, and survival rate “SR”). The formulation used the Boonyaratpalin (1989) equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = \frac{\text{Ln final weight} - \text{Ln initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab of weight}) - \text{initial weight}}$$

$$SR = \frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Table 1). To determine the effect of treatment on the test parameters, an analysis of Variance (ANOVA) was performed, followed by Duncan Multiple Range Test (DMRT) at a significance level of 95% ($p < 0.05$). All data were presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA). The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment with different carbohydrate and lipid ratio showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3).

Table 3. Molting, growth and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratio.

Parameters	Feed with different carbohydrate and lipid ratio			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	33.33±11.55 ^a	33.33±11.55 ^a	26.67±11.55 ^a	26.67±11.55 ^a
WG (g)	158.18±2.94 ^a	158.03±3.63 ^a	155.25±2.65 ^a	153.54±4.05 ^a
SGR (%)	1.79±0.14 ^a	1.78±0.05 ^a	1.72±0.41 ^a	1.71±0.04 ^a
PR (%)	17.98±0.49 ^a	16.74±0.89 ^a	16.43±0.89 ^a	16.31±0.38 ^a
LR (%)	9.19±0.10 ^a	8.97±0.06 ^a	8.55±0.71 ^a	8.65±0.49 ^a
FCR	1.94±0.16 ^a	2.03±0.14 ^a	2.09±0.12 ^a	2.10±1.09 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	86.67±11.55 ^a	73.33±11.55 ^a

- MP (molting percentage), WG (weight gain), SGR (specific growth rate), PR (protein retention), LR (lipid retention), FCR (feed conversion ratio), SR (survival rate)
- The results of the analysis of variance showed that the treatments did not significantly affect ($P>0.05$) the test parameters.
- Mean values followed by the same letter in the same row indicate not significantly different ($P>0.01$) based on Duncan Multiple Range Test (DMRT)

Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased, but the lipid and carbohydrate content decreased (Table 4).

Table 4. Chemical/proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) initial and final the research.

	Feed with different carbohydrate and lipid ratios	Proximate composition		
		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
Final meat	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation were shown in Table “5”.

Table 5. Physico-chemical variable, measurement result, optimal level and measuring tools.

Physico-chemical Variable	Measurement Results	Optimal Level	Measuring
Temperature	10 to 30°C	29°C (Hastuti et al., 2019); 20 to 30°C (Alberts-Hubatsch et al., 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti et al., 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti et al., 2015); 20 ppt ((Suyono, 2021))	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman et al., 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti et al., 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti et al., 2019)	Spectrofotometer

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3). The chemical (proximate) test results of the four feed treatments contained 34.49 - 34.97% protein, 30.77 to 35.29% carbohydrate, 6.31 to 9.94% fat. The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters. The protein content of the feed in this study was lower than the needs of mangrove crabs according to (**Unnikrishnan and Paulraj, 2010**), i.e. 46.90 to 47.03% and 32 to 40% (**Catacutan, 2002**), so it is thought to cause molting, growth and feed efficiency are not optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to **Furuichi and Yone (1982)**, carbohydrate requirements vary among species, carnivorous species require lower carbohydrates than herbivorous species. Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to **Mente (2006)**, protein functions as a source of energy for growth and tissue repair. Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (**Radhakrishnan et al., 2020**).

The carbohydrate content of the feed in this study was 30.77 to 35.29%, higher than the crustacean requirement level, which is only about 20 to 30% (**Wang and Chen, 2016**), 18 to 20% in mud crab (**Zhang et al., 2022**); *Litopenaeus vannamei* (**Wang et al., 2014**); *Scylla*

serrata (Truong et al., 2008); and *Penaeus monodon* (Fabricius) (Alava and Pascual, 1987). The high carbohydrate content in this study aims to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the test parameters. Crab growth is a function of molting (Tamsil and Hasnidar, 2018). The more often a crab molts, the faster its growth. The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to metabolize carbohydrates properly so that molting and growth are less than optimal. The same thing was also stated by (Zhang et al., 2022), that carbohydrate levels of 24.0 to 30.0% in feed negatively affect growth, prolong the molting cycle; and affect crab health (Li et al., 2013). The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. According to Stevens and Jewett (2014), crabs begin physiological preparations for molting about 2 months before releasing their shells. Meanwhile, mangrove crabs grow rapidly through molting occurring every 5-30 days depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth. However, feed with quantity and quality below the optimum will negatively affect growth (Nguyen et al., 2014).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely premolting, molting (ecdysis), postmolting, intermolting, the longest stage is intermolting. If the energy from food is sufficient for growth, the intermolting phase will be faster, and vice versa. Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. When the old carapace is detached (ecdysis), the new carapace will replace it and have a larger size. Therefore, the increase in crab size only occurs during molting. According to Stevens and Jewett (2014), the increase in size of crab every time it molts is around 20% to 25%. Therefore, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions, but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (Tocher, 2003); increase metabolism and respiration in mitochondria (Yuan et al., 2019). Feed lipid levels in this study of 6.31 to 9.94% are levels that can support good crab growth, which is 2 to 10% (Sheen and D'Abramo, 1991; Zhao et al., 2014) or 2 to 12% of the dry weight of food (Catacutan, 2002). The lipid requirements of organisms vary widely depending on species and environmental conditions (Holme et al., 2007). According to Bowser and Rosemark (1981), if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*; inhibit the development of *Marsupenaeus japonicus* larvae (Teshima et al., 1986). Furthermore, high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (Sun et al., 2020). Lipid metabolism in crabs still needs to be studied more deeply because there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells and daily metabolic processes. The results showed that the

treatment had no effect on protein retention. Factors that affect protein retention include size, water temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (**Suhenda *et al.*, 2004; Jobling, 2012**). Because of the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components including sex hormones, and molting activity (**Sheen and Wu, 2003**); growth and survival (**Aaqillah-amr *et al.*, 2021**); reproduction and egg quality (**Islam and Yahya, 2016**). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.10; lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5 (**Hastuti *et al.*, 2016**); 2.57 to 3.60 (**Hudita *et al.*, 2020**). The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals. The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth. The feed given must have a compact texture and not easily crushed. The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, this result is higher than the research reported by **Hastuti *et al.*, (2016)** which is 0 to 83%; 63 to 80% (**Yulianto *et al.*, 2019**). This difference is thought to be due to several factors including treatment, internal and external factors of the test animals. The high survival rate of the results of this study is thought to be because the feed provided produces enough energy to sustain the life of the crab. Environmental conditions, especially salinity, are at a less than optimal level, but mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5), showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. This is thought to be because crabs are less able to metabolize carbohydrates compared to fats. Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. This is thought to be because the energy from carbohydrates and lipids is used to adapt to the non-optimal physical-chemical environment of water, especially salinity (Table 5). According to **Tseng and Hwang (2008)** and **Wang *et al.*, (2012)**, carbohydrates can fulfill the high energy needs of aquatic animals during stressful conditions. Furthermore

Wang *et al.*, (2014), optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vanname shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth, and feed efficiency of mud crabs. However, there is a trend that the treatment with a carbohydrate and lipid ratio of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, 35 and 6%. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

REFERENCES

- Aaqillah-amr M.A.; Hidir A.; Azra M.N. and Noordiyana M.N** (2021). Interactions between food, feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava, V.R. and Pascual F.P.** (1987). Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita, P.** (2020). Performance of oven shelf type drying on BSF larva. Sriwijaya University.
- Apine, E.; Ramappa, P. and Bhatta, R.** (2023). Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit, I.; Liland, N.S.; and Gjesdal, P.** (2019). Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December): pp.609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin, M.** (1989). Methodologies for vitamin requirement studies. pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser, P.R.; and Rosemark, R.** (1981). Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.
- Bruni, L.; Belghit, I. and Lock, E.J.** (2020). Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado, M. P.; Álvarez-González, C. A.; Martínez-García, R.; Peña-Marin, E. S.; Rodríguez-Estrada, U.; Macias, E. B.; Civera-Cerecedo, R. and Civera-Cerecedo, R.** (2022). *Pterygoplichthys spp. (Siluriformes: Loricariidae) meal is suitable for the culture of Nile tilapia Oreochromis niloticus (Cichlidae) juveniles. June 2021.* <https://doi.org/10.1590/0103-8478cr20200801>
- Catacutan, M.R.** (2002). Growth and body composition of juvenile mud crab, *Scylla serrata*,

- fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8
- Cerbule, K.; Grimsmo, L.; Herrmann B. and Grimaldo, E.** (2023). Increasing sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820
- Chepkirui-Boit, V.; Ngugi, C.; Bowman, J.; Oyoo-Okoth, E.; Rasowo, J.; Bundi, J. and Cherop, L.** (2011). *Growth performance, survival, feed utilization and nutrient utilization of African catfish (Clarias gariepinus) larvae co-fed Artemia and a micro-diet containing freshwater atyid shrimp (Caridina nilotica) during weaning.* <https://doi.org/10.1111/j.1365-2095.2009.00737.x>
- Day, L.; Cakebread, J.A.; Loveday, S.M.** (2022). Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Fasakin, E.A.; Balogun, A.M. and Ajayi, O.O.** (2003). Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole, F.J.; Ayodeji, A.; Tihamiyu, L.O.; Ajala, K.; Obadara, S. O. and Ganiyu, I. O..** (2020). Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518** (October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi, M. and Yone, Y.** (1982). Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi* (Japanese Edition) **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen, S.H.M.; Crombag, J.J.R.; Senden, J.M.G.; Waterval, W.A.H.; Bierau, J.; Verdijk, L.B. and Van, Loon, L.J.C.** (2018). Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids*. 2018 Dec;50(12):1685-1695. doi: 10.1007/s00726-018-2640-5. Epub 2018 Aug 30. PMID: 30167963; PMCID: PMC6245118.
- Hasnidar, H.; Tamsil, A.; Saenong, M.; Akram, A.M. and Ardinasyah, M.** (2022). Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti, Y.P.; Nadeak, H.; Affandi, R. and Faturrohman, K.** (2016). Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich, K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati, V. E.; Susilo, A.; Pinandoyo.; Hutabarat, J.; Sugianto, D. N.; Wirasatriya, A. and Radjasa, O. K.** (2019). Optimization Of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) For Growth and Feed Utilization Efficiency Of Juvenile *Litpenaeus Vannamei*. *Asian Jr. of Microbiol. Biotech. Env*, **21**(2), 284–297.

- Herwati, V. E.; Pinandoyo.; Windarto, S.; Hariyadi, P.; Hutabarat, J.; Darmanto, Y.; Nurmanita.; Rismaningsih.; Prayitno, S. B. and Radjasa, O. K.** (2020). Maggot Meal (*Hermetia illucens*) Substitution on Fish Meal to Growth Performance, and Nutrient Content of Milkfish (*Chanos chanos*). *Hayati Journal of Biosciences*, 27(2), 154–165. <https://doi.org/10.4308/hjb.27.2.154>
- Holme, M.H.; Southgate, P.C. and Zeng, C.** (2007). Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme, M.H.; Zeng, C. and Southgate, P.C.** (2009). A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu, Y.; Huang, Y.; Tang T.; Zhong, L.; Chu, W.; Dai, Z.; Chen, K. and Hu, Yi.** (2020). Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345
- Hudita, K.; Agustoni and Lokapirnasari, W.P.** (2020). Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) dddition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam, M.L. and Yahya, K.** (2016). Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam, T.; Saha, D.; Bhowmik, S.; Nordin, N.; Islam, S.; Nur, As-Sd.U. and Begum, M.** (2022). Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling, M.** (2012). National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir, K.A.; Verdegem, M.C.J.; Verreth, J.A.J.; Philps, M. and Schrama, J.W.** (2020). Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan, C.P.** (1999). Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. ACIAR Proceedings.
- Li, X.F.; Wang, Y.; Liu W.B.; Zhen, J.G. and Zhu, J.** (2013). Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell, R.T.; Cho, C.Y.; and Cowey, C.B.** (1993). Nutrient Requirements of Fish. DOI: 10.17226/2115

- Manam, V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente, E.** (2006). Protein nutrition in crustaceans. CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources **1** (043): 1–7. DOI: 10.1079/PAVSNNR20061043
- Mukhlis.; Lausu and Majid, M.** (2020). Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3.
- Nguyen, N.T.B.; Chim, L.; Lemaire, P. and Wantiez, L.** (2014). Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI: 10.1016/j.aquaculture.2014.09.014
- Panase, P.; Uppapong, S.; Tuncharoen, S.; Tanitson, J.; Sootornprasit, K. and Intawicha, P.** (2018). Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekong giant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng, K.; Mo, W.; Xiao, H.; Wang, G and Huang, Y.** (2021). Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan, G.; Yashwanth, B.S.; Swamy, S.; Mannur, V.S.; Pinto, N.; Prathik, M.R. and Kulal, P.A.** (2020). Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.
- Sankian, Z.; Khosravi, S.; Kim, Y.O. and Lee, S.M.** (2018). Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen, S.S. and D'Abramo. L.R.** (1991). Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen, S.S. and Wu S.W.** (2003). Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forskål, 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva.; Raman, S.; Stringer, L.C.; Bruce, N.C. and Chong, C.S.** (2022). Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens, B.G. and Jewett, S.C.** (2014). Growth, Molting, and Feeding of King Crabs. DOI:

10.1201/b16664-12

- Suhenda, N. and Samsudin, R.** (2008). Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.
- Suhenda, N.; Tahapari, E.; Slembrouck, J. and Moreau, Y.** (2004). Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun, P.; Jin, M.; Jiao, L.; Monroig, O.; Navvarro, J. C.; Tocher, D. R.; Bentacor, M.B.; Wang, X.; Yuan, Y. and Zhou, Q.** (2020). Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil, A. and Hasnidar.** (2018). The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima, S.I.; Kanazawa, A. and Yamashita, M.** (1986). Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher, D.R.** (2003). Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong, P.H.; Anderson, A.J.; Mather, P.B.; Paterson, B.D. and Richardson, A.** (2008). Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng, Y.C. and Hwang, P.P.** (2008). Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148** (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan, U. and Paulraj, R.** (2010). Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang, X.; Li, E. and Chen L.** (2016). A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang, X.; Li, E.; Qin, J.G.; Wang, S.; Chen, X.; Cai, Y.; Chen, K.; Hou, Y.; Yu, N.; Zhang, M.; Du, Z. and Chen, L.** (2014). Growth, body composition, and ammonia tolerance of juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220
- Wang, Y.; Li, E.; Yu, N.; Wang, X.; Cai, C.; Tang, B.; Cheng, L. and Wormhoudt, A.V.** (2012). Characterization and expression of glutamate dehydrogenase in response to

acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. PLoS ONE **7** (5): 1–10. DOI: 10.1371/journal.pone.0037316

- Yildirim-Aksoy, M.; Eljack, R.; Schrimsher, C. and Beck, B.H.** (2020). Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. *Aquaculture Reports* **17** (May): 100373. DOI: 10.1016/j.aqrep.2020.100373
- Yuan, Y.; Sun, P.; Jin, M.; Wang, X. and Zhou, Q.** (2019). Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Frontiers in Physiology* **10** (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto, B.; Sunaryo.; Taufiq, N.S.P.; Djunaedi, A.; Subagiyo.; Santosa, A. and Soegianto, A.** (2019). Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. *Ecology Environment and Conservation Journal* **25** (july suppl.): 92–101.
- Zainuddin.; Haryati. and Aslamyiah, S.** (2014). Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. *Journal of Aquaculture Research & Development* **05** (06). DOI: 10.4172/2155-9546.1000274
- Zhang, X.; Jin, M.; Luo, J.; Guo, C.; Zhu, T.; Hu, X.; Yuan, Y. and Zhou, Q.** (2022). Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Aquaculture Nutrition* **2022**: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao, J.; Wen, X.; Li, S.; Zhu, D. and Li, Y.** (2014). Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). *Aquaculture Research* **47** (1–13): 887–899. DOI: 10.1111/are.12547



Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Re: 3892 -P

Magdy Khalil <mtkhalil52@hotmail.com>

2 Maret 2024 pukul 14.59

Kepada: Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Dear Authors,
Your article is in the final process; English language reviewing.
Regards



Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

Issue 5: 3892 -P

1 pesan

Magdy Khalil <mtkhalil52@hotmail.com>
Kepada: Hasnidar Yasin <hasnidar.yasin@umi.ac.id>

14 Maret 2024 pukul 17.14

Dear Authors,
Congratulations....Your article has been published in the EJABF Journal.

Regards

 **Issue 5-3892 P.pdf**
559K

Egyptian Journal of Aquatic Biology & Fisheries
 Zoology Department, Faculty of Science,
 Ain Shams University, Cairo, Egypt.
 ISSN 1110 – 6131
 Vol. 28 (2):67 – 81 (2024)
www.ejabf.journals.ekb.eg



Impact of Iso-Protein Feed with Different Carbohydrate and Lipid Ratios on Growth and Feed Efficiency of the Mud Crab, *Scylla serrata*

Hasnidar^{1*}, Andi Tamsil¹, Nursyahran²

¹Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90231

²Institute of Maritime Technology and Business Balikpapan, Makassar 90245

*Corresponding Author: hasnidar.yasin@umi.ac.id

ARTICLE INFO

Article History:

Received: Dec. 22, 2023

Accepted: Feb. 25, 2024

Online: March 11, 2024

Keywords:

Mud crab,
 Protein
 carbohydrate,
 Lipid ratio

ABSTRACT

The use of artificial feed in mud crab aquaculture continues to be developed considering that feed has an important role in growth, survival, and production costs. The largest production cost comes from feed, especially feed protein. The utilization of carbohydrates and lipids as feed protein savers in mud crab aquaculture has not been widely studied. This study aimed to analyze iso-protein feed with different carbohydrate and lipid ratios on molting, growth, and feed efficiency of mud crab, *Scylla serrata*. Mud crabs with a weight of 70 ± 4.7 g/head and a carapace length of 5 ± 0.5 cm were reared using crab boxes and placed in ponds for 60 days. *Pterygoplichthys pardalis* meal and maggot meal were used as feed protein sources. The treatments in this study were 35% iso-protein feeds with different carbohydrate and lipid ratios: treatments A (31 and 10%), B (33 and 9%), C (34 and 7%), and D (35 and 6%). This study used a completely randomized design consisting of four treatments and three replicates. The parameters observed were molting percentage (MP); growth (absolute weight growth -WGII and specific growth rate -SGRII); feed efficiency (protein retention -PRII & lipid retention -LRII, feed conversion ratio -FCRII, and survival -SRII) in addition to the physico-chemical parameters of the research media. Data were analyzed using a completely randomized design, while water physico-chemistry was analyzed descriptively. The results showed that 35% iso-protein feed with different ratios of carbohydrates and lipids had no significant effect on molting, growth, and feed efficiency. However, there is a trend that the quality of crab meat with the carbohydrate and lipid ratio treatment of 31 and 10% is better than the other treatments. Similar research is still needed with more comprehensive studies and conducted in controlled media so that water quality does not experience extreme fluctuations.

INTRODUCTION

The mud crab, *Scylla* spp. is a fishery commodity from the crustacean group other than shrimp that has high nutritional value (Islam *et al.*, 2022) and specialty flavors, as well as high prices. Producers' demand for crabs is increasing for food needs (Cerbule *et al.*, 2023). The demand for crabs is obtained from catches and aquaculture activities (Keenan, 1999). Crab culture is a profitable activity if conducted with low production costs (Apine *et al.*, 2023). Feed contributes more than half of the production costs (Lovell *et al.*, 1993; Holme *et al.*, 2009; Hasnidar *et al.*, 2022). Feeding with an appropriate nutrient content can result in good

growth and survival (**Chepkirui-Boit *et al.*, 2011**). The nutritional content of the feed is always related to the materials used (**Manam, 2023**).

Efforts to obtain cheap and quality feed ingredients are ongoing. Some of the feed ingredients currently developed as a source of feed protein are Amazon sailfin catfish, *P. pardalis* meal (**Castelnau, 1855; Cano-Salgado *et al.*, 2022; Hasnidar *et al.*, 2022**) and maggot meal (**Herawati *et al.*, 2019; Herawati *et al.*, 2020**). Both ingredients are expected to replace the commercial fishmeal, which is increasingly expensive. *P. pardalis* meal has a protein content of 38.6 to 54.48% and fat content of 6.23 to 15.64% (**Panase *et al.*, 2018; Hasnidar *et al.*, 2022**). Its use as a source of feed protein has been reported by several researchers such as in juvenile Mekong giant catfish, *Pangasianodon gigas* (**Panase *et al.*, 2018**), catfish, *Clarias gariepinus* (**Mukhlis *et al.*, 2020**) in addition to vaname shrimp, *Litopenaeus vannamei* (**Hasnidar *et al.*, 2022**). Apart from *P. pardalis* meal, maggot, *Chrysomya megacephala* meal can also be used as a protein source (**Siva Raman *et al.*, 2022**). Maggot is a fly larva *Hermetia illucens* or black soldier that has metamorphosed, with a protein content of 43.30 to 46.70% and a lipid of 7.00 to 14.35% (**Fasakin *et al.*, 2003**). Substitution of *P. pardalis* with maggot meal has a significant effect on the growth and feed efficiency of *Salmo salar* (**Belghit *et al.*, 2019**), Atlantic salmon (**Bruni *et al.*, 2020**), *Clarias gariepinus* (**Fawole *et al.*, 2020**), milkfish, *Chanos chanos* (**Herawati *et al.*, 2020**), *Monopterus albus* (**Hu *et al.*, 2020**), *Oreochromis niloticus* (**Yildirim-Aksoy *et al.*, 2020**) and *Micropterus salmoides* (**Peng *et al.*, 2021**).

Mud crabs are carnivorous organisms that tend to require high-protein feed. The protein requirement of mud crabs is between 32 to 40% (**Catacutan, 2002**), and 46.90 to 47.03% (**Unnikrishnan & Paulraj, 2010**). Combining different protein sources, i.e. *P. pardalis* and maggot meal, is expected to complement each other, especially the amino acid content that is lacking. The amino acid profile present in a particular protein varies depending on its source (**Gorissen *et al.*, 2018; Day *et al.*, 2022**) in addition to using cheap and quality protein sources for the efficient use of feed, it can also be done by reducing protein content and increasing the carbohydrate and lipid content of feed (**Kabir *et al.*, 2020**).

Protein is the most expensive feed component compared to carbohydrates and lipids. According to **Suhenda and Samsudin (2008)**, lipids and carbohydrates can be a protein-sparing effect, which is a substitute for protein as an energy source, hence the use of energy from protein can be optimized for growth. The reduction of feed protein content also has an impact on reducing the discharge of nitrogen waste into the environment. According to **Zainuddin *et al.* (2014)**, nitrogen waste discharge is higher at 30% feed carbohydrate content compared to 49% content. Furthermore, increasing feed carbohydrates can increase body glycogen levels which can be used for other metabolic activities at any time. Conversely, high feed fat content will cause fat storage in the body, decreased food consumption and growth, as well as liver degeneration. According to **Catacutan (2002)**, mud crabs grow well when given feed with a lipid content of 6 or 12%.

The utilization of carbohydrate and fat as a sparing effect of feed protein use in crabs is not yet available. Information is still limited to the level of protein and lipid of jambal fish with a ratio of carbohydrate and lipid of 32 and 8% (**Suhenda & Samsudin, 2008**), 55 and 7% in golden mandarin fish, *Siniperca scherzei* (**Sankian *et al.*, 2018**). Furthermore, in white

shrimp, *Litopenaeus vannamei* the use of carbohydrates with a content of 38% resulted in better growth.

MATERIALS AND METHODS

The research was conducted in a community aquaculture area in Minasa Upa Village, Bontoa District, Maros Regency, South Sulawesi, Indonesia. The series of research consisted of preparation, i.e. preparing mud crabs, *Scylla serrata* (Forsskal, 1775) research media, research containers (crab boxes), preparing feed ingredients and making feed (January to February, 2023), conducting research (March to April, 2023) & testing feed chemistry and crabs (May, 2023). The stages of the research are as follows:

Preparation of *P. pardalis* and maggot meal

P. pardalis specimens were collected from Lake Tempe, Wajo Regency, South Sulawesi, Indonesia. The fish were cleaned and gutted & then dried for five days. To reduce moisture during drying, the fish were then baked at 70°C for 24 hours (Panase *et al.*, 2018). The dried *P. pardalis* samples were then ground using a blender, and sieved using a 425µm mesh size sieve. Subsequently, the *P. pardalis* meal was ready for use. Maggots were obtained from a maggot farming in Makassar. Maggot was cleaned, and dried in the sun until dry. After drying, the maggot samples were put in an oven at 60°C for 12 hours (Alpionita, 2020), then mashed using a blender and sieved using a 425µm-sized sieve. Furthermore, maggot meal was ready for use.

Preparation of feed

Each feed ingredient used was weighed as needed. The ingredients and the amount needed for each treatment are presented in Table (1). The weighed feed ingredients were mixed evenly by stirring manually until the ingredients were homogeneous. Furthermore, small amounts of ingredients, such as fish oil, vitamin premix, and carboxy metil cellulose (CMC) were added and stirred again. After all the ingredients were evenly mixed, warm water was added to the mixture of feed ingredients until it formed a dough. The feed dough was stirred until it did not stick to the hands, then the dough was molded into pellets using a feed printer. The pelleted feed was then spread evenly on a tray and dried in the sun, then the feed was ready to be applied. The proximate test results of the test feed are presented in Table (2).

Table 1. Feed ingredients and quantities required for each treatment

Feed ingredient	Feed formulation (ratio of carbohydrate and lipid content of feed)			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
<i>P. pardalis</i> meal	30	30	30	30
Maggot meal	30	30	30	30
Soybean meal	24	24	24	24
Bran meal	5	5	5	5
Corn meal	0	2.5	5	7.5
Fish oil	7.5	5	2.5	0
Vitamin premix ¹	3	3	3	3
CMC ²	0.5	0.5	0.5	0.5
Total	100	100	100	100

1) Vitamin premix (in one kg of diet): vit. A 90.000IU, vit. D3 30.000IU, vit. K3 36mg, vit. E 225mg, vit. B1 90mg, vit. B2 135mg, vit. B6 90mg, vit. B12 90mg, vit. C 240mg, calcium D-pentathenate 120mg, folic acid 45mg, biotin 300mg, inositol 375mg, nicotinamide 600mg, and cholin chloride 450mg.

2) Carboxymethylcellulose.

Table 2. Proximate results of the experimental diets in mud crabs, *S. serrata* (Forsskal, 1775)

	Feed formulation (ratio of carbohydrate and lipid)			
	A(31 and 10)%	B(33 and 9)%	C(34 and 7)%	D(35 and 6)%
Feed proximate test result				
Protein (%)	34.49	34.48	34.96	34.97
Carbohydrate (%)	30.77	33.07	33.55	35.29
Lipid (%)	9.94	8.87	7.30	6.31
Ash (%)	17.21	17.75	17.44	18.25
Water (%)	6.27	6.83	7.15	8.70

Description: Analysis results. Lab. polytechnic tester, Pangkep. 2023. Except moisture content, all fractions are expressed in dry matter. *Carbohydrate: extractable material without nitrogen (BETN) + crude fiber.

Mud crabs, *S. serrata* with a weight of 70 ± 4.7 g/ crab and a carapace length of 5 ± 0.5 cm, were obtained from fishermen collecting in the location of the study site. The culture container used was a crab box measuring length $\times$ width $\times$ height that was $20 \times 15 \times 15$ cm as many as 60 pieces, and each crab box was filled with one crab. At the bottom and sides of the crab box, plastic nets with a mesh size of one inch were installed to prevent feed and feed residue from falling through the holes in the crab box.

Furthermore, crab boxes were placed in a row on a maintenance raft made of bamboo, with floats from the used plastic bottles. Water changes in the pond were made once a day. The feed had an iso-protein composition with varying ratios of carbohydrate and lipid, as detailed in Table (1) for each treatment. The crabs were fed 5% of the biomass weight per day, with a frequency of feeding of 2 per day at 7 am and 6 pm during 60 days of rearing.

Observations of the number of molted and dead crabs were made daily, and growth sampling was conducted every 15 days. Dead crabs were recorded and weighed. Crab weight gain was measured using digital scales (O-haus) with an accuracy of 0.1g, and the carapace width was measured using a sliding mister with an accuracy of 0.1mm. Measurement of

physical and chemical properties of water, such as temperature, pH, salinity, dissolved oxygen, ammonia, and nitrite were measured throughout the experiment. Temperature, pH, and salinity were measured daily, while measurements for dissolved oxygen, ammonia, and nitrite were measured at the beginning, middle and end of the experiment.

Observations of crab body chemical composition were made on crabs before treatment (initial) and crabs at the end of the experiment. A total of five crabs were taken randomly for body composition analysis. Proximate analysis of feed and crabs was conducted based on the outlines of the AOAC (**Helrich, 1990**). Dry matter content of samples was obtained by drying them in an oven at 105°C for 16 hours; crude protein was analyzed by micro-Kjeldahl, and fat was determined gravimetrically by chloroform methanol extraction. Crude fiber was analyzed by heating with alternating acid and alkaline washing, and ash by burning in a furnace at 550°C for 24 hours.

Parameters measured were molting (molting percentage –MP%), growth (weight gain –WG), and specific growth rate –SGR), and feed efficiency (protein retention –PR, lipid retention –LR, feed conversion ratio –FCR, and survival rate –SR). The formulation used the **Boonyaratpalin (1989)** equation as follows:

$$MP = \frac{\text{number of molting crabs}}{\text{initial number of crabs}} \times 100$$

$$WG = (\text{final weight of crabs} - \text{initial weight of crabs})$$

$$SGR = \frac{\text{Ln final weight} - \text{Ln initial weight}}{\text{cultivation time}} \times 100$$

$$PR = \frac{\text{increase in body protein weight}}{\text{feed protein weight}} \times 100$$

$$LR = \frac{\text{increase in body lipid weight}}{\text{feed lipid weight}} \times 100$$

$$FCR = \frac{\text{dry weight of feed given}}{(\text{final weight} + \text{dead crab of weight}) - \text{initial weight}}$$

$$SR = \frac{\text{Final number of crabs}}{\text{initial number of crabs}} \times 100$$

Statistical analysis

The experimental design used was a completely randomized design, with four treatments and three replications. The treatments consisted of four feed compositions (Table 1). To determine the effect of treatment on the test parameters, an analysis of variance (ANOVA) was performed, followed by Duncan multiple range test (DMRT) at a significance level of 95% ($P < 0.05$). All data were presented as mean and standard deviation. Statistical analysis was performed using SPSS software version 17 for Windows (SPSS Inc., Chicago, USA). The physical-chemical properties of water were analyzed descriptively.

RESULTS

Mud crabs reared for 60 days with feed composition treatment having different carbohydrate and lipid ratios showed no significant effect on MP, WG, SGR, PR, LR, FCR, and SR (Table 3).

Table 3. Molting, growth, and feed utilization response of mud crab, *S. serrata* (Forsskal, 1775) with different carbohydrate and lipid ratios

Parameter	Feed with different carbohydrate and lipid ratios			
	A (31 and 10)%	B (33 and 9)%	C (34 and 7)%	D (35 and 6)%
MP (%)	33.33±11.55 ^a	33.33±11.55 ^a	26.67±11.55 ^a	26.67±11.55 ^a
WG (g)	158.18±2.94 ^a	158.03±3.63 ^a	155.25±2.65 ^a	153.54±4.05 ^a
SGR (%)	1.79±0.14 ^a	1.78±0.05 ^a	1.72±0.41 ^a	1.71±0.04 ^a
PR (%)	17.98±0.49 ^a	16.74±0.89 ^a	16.43±0.89 ^a	16.31±0.38 ^a
LR (%)	9.19±0.10 ^a	8.97±0.06 ^a	8.55±0.71 ^a	8.65±0.49 ^a
FCR	1.94±0.16 ^a	2.03±0.14 ^a	2.09±0.12 ^a	2.10±1.09 ^a
SR (%)	86.67±11.54 ^a	80.00±0.00 ^a	86.67±11.55 ^a	73.33±11.55 ^a

- MP (molting percentage), WG (weight gain), SGR (specific growth rate), PR (protein retention), LR (lipid retention), FCR (feed conversion ratio), SR (survival rate).

-The results of the analysis of variance showed that the treatments did not significantly affect ($P > 0.05$) the test parameters.

- Mean values followed by the same letter in the same row indicate not significantly different ($P > 0.01$) based on Duncan multiple range test (DMRT)

Based on the proximate test results of crab meat before treatment (initial) and after treatment (final), the protein content of the crab increased, but the lipid and carbohydrate contents decreased (Table 4).

Table 4. Chemical/ proximate test results of mud crab meat, *Scylla serrata* (Forsskal, 1775) during initial and final research

		Proximate composition		
		Protein	Carbohydrate	Lipid
Initial meat		45.98	24.06	2.88
Final meat	A (31 and 10)%	54.20	11.02	2.62
	B (33 and 9)%	52.27	21.43	2.50
	C (34 and 7)%	49.21	23.29	2.47
	D (35 and 6)%	49.17	20.80	2.36

The physicochemical properties of the rearing media during the study and the optimal value for mangrove crab cultivation are shown in Table (5).

Table 5. Physico- chemical variable, measurement result, optimal level, and measuring tools

Physico- chemical variable	Measurement result	Optimal level	Measuring
Temperature	10 to 30 ⁰ C	29 ⁰ C (Hastuti <i>et al.</i> , 2019); 20 to 30 ⁰ C (Alberts-Hubatsch <i>et al.</i> , 2016)	Thermometers
pH	6.0 to 8.0	7.75 to 8.50 (Hastuti <i>et al.</i> , 2016)	pH meters
Salinity	5 to 15 ppt	25 ppt (Hastuti <i>et al.</i> , 2015); 20 ppt ((Suyono, 2021)	Refractometers
Dissolved oxygen (DO)	3 to 5 mg/l	±5.51 mg/l (Faturrohman <i>et al.</i> , 2017)	DO meter
Ammonia (NH ₃)	0.10 to 0.30 mg/l	< 0.5 (Hastuti <i>et al.</i> , 2019)	Spectrofotometer
Nitrite (NO ₂ ⁻)	0.00 to 0.002 mg/l	<0.05 (Hastuti <i>et al.</i> , 2019)	Spectrofotometer

DISCUSSION

The results of this study showed that iso-protein diets with different carbohydrate and lipid ratios did not significantly affect the molting percentage, absolute growth, growth rate, protein retention, fat retention, feed conversion, and survival of mud crabs (Table 3). The chemical (proximate) test results of the four feed treatments contained 34.49 to 34.97% protein, 30.77 to 35.29% carbohydrate & 6.31 to 9.94% fat. The low range of feed carbohydrates and fat is thought to be the reason why the treatments did not affect the test parameters. The protein content of the feed in this study was lower than the needs of mangrove crabs according to **Catacutan (2002)** (32 to 40%), and **Unnikrishnan and Paulraj (2010)** (46.90 to 47.03%). Therefore, it is thought to cause molting, growth, and feed efficiency not to be optimal. Feed with the right ratio of carbohydrate and fat is expected to supply the energy needed by the crabs and also have a protein-sparing effect. According to **Furuichi and Yone (1982)**, carbohydrate requirements vary among species; carnivorous species require lower carbohydrates than herbivorous species. Crabs are carnivorous species, presumably less able to metabolize carbohydrates to produce energy that can support molting and better crab growth. According to **Mente (2006)**, protein functions as a source of energy for growth and tissue repair. Body protein is always in a dynamic balance, and protein requirements vary greatly according to age and species (**Radhakrishnan et al., 2020**).

The carbohydrate content of the feed in this study was 30.77 to 35.29%, which is higher than the crustacean requirement level, which is only about 20 to 30% in *Penaeus monodon* (Fabricius) (**Alava & Pascual, 1987**), *Scylla serrata* (**Truong et al., 2008**), *Litopenaeus vannamei* (**Wang et al., 2014; Wang & Chen, 2016**), and 18 to 20% in mud crab (**Zhang et al., 2022**). The high carbohydrate content in this study aimed to evaluate the optimal carbohydrate requirement to replace protein as a source of energy. The results showed that the treatment did not significantly affect the test parameters. Crab growth is a function of molting (**Tamsil & Hasnidar, 2018**). The more often a crab molts, the faster it grows. The number of crabs molting in this study was 29.0 to 31.7%. Crabs are less able to

metabolize carbohydrates properly, hence molting and growth are less than optimal. The same thing was also stated by **Zhang et al. (2022)** who determined that carbohydrate levels of 24.0 to 30.0% in feed negatively affected growth, prolonged the molting cycle, and affected crab health (**Li et al., 2013**). The low number of molted crabs is also thought to be due to the relatively short research time of only 10 weeks. According to **Stevens and Jewett (2014)**, crabs begin physiological preparations for molting about 2 months before releasing their shells. Meanwhile, mangrove crabs grow rapidly through molting occurring every 5- 30 days, depending on the phase of development. Only a few studies have discussed the effect of feed quality on molting and crab growth. However, feed with quantity and quality below the optimum will negatively affect growth (**Nguyen et al., 2014**).

Molting in crabs is a complex interaction between internal and external factors. The molting process consists of four stages, namely pre-molting, molting (ecdysis), post-molting, and inter-molting; the longest stage is the inter-molting. If the energy from food is sufficient for growth, the inter-molting phase will be faster, and vice versa. Maximum weight growth will stimulate the crab to begin the molting process. This process begins with the formation of a new carapace under the old carapace. When the old carapace is detached (ecdysis), the new carapace will replace it with a larger size. Therefore, the increase in crab size only occurs during molting. According to **Stevens and Jewett (2014)**, the increase in size of crab every time it molts is around 20 to 25%. Hence, the growth pattern of crabs (especially the order decapoda) is discontinuous. High molting activity is not only influenced by the quantity and quality of food, environmental conditions but also the durability of the food when it is held by the crab's claws to be inserted into its mouth.

Lipids in crustaceans are assimilated in the hepatopancreas, distributed to other tissues via hemolymph. Lipids function to produce energy for molting, limb regeneration and growth (**Tocher, 2003**), as well as increase metabolism and respiration in mitochondria (**Yuan et al., 2019**). Feed lipid levels in this study recorded 6.31 to 9.94%, falling within the levels that can support good crab growth, which is 2 to 10% (**Sheen & D'Abramo, 1991; Zhao et al., 2014**) or 2 to 12% of the dry weight of food (**Catacutan, 2002**). The lipid requirements of organisms vary widely depending on species and environmental conditions (**Holme et al., 2007**). **Bowser and Rosemark (1981)**, in their study, denoted that if lipids (phospholipids) are not fulfilled in the feed, it will cause molting syndrome (molting failure) resulting in death in lobster, *Homarus americanus*, as well as inhibiting the development of *Marsupenaeus japonicus* larvae (**Teshima et al., 1986**). Furthermore, the high feed lipid levels inhibit growth and reduce feed utilization, lipolysis, lipid deposition in the hepatopancreas (**Sun et al., 2020**). Lipid metabolism in crabs still needs to be studied more deeply since there are many factors that affect it.

Protein retention is the amount of protein stored in the body and used to build new body tissues, repair damaged cells, and daily metabolic processes. The results showed that the treatment had no effect on protein retention. Factors that affect protein retention include size, water temperature, feed dosage, quantity and quality of feed, feed energy content and protein quality (**Suhenda et al., 2004; Jobling, 2012**). Due to the low feed protein content compared to the crab's needs, it is suspected that the energy for growth is also reduced.

Lipid retention is the amount of feed lipids that can be utilized and stored in the body during the rearing period. Similar to protein retention, the treatment did not significantly

affect lipid retention. Lipids serve as a source of energy, build cell membranes, precursors of physiological components, including sex hormones, molting activity (**Sheen & Wu, 2003**), reproduction and egg quality (**Islam & Yahya, 2016**), as well as growth and survival (**Aaqillah-amr et al., 2021**). The lipid content of the feed is appropriate for the crab's needs, but due to the low protein content of the feed, energy from lipids replaces protein energy to survive.

Feed conversion rate (FCR) is a description of the efficiency of feed utilization, the smaller the FCR value, the more efficient the feed utilization of an aquaculture organism. The FCR in this study was 1.94 to 2.10, which is lower than the FCR of mangrove crabs fed with raw fish, which were 7.20 to 21.5, according to **Hastuti et al. (2016)**, and 2.57 to 3.60 according to **Hudita et al. (2020)**. The difference in FCR value is thought to be due to differences in feed type, namely artificial feed (dry) and loose fish feed (wet), environmental factors and physiological conditions of the test animals. The use of artificial feed in crabs is still being researched in terms of quality, quantity, size, shape and texture of the feed. Crabs have unique eating habits, their claws function as hands to hold the food into the mouth. The feed given must have a compact texture and not easily crushed. The artificial feed in this study has a texture that is not optimal, easily crushed, which is thought to cause high FCR values.

The survival rate of crabs from this study was 73 to 87%, surpassing the findings of previous research by **Hastuti et al. (2016)** which is 0 to 83%, and exceeding those reported by **Yulianto et al. (2019)**, ranging from 63 to 80%. This difference is thought to be due to several factors, including treatment, internal and external factors of the test animals. The high survival rate of the results of this study is thought to be due to the feed provided produces enough energy to sustain the life of the crab. Environmental conditions, especially salinity, are at a less than the optimal level, but mangrove crabs are still able to survive well. Efforts to maintain the osmotic balance of body fluids and the environment (osmoregulation process) require high energy, which affects the low number of crabs molting and growth.

The results of the initial and post-study crab chemistry tests by feeding according to the treatment (Table 5) showed that the protein content of crab meat tended to increase after the study. Although the feed given did not significantly affect the test parameters, the meat protein content showed an increase. The treatment with a carbohydrate and lipid ratio of 31 and 10% has a higher protein content than the other treatments. This is thought to be ascribed to the fact that crabs are less able to metabolize carbohydrates compared to fats. Furthermore, the carbohydrate and lipid content of crab meat at the beginning and after the study tended to decrease. This may be related to the energy released from carbohydrates and lipids which is used to adapt to the non-optimal physical-chemical environment of water, especially salinity (Table 5). According to **Tseng and Hwang (2008)** and **Wang et al. (2012)**, carbohydrates can fulfill the high energy needs of aquatic animals during stressful conditions. Furthermore, **Wang et al. (2014)** stated that optimal carbohydrate levels can reduce the adverse effects of ammonia stress and low salinity on vannamee shrimp.

CONCLUSION

In conclusion, iso-protein feed with different carbohydrate and lipid ratios did not significantly affect molting, growth, and feed efficiency of mud crabs. However, there is a trend that the treatment with carbohydrate and lipid ratios of 31 and 10% is better than the treatments of 33 and 9%, 34 and 7%, as well as 35 and 6%. Additional research is required to further explore this topic comprehensively, ideally conducted in controlled environments to minimize fluctuations in water quality.

REFERENCES

- Aaqillah-amr M.A.; Hidir A.; Azra M.N. and Noordiyana M.N** (2021). Interactions between food, feeding and diets in crustaceans : A review. DOI: 10.20944/preprints202102.0018.v1
- Alava, V.R. and Pascual F.P.** (1987). Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. *Aquaculture* **61** (3–4): 211–217. DOI: 10.1016/0044-8486(87)90150-5
- Alpionita, P.** (2020). Performance of oven shelf type drying on BSF larva. Sriwijaya University.
- Apine, E.; Ramappa, P. and Bhatta, R.** (2023). Challenges and opportunities in achieving sustainable mud crab aquaculture in tropical coastal regions. *Ocean and Coastal Management* **242** (June): 106711. DOI: 10.1016/j.ocecoaman.2023.106711
- Belghit, I.; Liland, N.S.; and Gjesdal, P.** (2019). Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture* **503** (December): pp.609–619. DOI: 10.1016/j.aquaculture.2018.12.032
- Boonyaratpalin, M.** (1989). Methodologies for vitamin requirement studies. pp. 58–67. *In*: De Silva S.. (ed). *Fish Nutrition Research in Asia*. Asian Fisheries Society, Philippines, and International Development Research Centre of Canada, Manila, Philippines.
- Bowser, P.R.; and Rosemark, R.** (1981). Mortalities of cultured lobsters, *Homarus*, Associated with a molt death syndrome. *Aquaculture* **23**: 11–18.
- Bruni, L.; Belghit, I. and Lock, E.J.** (2020). Total replacement of dietary fish meal with black soldier fly (*Hermetia illucens*) larvae does not impair physical, chemical or volatile composition of farmed Atlantic salmon (*Salmo salar* L.). *Journal of the Science of Food and Agriculture* **100** (3): 1038–1047. DOI: 10.1002/jsfa.10108
- Cano-Salgado, M. P.; Álvarez-González, C. A.; Martínez-García, R.; Peña-Marin, E. S.; Rodríguez-Estrada, U.; Macias, E. B.; Civera-Cerecedo, R. and Civera-Cerecedo, R.** (2022). *Pterygoplichthys* spp. (Siluriformes: Loricariidae) meal is suitable for the culture of Nile tilapia *Oreochromis niloticus* (Cichlidae) juveniles. June 2021. <https://doi.org/10.1590/0103-8478cr20200801>
- Catacutan, M.R.** (2002). Growth and body composition of juvenile mud crab, *Scylla serrata*, fed different dietary protein and lipid levels and protein to energy ratios. *Aquaculture* **208** (1–2): 113–123. DOI: 10.1016/S0044-8486(01)00709-8
- Cerbule, K.; Grimsno, L.; Herrmann B. and Grimaldo, E.** (2023). Increasing

- sustainability in food production by using alternative bait in snow crab (*Chionoecetes opilio*) fishery in the barents sea. *Heliyon* **9** (3): e13820. DOI: 10.1016/j.heliyon.2023.e13820
- Chepkirui-Boit, V.; Ngugi, C.; Bowman, J.; Oyoo-Okoth, E.; Rasowo, J.; Bundi, J. and Cherop, L.** (2011). *Growth performance, survival, feed utilization and nutrient utilization of African catfish (Clarias gariepinus) larvae co-fed Artemia and a micro-diet containing freshwater atyid shrimp (Caridina nilotica) during weaning.* <https://doi.org/10.1111/j.1365-2095.2009.00737.x>
- Day, L.; Cakebread, J.A.; Loveday, S.M.** (2022). Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends in Food Science and Technology* **119** (December 2021): 428–442. DOI: 10.1016/j.tifs.2021.12.020
- Fasakin, E.A.; Balogun, A.M. and Ajayi, O.O.** (2003). Evaluation of full-fat and defatted maggot meals in the feeding of clariid catfish *Clarias gariepinus* fingerlings. *Aquaculture Research* **34** (9): 733–738. DOI: 10.1046/j.1365-2109.2003.00876.x
- Fawole, F.J.; Ayodeji, A.; Tihamiyu, L.O.; Ajala, K.; Obadara, S. O. and Ganiyu, I. O..** (2020). Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* **518** (October 2019): 734849. DOI: 10.1016/j.aquaculture.2019.734849
- Furuichi, M. and Yone, Y.** (1982). Availability of Carbohydrate in Nutrition of Carp and Red Sea Bream. *Nippon Suisan Gakkaishi (Japanese Edition)* **48** (7): 945–948. DOI: 10.2331/suisan.48.945
- Gorissen, S.H.M.; Crombag, J.J.R.; Senden, J.M.G.; Waterval, W.A.H.; Bierau, J.; Verdijk, L.B. and Van, Loon, L.J.C.** (2018). Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids*. 2018 Dec;50(12):1685-1695. doi: 10.1007/s00726-018-2640-5. Epub 2018 Aug 30. PMID: 30167963; PMCID: PMC6245118.
- Hasnidar, H.; Tamsil, A.; Saenong, M.; Akram, A.M. and Ardinasyah, M.** (2022). Penggunaan Tepung Ikan Sapu-sapu (*Pterygoplichthys pardalis*) Sebagai sumber Protein Pakan Pada Pentokolan Udang Vaname (*Litopenaeus vannamei*). *Journal of Indonesian Tropical Fisheries* **5** (2): 241–252.
- Hastuti, Y.P.; Nadeak, H.; Affandi, R. and Faturrohman, K.** (2016). Penentuan pH optimum untuk pertumbuhan kepiting bakau *Scylla serrata* dalam wadah terkontrol. *Jurnal Akuakultur Indonesia* **15** (2): 171. DOI: 10.19027/jai.15.2.171-179
- Helrich, K.** 1990. AOAC: Official Methods of Analysis. DOI: 10.7312/seir17116-004
- Herawati, V. E.; Susilo, A.; Pinandoyo.; Hutabarat, J.; Sugianto, D. N.; Wirasatriya, A. and Radjasa, O. K.** (2019). Optimization Of Fish Meal Substitution With Maggot Meal (*Hermetia Illucens*) For Growth and Feed Utilization Efficiency Of Juvenile *Litopenaeus Vannamei*. *Asian Jr. of Microbiol. Biotech. Env*, 21(2), 284–297.
- Herwati, V. E.; Pinandoyo.; Windarto, S.; Hariyadi, P.; Hutabarat, J.; Darmanto, Y.; Nurmanita.; Rismarningsih.; Prayitno, S. B. and Radjasa, O. K.** (2020). Maggot Meal (*Hermetia illucens*) Substitution on Fish Meal to Growth Performance, and Nutrient Content of Milkfish (*Chanos chanos*). *Hayati Journal of Biosciences*, 27(2), 154–165. <https://doi.org/10.4308/hjb.27.2.154>

- Holme, M.H.; Southgate, P.C. and Zeng, C.** (2007). Survival, development and growth response of mud crab, *Scylla serrata*, megalopae fed semi-purified diets containing various fish oil:corn oil ratios. *Aquaculture* **269** (1–4): 427–435. DOI: 10.1016/j.aquaculture.2007.05.024
- Holme, M.H.; Zeng, C. and Southgate, P.C.** (2009). A review of recent progress toward development of a formulated microbound diet for mud crab, *Scylla serrata*, larvae and their nutritional requirements. *Aquaculture* **286** (3–4): 164–175. DOI: 10.1016/j.aquaculture.2008.09.021
- Hu, Y.; Huang, Y.; Tang T.; Zhong, L.; Chu, W.; Dai, Z.; Chen, K. and Hu, Yi.** (2020). Effect of partial black soldier fly (*Hermetia illucens* L.) larvae meal replacement of fish meal in practical diets on the growth, digestive enzyme and related gene expression for rice field eel (*Monopterus albus*). *Aquaculture Reports* **17** (February). DOI: 10.1016/j.aqrep.2020.100345
- Hudita, K.; Agustoni and Lokapirnasari, W.P.** (2020). Penambahan crude fish oil (CFO) pada pakan terhadap pertumbuhan dan rasio konversi pakan pada kepiting bakau (*Scylla serrata*) addition of Crude Fish Oil (CFO) on feed toward growth and Feed Conversion Ratio of Mud Crab (*Scylla serrata*). *Journal of Marine and Coastal Science* **9** (1): 30–40.
- Islam, M.L. and Yahya, K.** (2016). Successive reproductive performance and amino acid profiles in the newly hatched larvae of green mud crab (*Scylla paramamosain*) under captive condition. *International Journal of Fisheries and Aquatic Studies* **4** (5): 270–278.
- Islam, T.; Saha, D.; Bhowmik, S.; Nordin, N.; Islam, S.; Nur, As-Sd.U. and Begum, M.** (2022). Nutritional properties of wild and fattening mud crab (*Scylla serrata*) in the south-eastern district of Bangladesh. *Heliyon* **8** (6): e09696. DOI: 10.1016/j.heliyon.2022.e09696
- Jobling, M.** (2012). National Research Council (NRC): Nutrient requirements of fish and shrimp. *Aquaculture International* **20** (3): 601–602. DOI: 10.1007/s10499-011-9480-6
- Kabir, K.A.; Verdegem, M.C.J.; Verreth, J.A.J.; Philips, M. and Schrama, J.W.** (2020). Effect of dietary carbohydrate to lipid ratio on performance of Nile tilapia and enhancement of natural food in pond aquaculture. *Aquaculture Research* **51** (5): 1942–1954. DOI: 10.1111/are.14546
- Keenan, C.P.** (1999). Aquaculture of the mud crab, genus *scylla* — past, present and future. Pp. 9–13. ACIAR Proceedings.
- Li, X.F.; Wang, Y.; Liu W.B.; Zhen, J.G. and Zhu, J.** (2013). Effects of dietary carbohydrate/lipid ratios on growth performance, body composition and glucose metabolism of fingerling blunt snout bream *Megalobrama amblycephala*. *Aquaculture Nutrition* **19** (5): 701–708. DOI: 10.1111/anu.12017
- Lovell, R.T.; Cho, C.Y.; and Cowey, C.B.** (1993). Nutrient Requirements of Fish. DOI: 10.17226/2115
- Manam, V.K.** 2023. Fish Feed Nutrition and its Management in Aquaculture Fish feed nutrition and its management in aquaculture. (April). DOI: 10.22271/fish.2023.v11.i2a.2791
- Mente, E.** (2006). Protein nutrition in crustaceans. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **1** (043): 1–7. DOI:

10.1079/PAVSNNR20061043

- Mukhlis.; Lausu and Majid, M.** (2020). Pengaruh pemberian pakan buatan berupa tepung ikan sapu-sapu (*Hypostomus* sp) terhadap pertumbuhan dan sintasan lele dumbo (*Clarias gariepinus*). Pp. 185–191. Prosiding Seminar Nasional Pertanian Peternakan Terpadu ke-3.
- Nguyen, N.T.B.; Chim, L.; Lemaire, P. and Wantiez, L.** (2014). Feed intake, molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of prote. *Aquaculture* **434** (October): 499–509. DOI: 10.1016/j.aquaculture.2014.09.014
- Panase, P.; Uppapong, S.; Tuncharoen, S.; Tanitson, J.; Sootornprasit, K. and Intawicha, P.** (2018). Partial replacement of commercial fish meal with Amazon sailfin catfish *Pterygoplichthys pardalis* meal in diets for juvenile Mekong giant catfish *Pangasianodon gigas*. *Aquaculture Reports* **12** (April): 25–29. DOI: 10.1016/j.aqrep.2018.08.005
- Peng, K.; Mo, W.; Xiao, H.; Wang, G and Huang, Y.** (2021). Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports* **20**: 100737. DOI: 10.1016/j.aqrep.2021.100737
- Radhakrishnan, G.; Yashwanth, B.S.; Swamy, S.; Mannur, V.S.; Pinto, N.; Prathik, M.R. and Kulal, P.A.** (2020). Dietary protein requirement for maintenance, growth, and reproduction in fish: A review. *Journal of Entomology and Zoology Studies* **8** (4): 208–215.
- Sankian, Z.; Khosravi, S.; Kim, Y.O. and Lee, S.M.** (2018). Effects of dietary inclusion of yellow mealworm (*Tenebrio molitor*) meal on growth performance, feed utilization, body composition, plasma biochemical indices, selected immune parameters and antioxidant enzyme activities of mandarin fish (*Siniperca scherzeri*) juveniles. *Aquaculture* **496**: 79–87. DOI: 10.1016/j.aquaculture.2018.07.012
- Sheen, S.S. and D’Abramo. L.R.** (1991). Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93** (2): 121–134. DOI: 10.1016/0044-8486(91)90211-O
- Sheen, S.S. and Wu S.W.** (2003). Essential fatty acid requirements of juvenile mud crab *Scylla serrata* (Forskål, 1775) (Decapoda, Scyllaridae). *Crustaceana* **75** (11): 1387–1401.
- Siva.; Raman, S.; Stringer, L.C.; Bruce, N.C. and Chong, C.S.** (2022). Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production* **373** (March): 133802. DOI: 10.1016/j.jclepro.2022.133802
- Stevens, B.G. and Jewett, S.C.** (2014). Growth, Molting, and Feeding of King Crabs. DOI: 10.1201/b16664-12
- Suhenda, N. and Samsudin, R.** (2008). Pemanfaatan pakan iso protein dengan kadar karbohidrat dan lemak yang berbeda untuk pertumbuhan benih ikan patin jambal (*Pangasius djambal*). *Jurnal Riset Akuakultur* **3** (4): 215–224.

- Suhenda, N.; Tahapari, E.; Slembrouck, J. and Moreau, Y.** (2004). Retensi protein dan pemanfaatan energi pada benih ikan patin jambal (*Pangasius djambal*) yang diberi pakan berprotein tinggi. *Jurnal Penelitian Perikanan Indonesia* **10** (5): 65–69. DOI: 10.15578/jppi.10.5.2004.65-69
- Sun, P.; Jin, M.; Jiao, L.; Monroig, O.; Navvarro, J. C.; Tocher, D. R.; Bentacor, M.B.; Wang, X.; Yuan, Y. and Zhou, Q.** (2020). Effects of dietary lipid level on growth, fatty acid profiles, antioxidant capacity and expression of genes involved in lipid metabolism in juvenile swimming crab, *Portunus trituberculatus*. *British Journal of Nutrition* **123** (2): 149–160. DOI: 10.1017/S0007114519002563
- Tamsil, A. and Hasnidar.** (2018). The effect of molting hormone (20-hydroxyecdysone) on molting of mud crab (*Scylla olivacea* Herbst, 1976). *Ecology, Environment and Conservation* **24** (2): 960–967.
- Teshima, S.I.; Kanazawa, A. and Yamashita, M.** (1986). Dietary value of several proteins and supplemental amino acids for larvae of the prawn *Penaeus japonicus*. *Aquaculture* **51** (3–4): 225–235. DOI: 10.1016/0044-8486(86)90314-5
- Tocher, D.R.** (2003). Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11** (2): 107–184. DOI: 10.1080/713610925
- Truong, P.H.; Anderson, A.J.; Mather, P.B.; Paterson, B.D. and Richardson, A.** (2008). Effect of selected feed meals and starches on diet digestibility in the mud crab, *Scylla serrata*. *Aquaculture Research* **39** (16): 1778–1786. DOI: 10.1111/j.1365-2109.2008.02056.x
- Tseng, Y.C. and Hwang, P.P.** (2008). Some insights into energy metabolism for osmoregulation in fish. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology* **148** (4): 419–429. DOI: 10.1016/j.cbpc.2008.04.009
- Unnikrishnan, U. and Paulraj, R.** (2010). Dietary protein requirement of giant mud crab *Scylla serrata* juveniles fed iso-energetic formulated diets having graded protein levels. *Aquaculture Research* **41** (2): 278–294. DOI: 10.1111/j.1365-2109.2009.02330.x
- Wang, X.; Li, E. and Chen L.** (2016). A Review of Carbohydrate Nutrition and Metabolism in Crustaceans. *North American Journal of Aquaculture* **78** (2): 178–187. DOI: 10.1080/15222055.2016.1141129
- Wang, X.; Li, E.; Qin, J.G.; Wang, S.; Chen, X.; Cai, Y.; Chen, K.; Hou, Y.; Yu, N.; Zhang, M.; Du, Z. and Chen, L.** (2014). Growth, body composition, and ammonia tolerance of juvenile white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate levels at low salinity. *Journal of Shellfish Research* **33** (2): 511–517. DOI: 10.2983/35.033.0220
- Wang, Y.; Li, E.; Yu, N.; Wang, X.; Cai, C.; Tang, B.; Cheng, L. and Wormhoudt, A.V.** (2012). Characterization and expression of glutamate dehydrogenase in response to acute salinity stress in the Chinese mitten crab, *Eriocheir sinensis*. *PLoS ONE* **7** (5): 1–10. DOI: 10.1371/journal.pone.0037316
- Yildirim-Aksoy, M.; Eljack, R.; Schrimsher, C. and Beck, B.H.** (2020). Use of dietary frass from black soldier fly larvae, *Hermetia illucens*, in hybrid tilapia (Nile x Mozambique, *Oreochromis niloticus* x *O. mozambique*) diets improves growth and resistance to bacterial diseases. *Aquaculture Reports* **17** (May): 100373. DOI: 10.1016/j.aqrep.2020.100373

-
- Yuan, Y.; Sun, P.; Jin, M.; Wang, X. and Zhou, Q.** (2019). Regulation of dietary lipid sources on tissue lipid classes and mitochondrial energy metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Frontiers in Physiology* **10** (APR): 1–13. DOI: 10.3389/fphys.2019.00454
- Yulianto, B.; Sunaryo.; Taufiq, N.S.P.; Djunaedi, A.; Subagiyo.; Santosa, A. and Soegianto, A.** (2019). Survival and growth of mud crab (*Scylla serrata* Forsskal, 1775) reared in crab bucket recirculating aquaculture system. *Ecology Environment and Conservation Journal* **25** (july suppl.): 92–101.
- Zainuddin.; Haryati. and Aslamyah, S.** (2014). Effect of dietary carbohydrate levels and feeding frequencies on growth and carbohydrate digestibility by white shrimp *Litopenaeus vannamei* under laboratory conditions. *Journal of Aquaculture Research & Development* **05** (06). DOI: 10.4172/2155-9546.1000274
- Zhang, X.; Jin, M.; Luo, J.; Guo, C.; Zhu, T.; Hu, X.; Yuan, Y. and Zhou, Q.** (2022). Effects of dietary carbohydrate level on the growth and glucose metabolism of juvenile swimming crab, *Portunus trituberculatus*. *Aquaculture Nutrition* **2022**: 1–15. DOI: <https://doi.org/10.1155/2022/7110052>
- Zhao, J.; Wen, X.; Li, S.; Zhu, D. and Li, Y.** (2014). Effects of different dietary lipid sources on tissue fatty acid composition, serum biochemical parameters and fatty acid synthase of juvenile mud crab *Scylla paramamosain* (Estampador 1949). *Aquaculture Research* **47** (1–13): 887–899. DOI: 10.1111/are.12547