

BUKTI KORESPONDENSI

ARTIKEL JURNAL INTERNASIONAL BEREPUTASI

Judul artikel : Heavy metals landfill effect on people and organism health risk:
atomic absorption spectroscopy approach

Jurnal : Oxidation Communication, 2023.

Penulis : Andi Artiningsih, Sri Widodo

No.	Perihal	Tanggal
1.	Bukti konfirmasi submit artikel dan artikel yang disubmit	28 April 2023
2.	Bukti konfirmasi review dan hasil review pertama	27 Mei 2023
3.	Bukti konfirmasi submit revisi pertama, respon kepada reviewer, dan artikel yang diresubmit	9 Juni 2022
4.	Bukti konfirmasi naskah diterima	3 Juli 2023
5.	Bukti konfirmasi artikel di published	5 Agustus 2023

**1. Bukti konfirmasi submit artikel dan
artikel yang disubmit
(28 April 2023)**



Andi Artiningsih <andi.artiningsih@umi.ac.id>

Fri, Apr
28,
1:54 PM

to scibulcom2

Dear Editor(s)

We wish to submit an original research article entitled “**Heavy Metals landfill effect on people and organism Health Risk: The Atomic Absorption Spectroscopy Approach**” for consideration by *Oxidation Communication*. This is an original study and has not been published elsewhere, nor is it currently under consideration for publication elsewhere.

Sincerely,

Heavy Metals landfill effect on people and organism Health Risk: The Atomic Absorption Spectroscopy Approach

Andi Artiningsih¹, Sri Widodo²

¹Faculty of Engineering, Universitas Muslim Indonesia, South Sulawesi Indonesia

²Faculty of Engineering, Hasanuddin University, South Sulawesi Indonesia

Corresponding Author: andi.artiningsih@umi.ac.id

Some urban city faced to soils around landfill. This study investigates the heavy metals contamination around the Tamangapa Antang Final Disposal Site (FDS). The sample were divided into 4 Points consisting of 2 Points that align with the groundwater flow (Point A and Point C), and 2 Points that are vertical to the groundwater flow (Point B and Point D) in the Antang landfill. the Atomic Absorption Spectroscopy (AAS) was used to examine heavy metal around sample observation. The results of the analysis shown that the soils of landfill have spread across the topsoil vertically with the highest concentration, it reflecting the vast amount of trash dumped to the FDS. The second highest heavy metal concentration was which was at its highest at Point A and Point D exceeding the limit, but remained under the limit recommended by the Indonesian Food and Drug Authority at Point B and Point C. Meanwhile, Point A, B, and C heavy metal concentrations were lower. The findings showed that in general, FDS has been contaminated hence need to concern for regional disaster management. Specifically for people which lives around the Antang landfill.

Contributions: The construction of regional centers for recycling and reuse activities, and consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities.

Keywords: Landfill, Soils, Vertical, AAS, Heavy Metals

Introduction

The main issue in developing countries is how to select a suitable area for a final disposal site (FDS). Because of the rapid population growth and urbanization, the demand for disposal products has increased by a factor of two (Mokoele 2023; Njoku et al. 2019; Weldeyohanis et al. 2022). It caused a greater need for landfill space. Hence, an integration approach for waste disposal and waste management plays an important role in reducing the impact of solid waste and heavy meal on the environment and the people who live in and around landfill areas. People who reside nearby landfills are exposed to a number of harmful chemicals that are created during the landfilling process (Sasaki et al. 2019). Litter, dust, excessive rodents, unforeseen landfill fires are some additional contaminants linked to the deposition of trash on landfills. The type and quantity of garbage placed, the age of the landfill, and the climatic conditions of the landfill sites are all factors that affect the by-product or emissions from landfills. Heavy metals, particulate debris, and a

number of gaseous contaminants (including dioxins and polycyclic aromatic hydrocarbons) are regularly formed in the landfill as a result of intricate chemical and microbiological processes (Abdelli et al. 2017).

Urbanization is a key issue in developing nations like Indonesia, which has faced various environmental challenges, including the percentage of solid waste, due to social issues including socioeconomic, economic, and population inequality. Traditional solid waste management methods used in Indonesian towns have not been designed to advance environmental protection, public health, or renewable energy sources (Hoque & Rahman 2020; Mester et al. 2022). These methods of managing solid waste are now widely recognized as the primary contributors to contamination of surface and groundwater as well as a degradation in the cleanliness of cities and towns (Morais et al. 2021). Makassar is one of the places, and it deals with concerns including how to manage heavy metal problems in final disposal sites and poorly managed solid waste operations (FDS). Makassar City has been undergoing a significantly rapid development which unfortunately contributes to the increasing amount of waste produced. Waste is an undeniably serious issue which grows with the population in Makassar City.

The Final Disposal Site (FDS) is a vital necessity to create a convenient environment for the community in big cities. If waste from all areas of Big City is not transferred to the FDS even for one day, the city will be full of waste, and it annoys the convenience of the community. It is an undeniable truth that the current conditions of FDS in some big cities have not met the standards of environmental safety. In fact, some FDSs are shut down because they are located too close to people residence and the resulting air pollution from waste deterioration has aggravated the community (Manimekalai & Vijayalakshmi 2014). Poorly managed FDS is critically harmful to human health and potentially damaging the environment. The human activities have caused serious damage to the environment, and significantly increased biological toxicity of heavy metals in soils. Some heavy metals excessive concentration may harm the organisms around landfill. The heavy metal contamination also damages soils and animals (Abiola et al. 2021; Chibuike & Obiora 2014; Tshelaza et al. 2021).

Soils may be contaminated with heavy metals that are typically without color nor odor, thus difficult to be controlled. Heavy metal contamination may not instantly be damaging to environment; however, Nawaz et al. (2023) reported that when the concentration of heavy metals has exceeded the tolerance level of the environment, they will be activated and trigger ecological damage. Previous researchers have investigated heavy metal concentration in the soils landfill in Slovakia (Andras et al. 2013). Similarly, Esfahani et al. (2021) concluded heavy metal in landfill may removal by modified membrane.

Methods

Research Area

The administrative area of the research encompassed the regional Bangkala Sub-district at Manggala District, Makassar City, South Sulawesi Province.




Figure 1 The Location Map and Sampling Points taken vertically for Point A, Point B, Point C, and Point D at the depth of 0-10 cm, 10-20 cm, and 20-30 cm, around Antang Final Disposal Site in Makassar, Sulawesi Selatan.

Sampling Point

The samples used in this study were the topsoil collected from the area of Antang Final Disposal Site, Makassar City, South Sulawesi. There were four sampling points: two points (Point A and Point C) aligned with the groundwater flow, and two points (Point B and Point D) were vertical to

the groundwater flow. All points were measured vertically at the depth of 0-10, 10-20, and 20-30 cm deep. There were a total of 12 samples subjected to analysis of heavy metal Cu, Pb, and Zn using the *Atomic Absorption Spectroscopy* (AAS).

Results and Discussion

Results

The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point A at the depth of 0-10, 10-20, and 20-30 are presented in Figure 2 below:




Figure 2: Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point A at the depth of 0-10, 10-20, and 20-30.

Point A shows that the concentration of Cu at the depth of 0-10cm is 138 ppm, then increases to 168 ppm and 177 ppm at the depth of 10-20 and 20-30 cm, respectively. The concentration of Pb at 0-10cm deep is around 104 ppm, then increases to 11 ppm at 10-20 cm deep, and increases to 105 ppm at 20-30 cm deep. Meanwhile, Zn concentration at 0-10cm was 128 ppm which decreases to 126 and 118 ppm at the depth of 10-20 and 20-30 cm, respectively. The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point B at the depth of 0-10, 10-20, and 20-30 are presented in Figure 3:




Figure 3 Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point B at the depth of 0-10, 10-20, and 20-30.

Point B shows that the concentration of Cu at the depth of 0-10 cm is 67 ppm, then increases to 72 ppm (10-20 cm) and decreases to 64 ppm (20-30 cm). Pb concentration at 0-10cm deep is around 52 ppm, then decreases to 43 ppm (10-20 cm) deep and increases to 63 ppm (20-30 cm) deep. Meanwhile, Zn concentration at 0-10cm was 61 ppm, then increases to 64 ppm (10-20 cm), and decrease back to 61 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point C at the depth of 0-10, 10-20, and 20-30 are presented in Figure 4:




Figure 4 Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point C at the depth of 0-10, 10-20, and 20-30.

Point C shows that the concentration of Cu at the depth of 0-10 cm is 65 ppm, then increases to 84 ppm (10-20 cm) and decreases to 43 ppm (20-30 cm). Pb concentration at 0-10cm deep is around 61 ppm, then increases to 91 ppm (10-20 cm) deep and down to 31 ppm (20-30 cm) deep. Meanwhile, Zn concentration at 0-10cm was 50 ppm, then increases to 58 ppm (10-20 cm), and

decrease to 4 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point CD at the depth of 0-10, 10-20, and 20-30 are presented in Figure 5.




Figure 5 Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point D at the depth of 0-10, 10-20, and 20-30.

Point D shows that the concentration of Cu at the depth of 0-10 cm is 158 ppm, then increases to 161 ppm (10-20 cm) and decreases to 149 ppm (20-30 cm). Pb concentration at 0-10cm deep is around 73 ppm, then decreases to 61 ppm (10-20 cm) deep and back up again to 73 ppm (20-30 cm) deep. Meanwhile, Zn concentrations at 0-10, 10-20, and 20-30 cm are around 132 ppm, 142 ppm dan 142 ppm, respectively.

Discussion

Heavy metals, such as Cu, Pb, and Zn in Antang Final Disposal Site, Makassar City, South Sulawesi have vertically distributed across the topsoil with a variety of concentrations at each depth which has exceeded the limit of contaminant in soil environment, which is acidic. The recommended limit of heavy metal concentrations by the Government of the Republic of Indonesia and Food and Drug Authority is 100 ppm of Cu, 100 ppm of Pb, and 70 ppm of Zn.

Heavy metal Cu is strongly associated with organic matters, iron and manganese oxidation, silica clay and a slight amount of mineral. Cu is particularly absorbed and slightly mobile in the soils. Organic matter is the dominant factor that control the retention of heavy metal Cu. Most Cu is soluble in the soil surface in a complex form of Cu^{2+} . It means that, if Pb heavy metal in the soils functions as Pb^{2+} oxidizer. If pH increases, the Pb^{2+} ion becomes less soluble oxidizer. A higher pH will form a complex form which allows the organic matters settle as carbonate, hydroxide, and

phosphate. Lead is one of heavy metals that poses significant harms to living organisms because it is carcinogenic. The recent study found a varying levels of Zn and the average pH was indicative of acidity. The oxidation of Zn in the soils shows Zn^{2+} under oxidized acid conditions. Zn^{2+} is one of heavy metals that is the most soluble and mobile in the soil, hydrocarbon oxide, and organic matters. Zn solubility in the soils is highly potential at acidic pH (Faborode 2022).

The highest concentrations of heavy metals in this study which also have surpassed the limit are tolerance. High levels of Cu in Final Disposal Site reflect the high quantity of waste dumped to the FDS (Sulaiman et al. 2016). The enrichment of heavy metal in the Antang landfill has been highly accumulated because of the anthropogenic activities or human activities, Leachate, and soil characteristics (Bityukova et al. 2000) and supported by a very acidic pH, the mobility of acid in heavy metals in the soils is relatively high (Schmitt & Sticker 1991). Heavy metals will be carried by Leachate and deposited in the soils, and as a result, heavy metals like Cu, Pb, and Zn will remain deposited in the soils.

Relatively acid soils have an increasing solubility and mobility of nutrients, thus increasing the concentration of heavy metal in the soils (Weldeyohanis et al. 2022; Chibuike & Obiora, 2014). If the soil solution is too acidic, plants cannot utilize and other nutrients they need to grow. Meanwhile, acidic soils (low pH) is dominated by ion Al, Fe, and Mn which will bind the nutrients vital for the plants, especially P, K S, and Mg. As a result, plants cannot absorb their food well despite the richness of nutrients in the soil. In addition, acidic soils contain high level of micro nutrients, which are very dangerous to people's health (Sharma & Jain 2020) particularly those living around the landfill because it causes cancer and lung damage in addition to IQ drop among children, and potential death to human in general.

CONCLUSION

The current study used a multifaceted approach to examine the key elements affecting environmental appropriateness and choosing the best location for the final disposal site (FDS) in the City of Makassar in Indonesia. The determination of sub-criteria and criteria values as well as the choice of the final location utilize a method that combines GIS and an analytical hierarchy

process with data from remote sensing and laboratory analysis. The investigation of environmental suitability, comparison of the identified heavy metal and solid waste disposal sites, and final site rankings in accordance with the specified criteria are all included in the study. The issue of sustainable waste management has come into sharper focus as a result of several studies and scientific investigations. The increased manufacturing and widespread availability of different consumer items have an impact on the continuous rise in garbage output. The purpose of this work was to propose a multi-criteria analysis technique that is integrated with geographic information systems and to highlight the benefits that this approach provides for decision-making in environmental studies involving heavy metal analyses. By taking into account many facets of the problem and different stakeholders in the decision-making process, multi-criteria approaches offer a multi-dimensional approach to problems.

Therefore, the methodological properties of multi-criteria analysis, especially from the perspective of environmental planning and environmental management, permit its successful application in environmental and geographical sciences. The use of these approaches and the outcomes attained are in line with previously established waste management practices. Municipal waste management and the effects of heavy metals on the environment and community health in Indonesia can be significantly improved by even just coordinated work and integrated activities of experts, researchers, government and municipal officials, as well as citizens and other stakeholders. The expansion of the municipal trash collection system, the construction of regional centers for recycling and reuse activities, and finding practical and affordable alternatives to locate consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities. In addition to keeping people and the environment safe, this can offer financial payback and the chance to partially recoup maintenance costs for waste management systems. As it gathers useful informational bases required for the development of environmental protection strategies, city development strategies, and information for spatial planning in the Indonesia region, this research can be applied by the city's official agencies for environmental management and protection as well as spatial planning.

REFERENCES

- Abdelli, I.S., Asnoune, M., Arab, Z., Abdelmalek, F. & Addou, A. 2017. 'Management of household waste in sanitary landfill of Mostaganem district (Western Algeria)'. *Journal of Material Cycles and Waste Management* 19, 265–281. <https://doi.org/10.1007/s10163-015-0415-6>
- Andras, P., Turisova, L.E., Kharbish, S., Krnac, J. & Cmielova, L. 2013. 'Environmental risk due to heavy metal contamination caused by old copper mining activity at Lubietova Deposit, Slovakia'. *Energy and Environment Research* 3(2), 1–15. <https://doi.org/10.5539/eer.v3n2p182>
- Bityukova L, Shogenova, A. & Birke, M. 2000. 'Urban Geochemistry: A study of element distribution in the soil of Tallin (Estonia)'. *Environmental Geochemistry and Health*. 22; 173-193. <https://doi.org/10.1023/A:1006754326260>
- Chibuikwe, G.U. & Obiora, S.C. 2014. 'Heavy metal polluted soil: Effect on plants and bioremediation methods'. *Hindawi Publishing Corporation; Applied and Environmental soil science*, 1-13. <https://doi.org/10.1155/2014/752708>.
- Esfahani, A.R., Zhai, L. & Sadmani, A.H.M.A. 2021. 'Removing heavy metals from landfill leachate using electrospun polyelectrolyte fiber mat-laminated ultrafiltration membrane'. *Journal of Environmental Chemical Engineering* 9(4), 105355. <https://doi.org/10.1016/j.jece.2021.105355>
- Faborode, H.F.B. 2022. 'Gender assessment of poultry waste utilization among small-scale poultry farmers in Osun state, Nigeria: Exploring the untapped potentials'. *African Journal of Food, Agriculture and Nutrition Development* 22(4), 20259–20279. <https://doi.org/10.18697/ajfand.109.19780>
- Hoque, M.M. & Rahman, M.T.U. 2020. 'Landfill area estimation based on solid waste collection prediction using NN model and final waste disposal options'. *Journal of Cleaner Production* 256, 1–12. <https://doi.org/10.1016/j.jclepro.2020.120387>
- Manimekalai, B. & Vijayalakshmi, P. 2014. 'Analysis of Leachate Contamination Potential of a Municipal Landfill Using Leachate Pollution Index'. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 2(1), 16–39. <https://doi.org/10.9790/2402-0211639>
- Mester, T., Szabó, G., Sajtos, Z., Baranyai, E., Szabó, G. & Balla, D. 2022. 'Environmental hazards of an recultivated liquid waste disposal site on soil and groundwater'. *Water* 14, 1–17. <https://doi.org/10.3390/w14020226>
- Mokoele, N.J., 2023, 'Complexities of contemporary urban planning in local government in the City of Polokwane, Limpopo province', *Jambá: Journal of Disaster Risk Studies* 15(1), a132. <https://doi.org/10.4102/jamba.v15i1.1326>
- Morais, L., Nascimento, V., Simões, S. & Ometto, J. 2021. 'Regional distance routes estimation for municipal solid waste disposal, case study São Paulo State, Brazil'. *Energies* 14, 1–14. <https://doi.org/10.3390/en14133964>
- Nawaz, R., Aslam, M., Nasim, I., Irshad, M.A., Ahmad, S., Latif, M. & Hussain, F. 2023. 'Air Pollution Tolerance Index and Heavy Metals Accumulation of Tree Species for Sustainable

Environmental Management in Megacity of Lahore'. *Air* 1(1), 55-68.
<https://doi.org/10.3390/air1010004>

- Njoku, P.O., Edokpayi, J.N. & Odiyo, J.O., 2019. 'Health and environmental risks of residents living close to a landfill: A case study of Thohoyandou landfill, Limpopo Province, South Africa', *International Journal of Environmental Research and Public Health* 16, 1–27.
<https://doi.org/10.3390/ijerph16122125>
- Parth, V., Murthy, N.N. & Raj Saxena, P. 2011. 'Assessment of heavy metal contamination in soil around hazardous waste disposal sites in Hyderabad city (India): natural and anthropogenic implications'. *E3 Journal of Environmental and Research Management* 2(2), 27-34.
- Sasaki, S., Watanabe, K., Widyaningsih, N. & Araki, T. 2019. 'Collecting and dealing of recyclables in a final disposal site and surrounding slum residence: The case of Bantar Gebang, Indonesia'. *Journal of Material Cycles and Waste Management* 21, 375–393.
<https://doi.org/10.1007/s10163-018-0798-2>
- Schmitt, H.S. & Sticker, H. 1991. 'Heavy metal compounds in the soil'. Merian, E. (Ed.), *Metals and their compounds in the Environment*. 311-326. Weinheim VCH.
- Sharma, K.D. & Jain, S. 2020. 'Municipal solid waste generation, composition, and management: The global scenario'. *Social Responsibility Journal* 16(6), 917–948. 10.1108/SRJ-06-2019-0210.
- Sulaiman, M.B., Maigari, A.S. & Danladi, S. 2016. 'Impact of municipal solid waste dumps on surrounding soil and groundwater in Gombe, Nigeria'. *International Journal of Science Environmental* 5(5), 3059-3068.
- Tsheleza, V., Ndhleve, S., Kabiti, H.M., Musampa, C.M. & Nakin, M.D.V., 2019, 'Vulnerability of growing cities to solid waste-related environmental hazards: The case of Mthatha, South Africa', *Jàmá: Journal of Disaster Risk Studies* 11(1), a632. <https://doi.org/10.4102/jamba.v11i1.632>
- Weldeyohanis, Y.H., Aneseyee, A.B. & Sodango, T.H. 2022. 'Evaluation of current solid waste disposal site based on socio-economic and geospatial data: A case study of Wolkite town, Ethiopia'. *Geo Journal* 87, 585–601. <https://doi.org/10.1007/s10708-020-10268-3>

**2. Bukti konfirmasi review dan hasil review
pertama
(27 Mei 2023)**

prof.Slavi Ivanov <scibulcom2@abv.bg>

May 27,
2023,
11:28 PM

to me

Dear authors,

Please find as attachment the letter of acceptance for your paper. The details for the fee (250 Euros) transfer are the following:

Bank account:

Scibulcom 2

Ltd.

SWIFT BIC: STSABGSF

IBAN: BG96STSA93000023294464

DSKBank, k-s Geo Milev

1113 Sofia, Bulgaria

VAT No: BG203908726

Currency: EURO

Thank you in advance

Sincerely yours

Dr. M.Boneva

HEAVY METALS LANDFILL EFFECT ON PEOPLE AND ORGANISM HEALTH RISK: ATOMIC ABSORPTION SPECTROSCOPY APPROACH

ANDI ARTININGSIH^{a*}, SRI WIDODO^b

^aFaculty of Engineering, Universitas Muslim Indonesia, South Sulawesi, Indonesia

E-mail: andi.artiningsih@umi.ac.id

^bFaculty of Engineering, Hasanuddin University, South Sulawesi, Indonesia

ABSTRACT

Some urban city faced to soils around landfill. This study investigates the heavy metals contamination around the Tamangapa Antang Final Disposal Site (FDS). The sample were divided into 4 points consisting of 2 points that align with the groundwater flow (Point A and Point C), and 2 points that are vertical to the groundwater flow (Point B and Point D) in the Antang landfill. Atomic Absorption Spectroscopy (AAS) was used to examine heavy metal around sample observation. The results of the analysis show that the soils of landfill have spread across the topsoil vertically with the highest concentration, reflecting the vast amount of trash dumped to the FDS. The second highest heavy metal concentration was at its highest at Point A and Point D exceeding the limit, but remaining under the limit recommended by the Indonesian Food and Drug Authority at Point B and Point C. Meanwhile, Point A, B, and C heavy metal concentrations were lower. The findings showed that in general, FDS has been contaminated, hence there is a need to concern for regional disaster management, specifically for people which live around the Antang landfill. The construction of regional centres for recycling and reuse activities, and consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities.

Keywords: landfill, soils, vertical, AAS, heavy metals.

The citation of references in the text must be consecutively, by Arabic superscripts with numbers corresponding to those in the final list – we corrected the first 3 sources!

AIMS AND BACKGROUND

The main issue in developing countries is how to select a suitable area for a final disposal site (FDS). Because of the rapid population growth and urbanization, the demand for disposal products

* For correspondence.

has increased by a factor of two¹⁻³ (Mokoele 2023; Njoku et al. 2019; Weldeyohanis et al. 2022). It caused a greater need for landfill space. Hence, an integration approach for waste disposal and waste management plays an important role in reducing the impact of solid waste and heavy metal on the environment and the people who live in and around landfill areas. People who reside nearby landfills are exposed to a number of harmful chemicals that are created during the landfilling process (Sasaki et al. 2019 –Re No???). Litter, dust, excessive rodents, unforeseen landfill fires are some additional contaminants linked to the deposition of trash on landfills. The type and quantity of garbage placed, the age of the landfill, and the climatic conditions of the landfill sites are all factors that affect the by-product or emissions from landfills. Heavy metals, particulate debris, and a number of gaseous contaminants (including dioxins and polycyclic aromatic hydrocarbons) are regularly formed in the landfill as a result of intricate chemical and microbiological processes (Abdelli et al. 2017 –Re No???).

Urbanization is a key issue in developing nations like Indonesia, which has faced various environmental challenges, including the percentage of solid waste, due to social issues including socioeconomic, economic, and population inequality. Traditional solid waste management methods used in Indonesian towns have not been designed to advance environmental protection, public health, or renewable energy sources (Hoque & Rahman 2020; Mester et al. 2022 –Re No???). These methods of managing solid waste are now widely recognised as the primary contributors to contamination of surface and groundwater as well as a degradation in the cleanliness of cities and towns (Morais et al. 2021 –Re No???). Makassar is one of the places, and it deals with concerns including how to manage heavy metal problems in final disposal sites and poorly managed solid waste operations (FDS). Makassar City has been undergoing a significantly rapid development which unfortunately contributes to the increasing amount of waste produced. Waste is an undeniably serious issue which grows with the population in Makassar City.

The Final Disposal Site (FDS) is a vital necessity to create a convenient environment for the community in big cities. If waste from all areas of Big City is not transferred to the FDS even for one day, the city will be full of waste, and it annoys the convenience of the community. It is an undeniable truth that the current conditions of FDS in some big cities have not met the standards of environmental safety. In fact, some FDSs are shut down because they are located too close to people residence and the resulting air pollution from waste deterioration has aggravated the community (Manimekalai & Vijayalakshmi 2014 –Re No???). Poorly managed FDS is critically

harmful to human health and potentially damaging the environment. The human activities have caused serious damage to the environment, and significantly increased biological toxicity of heavy metals in soils. Some heavy metals in excessive concentration may harm the organisms around landfill. The heavy metal contamination also damages soils and animals (Abiola et al. 2021; Chibuike & Obiora 2014; Tsheleza et al. 2021–Re No???).

Soils may be contaminated with heavy metals that are typically without colour or odour, thus are difficult to be controlled. Heavy metal contamination may not instantly be damaging to environment; however, Nawaz et al. (2023) –Re No??? reported that when the concentration of heavy metals has exceeded the tolerance level of the environment, they will be activated and trigger ecological damage. Previous researchers have investigated heavy metal concentration in the soils landfill in Slovakia (Andras et al. 2013–Re No???). Similarly, Esfahani et al. (2021) –Re No??? reported that heavy metals in landfill may be removed by modified membrane.

EXPERIMENTAL

Research area. The administrative area of the research encompassed the regional Bangkala Sub-district at Manggala District, Makassar City, South Sulawesi Province.




Fig. 1. Location map and sampling points taken vertically for Point A, Point B, Point C, and Point D at the depth of 0-10, 10-20, and 20-30 cm, respectively, (4 points, 3- depths????) around Antang Final Disposal Site in Makassar, Sulawesi Selatan – cite in the text!!!

Sampling point. The samples used in this study were the topsoil collected from the area of Antang Final Disposal Site, Makassar City, South Sulawesi. There were four sampling points: two points (Point A and Point C) aligned with the groundwater flow, and two points (Point B and Point D) were vertical to the groundwater flow. All points were measured vertically at the depth of 0-10, 10-20, and 20-30 cm deep. There were a total of 12 samples subjected to analysis of heavy metal Cu, Pb, and Zn using Atomic Absorption Spectroscopy (AAS).

RESULTS AND DISCUSSION

The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point A at the depth of 0-10, 10-20, and 20-30 are presented in Fig. 2.




Fig. 2. Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point A at the depth of 0-10, 10-20, and 20-30

Point A shows that the concentration of Cu at the depth of 0-10 cm is 138 ppm, then increases to 168 and 177 ppm at the depth of 10-20 and 20-30 cm, respectively. The concentration of Pb at 0-10 cm depth is around 104 ppm, then increases to 11 ppm at 10-20 cm depth, and to 105 ppm at 20-30 cm depth. Meanwhile, Zn concentration at 0-10 cm was 128 ppm which decreases to 126 and 118 ppm at the depth of 10-20 and 20-30 cm, respectively. The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point B at the depth of 0-10, 10-20, and 20-30 are presented in Fig. 3.




Fig. 3. Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point B at the depth of 0-10, 10-20, and 20-30 cm

Point B shows that the concentration of Cu at the depth of 0-10 cm is 67 ppm, then increases to 72 ppm (10-20 cm) and decreases to 64 ppm (20-30 cm). Pb concentration at 0-10cm depth is around 52 ppm, then decreases to 43 ppm (10-20 cm) and increases to 63 ppm (20-30 cm). Meanwhile, Zn concentration at 0-10 cm was 61 ppm, then increases to 64 ppm (10-20 cm), and decreases back to 61 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point C at the depth of 0-10, 10-20, and 20-30 are presented in **Fig. 4**.




Fig. 4. Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point C at the depth of 0-10, 10-20, and 20-30 cm

Point C shows that the concentration of Cu at the depth of 0-10 cm is 65 ppm, then increases to 84 ppm (10-20 cm) and decreases to 43 ppm (20-30 cm). Pb concentration at 0-10 cm depth is around 61 ppm, then increases to 91 ppm (10-20 cm) and decreases to 31 ppm (20-30 cm). Meanwhile, Zn concentration at 0-10 cm was 50 ppm, then increases to 58 ppm (10-20 cm), and decreases to 4 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point CD at the depth of 0-10, 10-20, and 20-30 are presented in Fig. 5.




Fig. 5. Results of analysis of heavy metals concentrations of Cu, Pb, and Zn at Point D at the depth of 0-10, 10-20, and 20-30 cm

Point D shows that the concentration of Cu at the depth of 0-10 cm is 158 ppm, then increases to 161 ppm (10-20 cm) and decreases to 149 ppm (20-30 cm). Pb concentration at 0-10 cm depth is around 73 ppm, then decreases to 61 ppm (10-20 cm) and back up again to 73 ppm (20-30 cm). Meanwhile, Zn concentrations at 0-10, 10-20, and 20-30 cm are around 132, 142 and 142 ppm, respectively.

Heavy metals, such as Cu, Pb, and Zn in Antang Final Disposal Site, Makassar City, South Sulawesi have vertically distribution across the topsoil with a variety of concentrations at each depth which has exceeded the limit of contaminant in soil environment, which is acidic. The recommended limit of heavy metal concentrations by the Government of the Republic of Indonesia and Food and Drug Authority is 100 ppm of Cu, 100 ppm of Pb, and 70 ppm of Zn.

Heavy metal Cu is strongly associated with organic matters, iron and manganese oxidation, silica clay and a slight amount of mineral. Cu is particularly absorbed and is slightly mobile in the soils. Organic matter is the dominant factor that controls the retention of Cu. Most Cu is soluble in the soil surface in a complex form of Cu^{2+} . It means that, if Pb heavy metal in the soils functions as Pb^{2+} oxidiser. If pH increases, the Pb^{2+} ion becomes less soluble oxidiser. A higher pH will form a complex form which allows the organic matters settle as carbonate, hydroxide, and phosphate. Lead is one of heavy metals that poses significant harms to living organisms because it is carcinogenic. The recent study found varying levels of Zn and the average pH was indicative of acidity. The oxidation of Zn in the soils shows $^{2+}$??? under oxidised acid conditions. Zn^{2+} is one of the heavy metals that is the most soluble and mobile in the soil, hydrocarbon oxide, and organic matters. Zn solubility in the soils is highly potential at acidic pH (Faborode 2022–Re No???).

The highest concentrations of heavy metals in this study which also have surpassed the limit are tolerance.English!!! High levels of Cu in Final Disposal Site reflect the high quantity of waste dumped to the FDS (Sulaiman et al. 2016–Re No???). The enrichment of heavy metal in the Antang landfill has been highly accumulated because of the anthropogenic activities or human activities, Leachate, and soil characteristics (Bityukova et al. 2000–Re No???) and supported by a very acidic pH, the mobility of acid in heavy metals in the soils is relatively high (Schmitt & Sticker 1991–Re No???).English!!! Heavy metals will be carried by leachate and deposited in the soils, and as a result, heavy metals like Cu, Pb, and Zn will remain deposited in the soils.

Relatively acid soils have an increasing solubility and mobility of nutrients, thus increasing the concentration of heavy metals in the soils (Weldeyohanis et al. 2022; Chibuike & Obiora, 2014–Re No???). If the soil solution is too acidic, plants can not utilise and other nutrients they need to grow. Meanwhile, acidic soils (low pH) are dominated by ion Al, Fe, and Mn which will bind the nutrients vital for the plants, especially P, K S, and Mg. As a result, plants can not absorb their food well despite the richness of nutrients in the soil. In addition, acidic soils contain high level of micro nutrients, which are very dangerous to people's health (Sharma & Jain 2020–Re No???), particularly those living around the landfill because it causes cancer and lung damage in addition to IQ drop among children, and potential death to human in general.

CONCLUSIONS – this section must be shortened and only the main contributions must be outlined!!!

The current study used a multifaceted approach to examine the key elements affecting environmental appropriateness and choosing the best location for the final disposal site (FDS) in the City of Makassar in Indonesia. The determination of sub-criteria and criteria values as well as the choice of the final location utilise a method that combines GIS and an analytical hierarchy process with data from remote sensing and laboratory analysis. Is this true??? The investigation of environmental suitability, comparison of the identified heavy metal and solid waste disposal sites, and final site rankings in accordance with the specified criteria are all included in the study. The issue of sustainable waste management has come into sharper focus as a result of several studies and scientific investigations. The increased manufacturing and widespread availability of different consumer items have an impact on the continuous rise in garbage output. The purpose of this work was to propose a multi-criteria analysis technique that is integrated with geographic information systems and to highlight the benefits that this approach provides for decision-making in environmental studies involving heavy metal analyses. By taking into account many facets of the problem and different stakeholders in the decision-making process, multi-criteria approaches offer a multi-dimensional approach to problems.

Therefore, the methodological properties of multi-criteria analysis, especially from the perspective of environmental planning and environmental management, permit its successful application in environmental and geographical sciences. The use of these approaches and the outcomes attained are in line with previously established waste management practices. Municipal waste management and the effects of heavy metals on the environment and community health in Indonesia can be significantly improved by even just coordinated work and integrated activities of experts, researchers, government and municipal officials, as well as citizens and other stakeholders. The expansion of the municipal trash collection system, the construction of regional centres for recycling and reuse activities, and finding practical and affordable alternatives to locate consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities. In addition to keeping people and the environment safe, this can offer financial payback and the chance to partially recoup maintenance costs for waste management systems. As it gathers useful informational bases required for the development of environmental protection strategies, city development strategies, and information for spatial planning in the Indonesia region, this research can be applied by the city's official agencies for environmental management and protection as well as spatial planning.

The references must be arranged properly according their citation in the text and presented as the first 3 highlighted in yellow sources!!!

REFERENCES

1. N. J. MOKOELE: Complexities of Contemporary Urban Planning in Local Government in the City of Polokwane, Limpopo Province. *Jàmbá J Disaster Risk Stud*, **15** (1), a132 (2023). <https://doi.org/10.4102/jamba.v15i1.1326>.
 2. P. O. NJOKU, J. N. EDOKPAYI, J. O. ODIYO: Health and Environmental Risks of Residents Living Close to a Landfill: a Case Study of Thohoyandou Landfill, Limpopo Province, South Africa. *International Journal of Environmental Research and Public Health ((IJERPH)*, **16**, 1 (2019). <https://doi.org/10.3390/ijerph16122125>.
 3. Y. H. WELDEYOHANIS, A. B. ANESEYEE, T. H. SODANGO: Evaluation of Current Solid Waste Disposal Site Based on Socio-economic and Geospatial Data: a Case Study of Wolkite Town, Ethiopia. *Geo J*, **87**, 585 (2022). <https://doi.org/10.1007/s10708-020-10268-3>.
- Abdelli, I.S., Asnoune, M., Arab, Z., Abdelmalek, F. & Addou, A. 2017. ‘Management of household waste in sanitary landfill of Mostaganem district (Western Algeria)’. *Journal of Material Cycles and Waste Management* 19, 265–281. <https://doi.org/10.1007/s10163-015-0415-6>
- Andras, P., Turisova, L.E., Kharbish, S., Krnac, J. & Cmielova, L. 2013. ‘Environmental risk due to heavy metal contamination caused by old copper mining activity at Lubietova Deposit, Slovakia’. *Energy and Environment Research* 3(2), 1–15. <https://doi.org/10.5539/eer.v3n2p182>
- Bitjukova L, Shogenova, A. & Birke, M. 2000. ‘Urban Geochemistry: A study of element distribution in the soil of Tallin (Estonia)’. *Environmental Geochemistry and Health*. 22; 173-193. <https://doi.org/10.1023/A:1006754326260>

- Chibuikwe, G.U. & Obiora, S.C. 2014. 'Heavy metal polluted soil: Effect on plants and bioremediation methods'. *Hindawi Publishing Corporation; Applied and Environmental soil science*, 1-13. <https://doi.org/10.1155/2014/752708>.
- Esfahani, A.R., Zhai, L. & Sadmani, A.H.M.A. 2021. 'Removing heavy metals from landfill leachate using electrospun polyelectrolyte fiber mat-laminated ultrafiltration membrane'. *Journal of Environmental Chemical Engineering* 9(4), 105355. <https://doi.org/10.1016/j.jece.2021.105355>
- Faborode, H.F.B. 2022. 'Gender assessment of poultry waste utilization among small-scale poultry farmers in Osun state, Nigeria: Exploring the untapped potentials'. *African Journal of Food, Agriculture and Nutrition Development* 22(4), 20259–20279. <https://doi.org/10.18697/ajfand.109.19780>
- Hoque, M.M. & Rahman, M.T.U. 2020. 'Landfill area estimation based on solid waste collection prediction using NN model and final waste disposal options'. *Journal of Cleaner Production* 256, 1–12. <https://doi.org/10.1016/j.jclepro.2020.120387>
- Manimekalai, B. & Vijayalakshmi, P. 2014. 'Analysis of Leachate Contamination Potential of a Municipal Landfill Using Leachate Pollution Index'. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 2(1), 16–39. <https://doi.org/10.9790/2402-0211639>
- Mester, T., Szabó, G., Sajtos, Z., Baranyai, E., Szabó, G. & Balla, D. 2022. 'Environmental hazards of an recultivated liquid waste disposal site on soil and groundwater'. *Water* 14, 1–17. <https://doi.org/10.3390/w14020226>
- Morais, L., Nascimento, V., Simões, S. & Ometto, J. 2021. 'Regional distance routes estimation for municipal solid waste disposal, case study São Paulo State, Brazil'. *Energies* 14, 1–14. <https://doi.org/10.3390/en14133964>
- Nawaz, R., Aslam, M., Nasim, I., Irshad, M.A., Ahmad, S., Latif, M. & Hussain, F. 2023. 'Air Pollution Tolerance Index and Heavy Metals Accumulation of Tree Species for Sustainable Environmental Management in Megacity of Lahore'. *Air* 1(1), 55-68. <https://doi.org/10.3390/air1010004>

- Parth, V., Murthy, N.N. & Raj Saxena, P. 2011. 'Assessment of heavy metal contamination in soil around hazardous waste disposal sites in Hyderabad city (India): natural and anthropogenic implications'. *E3 Journal of Environmental and Research Managagement* 2(2), 27-34.
- Sasaki, S., Watanabe, K., Widyaningsih, N. & Araki, T. 2019. 'Collecting and dealing of recyclables in a final disposal site and surrounding slum residence: The case of Bantar Gebang, Indonesia'. *Journal of Material Cycles and Waste Management* 21, 375–393. <https://doi.org/10.1007/s10163-018-0798-2>
- Schmitt, H.S. & Sticker, H. 1991. 'Heavy metal compounds in the soil'. Merian, E. (Ed.), *Metals and their compounds in the Environment*. 311-326. Weinheim VCH.
- Sharma, K.D. & Jain, S. 2020. 'Municipal solid waste generation, composition, and management: The global scenario'. *Social Responsibility Journal* 16(6), 917–948. 10.1108/SRJ-06-2019-0210.
- Sulaiman, M.B., Maigari, A.S. & Danladi, S. 2016. 'Impact of municipal solid waste dumps on surrounding soil and groundwater in Gombe, Nigeria'. *International Journal of Science Environmental* 5(5), 3059-3068.
- Tsheleza, V., Ndhleve, S., Kabiti, H.M., Musampa, C.M. & Nakin, M.D.V., 2019, 'Vulnerability of growing cities to solid waste-related environmental hazards: The case of Mthatha, South Africa', *Jàmbá: Journal of Disaster Risk Studies* 11(1), a632. <https://doi.org/10.4102/jamba.v11i1.632>

Received 29 April 2023
Revised

**3. Bukti konfirmasi submit revisi pertama,
tanggapan kepada saran reviewer dan
naskah yang di submit
(9 Juni 2023)**

Andi Artiningsih <andi.artiningsih@umi.ac.id>

Jun 9,
2023,
7:48 PM

to prof.Slavi

Dear Professor Slavi Ivanov

We would like to thank the Editor and Reviewers for their detailed and helpful feedback regarding our manuscript, and for providing us with the opportunity to revise our manuscript. We have revised the manuscript according to the Reviewers' recommendations. If there were specific ways, you would like us to address any remaining concerns, please let us know.

Sincerely

HEAVY METALS LANDFILL EFFECT ON PEOPLE AND ORGANISM HEALTH RISK: ATOMIC ABSORPTION SPECTROSCOPY APPROACH

ANDI ARTININGSIH^{a*}, SRI WIDODO^b

^a*Chemical Engineering Department, Faculty of Industrial Engineering, Universitas Muslim Indonesia, South Sulawesi, Indonesia*

E-mail: andi.artiningsih@umi.ac.id

^b*Mining Engineering Department, Faculty of Engineering, Hasanuddin University, South Sulawesi, Indonesia*

ABSTRACT

Some urban city faced to soils around landfill. This study investigates the heavy metals contamination around the Tamangapa Antang Final Disposal Site (FDS). The sample were divided into 4 areas consisting of 2 locations that align with the groundwater flow (area A and area C), and 2 locations that are vertical to the groundwater flow (area B and area D) in the Antang landfill. Atomic Absorption Spectroscopy (AAS) was used to examine heavy metal around sample observation. The results of the analysis show that the soils of landfill have spread across the topsoil vertically with the highest concentration, reflecting the vast amount of trash dumped to the FDS. The second highest heavy metal concentration was at its highest at area A and area D exceeding the limit, but remaining under the limit recommended by the Indonesian Food and Drug Authority at area B and area C. Meanwhile, area A, B, and C heavy metal concentrations were lower. The findings showed that in general, FDS has been contaminated, hence there is a need to concern for regional disaster management, specifically for people which live around the Antang landfill. The construction of regional centres for recycling and reuse activities, and consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities.

Keywords: landfill, soils, vertical, AAS, heavy metals.

AIMS AND BACKGROUND

* For correspondence.

The main issue in developing countries is how to select a suitable area for a final disposal site (FDS). Because of the rapid population growth and urbanization, the demand for disposal products has increased by a factor of two¹⁻³. It caused a greater need for landfill space. Hence, an integration approach for waste disposal and waste management plays an important role in reducing the impact of solid waste and heavy metal on the environment and the people who live in and around landfill areas. People who reside nearby landfills are exposed to a number of harmful chemicals that are created during the landfilling process⁴⁻⁵. Litter, dust, excessive rodents, unforeseen landfill fires are some additional contaminants linked to the deposition of trash on landfills. The type and quantity of garbage placed, the age of the landfill, and the climatic conditions of the landfill sites are all factors that affect the by-product or emissions from landfills. Heavy metals, particulate debris, and a number of gaseous contaminants (including dioxins and polycyclic aromatic hydrocarbons) are regularly formed in the landfill as a result of intricate chemical and microbiological processes⁶⁻⁷.

Urbanization is a key issue in developing nations like Indonesia, which has faced various environmental challenges, including the percentage of solid waste, due to social issues including socioeconomic, economic, and population inequality. Traditional solid waste management methods used in Indonesian towns have not been designed to advance environmental protection, public health, or renewable energy sources⁸⁻⁹. These methods of managing solid waste are now widely recognised as the primary contributors to contamination of surface and groundwater as well as a degradation in the cleanliness of cities and towns¹⁰⁻¹¹. Makassar is one of the places, and it deals with concerns including how to manage heavy metal problems in final disposal sites and poorly managed solid waste operations (FDS). Makassar City has been undergoing a significantly rapid development which unfortunately contributes to the increasing amount of waste produced. Waste is an undeniably serious issue which grows with the population in Makassar City.

The Final Disposal Site (FDS) is a vital necessity to create a convenient environment for the community in big cities. If waste from all areas of Big City is not transferred to the FDS even for one day, the city will be full of waste, and it annoys the convenience of the community. It is an undeniable truth that the current conditions of FDS in some big cities have not met the standards of environmental safety. In fact, some FDSs are shut down because they are located too close to people residence and the resulting air pollution from waste deterioration has aggravated the community¹². Poorly managed FDS is critically harmful to human health and potentially damaging

the environment. The human activities have caused serious damage to the environment, and significantly increased biological toxicity of heavy metals in soils. Some heavy metals in excessive concentration may harm the organisms around landfill. The heavy metal contamination also damages soils and animals¹³⁻¹⁴.

Soils may be contaminated with heavy metals that are typically without colour or odour, thus are difficult to be controlled. Heavy metal contamination may not instantly be damaging to environment; however, ¹⁵⁻¹⁶ reported that when the concentration of heavy metals has exceeded the tolerance level of the environment, they will be activated and trigger ecological damage. Previous researchers have investigated heavy metal concentration in the soils landfill in Slovakia¹⁷. Similarly, heavy metals in landfill may be removed by modified membrane¹⁸.

EXPERIMENTAL

Research area. The administrative area of the research encompassed the regional Bangkala Sub-district at Manggala District, Makassar City, South Sulawesi Province.




Figure. 1: Location map Antang landfill, Makassar city, South Sulawesi, Indoensia

Sampling point. The samples used in this study were the topsoil collected from the area of Antang Final Disposal Site, Makassar City, South Sulawesi. There were four sampling areas: two areas (A and C) aligned with the groundwater flow, and area B and D were vertical to the groundwater flow. All points were measured vertically at the depth of 0-10, 10-20, and 20-30 centimetres deep.

RESULTS AND DISCUSSION

The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area A at the depth of 0-10, 10-20, and 20-30 are presented in **Figure. 2**.




Figure. 2. Results of analysis of heavy metals concentrations of area A

Area A shows that the concentration of Cu at the depth of 0-10 cm is 138 ppm, then increases to 168 and 177 ppm at the depth of 10-20 and 20-30 cm, respectively. The concentration of Pb at 0-10 cm depth is around 104 ppm, then increases to 111 ppm at 10-20 cm depth, and to 105 ppm at 20-30 cm depth. Meanwhile, Zn concentration at 0-10 cm was 128 ppm which decreases to 126 and 118 ppm at the depth of 10-20 and 20-30 cm, respectively. The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area B at the depth of 0-10, 10-20, and 20-30 are presented in **Figure. 3**.




Figure. 3. Results of analysis of heavy metals concentrations of area B

Area B shows that the concentration of Cu at the depth of 0-10 cm is 67 ppm, then increases to 72 ppm (10-20 cm) and decreases to 64 ppm (20-30 cm). Pb concentration at 0-10cm depth is around 52 ppm, then decreases to 43 ppm (10-20 cm) and increases to 63 ppm (20-30 cm). Meanwhile, Zn concentration at 0-10 cm was 61 ppm, then increases to 64 ppm (10-20 cm), and decreases back to 61 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area C at the depth of 0-10, 10-20, and 20-30 are presented in [Figure. 4](#).




Fig. 4. Results of analysis of heavy metals concentrations of area C

Area C shows that the concentration of Cu at the depth of 0-10 cm is 65 ppm, then increases to 84 ppm (10-20 cm) and decreases to 43 ppm (20-30 cm). Pb concentration at 0-10 cm depth is 65 ppm, then increases to 84 ppm (10-20 cm) and decreases to 30 ppm (20-30 cm). Zn concentration at 0-10 cm was 50 ppm, then increases to 58 ppm (10-20 cm), and decreases to 48 ppm (20-30 cm).

around 61 ppm, then increases to 91 ppm (10-20 cm) and decreases to 31 ppm (20-30 cm). Meanwhile, Zn concentration at 0-10 cm was 50 ppm, then increases to 58 ppm (10-20 cm), and decreases to 4 ppm (20-30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area CD at the depth of 0-10, 10-20, and 20-30 are presented in **Figure. 5**.




Figure. 5. Results of analysis of heavy metals concentrations of area D

Area D shows that the concentration of Cu at the depth of 0-10 cm is 158 ppm, then increases to 161 ppm (10-20 cm) and decreases to 149 ppm (20-30 cm). Pb concentration at 0-10 cm depth is around 73 ppm, then decreases to 61 ppm (10-20 cm) and back up again to 73 ppm (20-30 cm). Meanwhile, Zn concentrations at 0-10, 10-20, and 20-30 cm are around 132, 142 and 142 ppm, respectively.

Heavy metals, such as Cu, Pb, and Zn in Antang Final Disposal Site, Makassar City, South Sulawesi have vertically distribution across the topsoil with a variety of concentrations at each depth which has exceeded the limit of contaminant in soil environment, which is acidic. The recommended limit of heavy metal concentrations by the Government of the Republic of Indonesia and Food and Drug Authority is 100 ppm of Cu, 100 ppm of Pb, and 70 ppm of Zn.

Heavy metal Cu is strongly associated with organic matters, iron and manganese oxidation, silica clay and a slight amount of mineral. Cu is particularly absorbed and is slightly mobile in the soils. Organic matter is the dominant factor that controls the retention of Cu. Most Cu is soluble in the soil surface in a complex form of Cu^{2+} . It means that, if Pb heavy metal in the soils functions as Pb^{2+} oxidiser. If pH increases, the Pb^{2+} ion becomes less soluble oxidiser. A higher pH will form

a complex form which allows the organic matters settle as carbonate, hydroxide, and phosphate. Lead is one of heavy metals that poses significant harms to living organisms because it is carcinogenic. The recent study found varying levels of Zn and the average pH was indicative of acidity. The oxidation of Zn in the soils shows Zn^{2+} under oxidised acid conditions. Zn^{2+} is one of the heavy metals that is the most soluble and mobile in the soil, hydrocarbon oxide, and organic matters. Zn solubility in the soils is highly potential at acidic pH¹⁹.

The concentrations of heavy metals in this study which also have surpassed the limit are tolerance. High levels of Cu in Final Disposal Site reflect the high quantity of waste dumped to the FDS²⁰. The enrichment of heavy metal in the Antang landfill has been highly accumulated because of the anthropogenic activities or human activities, Leachate, and soil characteristics and supported by a very acidic pH, the mobility of acid in heavy metals in the soils is relatively high²¹. Heavy metals will be carried by leachate and deposited in the soils, and as a result, heavy metals like Cu, Pb, and Zn will remain deposited in the soils.

Relatively acid soils have an increasing solubility and mobility of nutrients, thus increasing the concentration of heavy metals in the soils²². If the soil solution is too acidic, plants can not utilise and other nutrients they need to grow. Meanwhile, acidic soils (low pH) are dominated by ion Al, Fe, and Mn which will bind the nutrients vital for the plants, especially P, K S, and Mg. As a result, plants can not absorb their food well despite the richness of nutrients in the soil. In addition, acidic soils contain high level of micro nutrients, which are very dangerous to people's health²³, particularly those living around the landfill because it causes cancer and lung damage in addition to IQ drop among children, and potential death to human in general.

CONCLUSIONS

The increased manufacturing and widespread availability of different consumer items have an impact on the continuous rise in garbage output. Municipal waste management and the effects of heavy metals on the environment and community health in Indonesia and developing countries can be significantly improved by even just coordinated work and integrated activities of experts, researchers, government and municipal officials, as well as citizens and other stakeholders. The expansion of the municipal trash collection system, the construction of regional centres for recycling and reuse activities, and finding practical and affordable alternatives to locate consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic

priorities. In addition to keeping people and the environment safe, this can offer financial payback and the chance to partially recoup maintenance costs for waste management systems. As it gathers useful informational bases required for the development of environmental protection strategies, city development strategies, and information for spatial planning in the Indonesia region, this research can be applied by the city's official agencies for environmental management and protection as well as spatial planning.

REFERENCES

1. N. J. MOKOELE: Complexities of Contemporary Urban Planning in Local Government in the City of Polokwane, Limpopo Province. *Jàmbá J Disaster Risk Stud*, **15** (1), a132 (2023). <https://doi.org/10.4102/jamba.v15i1.1326>.
2. P. O. NJOKU, J. N. EDOKPAYI, J. O. ODIYO: Health and Environmental Risks of Residents Living Close to a Landfill: A Case Study of Thohoyandou Landfill, Limpopo Province, South Africa. *Int J Environmental Research Public Health (IJERPH)*, **16**, 1 (2019). <https://doi.org/10.3390/ijerph16122125>.
3. Y. H. WELDEYOHANIS, A. B. ANESEYEE, T. H. SODANGO: Evaluation of Current Solid Waste Disposal Site Based on Socio-economic and Geospatial Data: a Case Study of Wolkite Town, Ethiopia. *Geo J*, **87**, 585 (2022). <https://doi.org/10.1007/s10708-020-10268-3>.
4. S. SASAKI, K. WATANABE, N. WIDYANINGSIH, T. ARAKI: Collecting and dealing of recyclables in a final disposal site and surrounding slum residence: The case of Bantar Gebang, Indonesia. *J Material Cycles Waste Management*, **21**, (2019). <https://doi.org/10.1007/s10163-018-0798-2>
5. M. PAN, H. LI, J. WU: Study on Catalysed Treatment of Coking Wastewater by TIO₂. *Oxidation Communications*, **39**, 4, 3457–3461 (2016).
6. I. S. ABDELLI, M. ASNOUNE, Z. ARAB, F. ABDELMALEK, A. ADDOU: Management of household waste in sanitary landfill of Mostaganem district (Western Algeria). *J Material Cycles and Waste Management*, **19**, 265-281. (2017). <https://doi.org/10.1007/s10163-015-0415-6>

7. W. XIAOYUN, Z. YOUFEI, L. SHENGTAO, L. KESI: Research Status of Methane Emission in Landfill Sites. *Oxidation Communications*, **39**, 4, 3090-3097 (2016).
8. M. M. HOQUE, M. T. U. RAHMAN: Landfill area estimation based on solid waste collection prediction using NN model and final waste disposal options. *J Cleaner Production*, **256**, 1-12 (2020). <https://doi.org/10.1016/j.jclepro.2020.120387>
9. T. MESTER, G. SZABÓ, Z. SAJTOS, E. BARANYAI, G. SZABÓ, D. BALLA: Environmental hazards of a recultivated liquid waste disposal site on soil and groundwater. *Water*, **14**, 1-17 (2022) <https://doi.org/10.3390/w14020226>
10. L. MORAIS, V. NASCIMENTO, S. SIMÕES, J. OMETTO: Regional distance routes estimation for municipal solid waste disposal, case study São Paulo State, Brazil. *Energies*, **14**, 1-14 (2021). <https://doi.org/10.3390/en14133964>
11. R. POBLETE, N. K. SALIHOGLU, G. SALIHOGLU: Landfill Leachate Treatment Using Advanced Oxidation Processes. *Oxidation Communications*, **39**, 1, 493-502 (2016).
12. B. MANIMEKALAI, P. VIJAYALAKSHMI: Analysis of Leachate Contamination Potential of a Municipal Landfill Using Leachate Pollution Index. *IOSR J Environmental Science, Toxicology and Food Technology*, **2**, 1, 16-39 (2014). <https://doi.org/10.9790/2402-0211639>
13. G. U. CHIBUIKE, S. C. OBIORA: Heavy metal polluted soil: Effect on plants and bioremediation methods. *Hindawi Pub Corp; Applied and Environmental soil science*, 1-13 (2014). <https://doi.org/10.1155/2014/752708>.
14. V. TSHELEZA, S. NDHLEVE, H. M. KABITI, C. M. MUSAMPA, M. D. V. NAKIN: Vulnerability of growing cities to solid waste-related environmental hazards: The case of Mthatha, South Africa, *Jamba J Disaster Risk Stud*, **11**, 1, a632 (2019). <https://doi.org/10.4102/jamba.v11i1.632>
15. R. NAWAZ, M. ASLAM, I. NASIM, M. A. IRSHAD, S. AHMAD, M. LATIF, F. HUSSAIN: Air Pollution Tolerance Index and Heavy Metals Accumulation of Tree Species for Sustainable Environmental Management in Megacity of Lahore. *Air*, **1**, 1, 55-68 (2023). <https://doi.org/10.3390/air1010004>

16. P. LIU, C. WANG, M. HAO, Z. XU, Y. LI: Treatment of Oily Wastewater by the Fenton Reagent and Wet Hydrogen Peroxide Oxidation. *Oxidation Communications*, **39**, 2666-2675 (2016).
17. P. ANDRAS, L. E. TURISOVA, S. KHARBISH, J. KRNAC, L. CMIELOVA: Environmental Risk due to Heavy Metal Contamination Caused by Old Copper Mining Activity at Lubietova Deposit, Slovakia. *Energy and Environment Research*, **3**, 2, 1-15 (2016).
<https://doi.org/10.5539/eer.v3n2p182>
18. A. R. ESFAHANI, L. ZHAI, A. H. M. A. SADMANI: Removing Heavy Metals from Landfill Leachate using Electrospun Polyelectrolyte Fiber Mat-laminated Ultrafiltration Membrane. *J Env Chemical Engineering*, **9**, 4, 105355 (2021).
<https://doi.org/10.1016/j.jece.2021.105355>
19. F. F. B. FABORODE: Gender Assessment of Poultry Waste Utilization among Small-scale Poultry Farmers in Osun State, Nigeria: Exploring the Untapped Potentials. *African J of Food, Agriculture and Nutrition Development*, **22**, 4, 20259-20279 (2022).
<https://doi.org/10.18697/ajfand.109.19780>
20. M. B. Sulaiman, A. S. Maigari, S. Danladi: Impact of Municipal Solid Waste Dumps on Surrounding Soil and Groundwater in Gombe, Nigeria. *International J of Science Environmental*, **5**, 5, 3059-3068 (2016).
21. L. BITYUKOVA, A. SHOGENOVA, M. BIRKE: Urban Geochemistry: A Study of Element Distribution in the Soil of Tallin (Estonia). *Environmental Geochemistry and Health*, **22**, 173-193 (2000). <https://doi.org/10.1023/A:1006754326260>
22. I. Sazykin, L. Khmelevtsova, T. Azhogina, M. Sazykina: Heavy Metals Influence on the Bacterial Community of Soils: A Review. *Agriculture*, **13**, 3, 653 (2023).
<https://doi.org/10.3390/agriculture13030653>
23. K. D. Sharma, S. Jain: Municipal solid waste generation, composition, and management: The global scenario. *Social Responsibility Journal*, **16**, 6, 917-948 (2020). 10.1108/SRJ-06-2019-0210.

Received 29 April 2023
Revised

4. Bukti konfirmasi naskah diterima (3 Juli 2023)



prof.Slavi Ivanov <scibulcom2@abv.bg>

Jul 3, 2023,
11:03 PM

to me

Dear authors,

Please find as attachments the pdf.file of your paper, published in book 2 (2023) of Oxidation Communications, together with the covers of this issue.

In case you have some corrections concerning the title and the authors of the paper, we will insert them within 3 days from today, till 6th of July, 2023. Otherwise they will be noted as Errata in the next issue.

Thank you for your co-operation.

Sincerely yours

Dr. M.Boneva

5. Bukti konfirmasi naskah dipublish (5 Agustus 2023)

CONTENTS

Vol. 46 (2), 2023

Review article

- KARUNA CHOURASIA, NEHA MISHRA, MITHUN KORI, TAKKALUM KHAN, RITU YADAV. A Mini Review on Synthetic Route for Bioactive Heterocyclic Schiff Base Ligands .. 249
K. STANCHEVA, C. PASHA. Coulometry: Fundamentals and Applications 261

Molecular docking analysis

- S. PANDIYAN ARUN, R. THIRUCHELVI, K. RAJAKUMARI. *In silico* Studies of the Hepatitis B Viral Protein for Drug Discovery 272
V. K. SUKANYA, K. RAJAKUMARI, S. IVO ROMAULD. Molecular Docking and ADMET Studies of Bioactive Compounds of *Verbascum thapsus* against the Bacterial Protein CagA 283

Oxidative stress and public health

- N. NALINI, R. SIVARAMAN, K. BINDU KUMAR, MAMTA BHAT, D. ANAND, S. ARUNA KUMARI, D. SRAVANI. Impact of Oxidative Stress on Cellular Communication and Its Implications for Diseases 293

Allelopathic activity of medicinal plants

- A. K. SINGH, R. N. YADAVA. Allelopathic Effects of Stems Extract of *Cassia alata* Linn.... 303

Phytochemical analysis and antidiabetic activity

- K. KAVITHA, J. PRIYADHARSHINI, VIVEK PAZHAMALAI, S. IVO ROMAULD. Phytochemical Analysis and Antidiabetic Activity of *Mangifera indica* and *Ricinus communis*..... 309

Biological and antimicrobial activity

- I. S. CHAKRAPANI, S. ANISREE, V. KARTHIKA, VINJAMURI VENKATA KAMESH, NAGESWARA RAO MEDIKONDU, Y. ISMAIL, M. AARTHI. Exploring the Bioactive Potential, Antimicrobial Properties and Toxicity of Peptides and Proteins from *Nigella sativa* for Drug Discovery 318

Biological and biochemical oxidation processes and activity; medical image processing; neural networks and fuzzy logic techniques

- A. JOEL DICKSON, J. ARUL LINSELY, R. J. ALICE NINETA. Abnormality Detection of CT Liver Images Using Mask Recurrent Convolutional Neural Network Approach..... 328

Synthesis of a new type – sulphur-containing glycosides

- N. N. SIDAMONIDZE, L. V. TABATADZE, R. O. VARDIASHVILI, M. O. NUTSUBIDZE, D. J. IREMASHVILI, E. I. CHACHUA, N. G. SHENGELIA. Hydrosulphurisation Reactions of Allyl Derivatives of Altrapranose with Mercaptans 337

Synthesis of sodium selenate dihydrogen selenite

- TSVETAN TSENOV, TSVETELINA YOTOVA, RUMYANA YANKOVA. Hirshfeld Surface and Reduced Density Gradient Analyses of Sodium Selenate Dihydrogen Selenite..... 345

Synthesis of organo-tellurium compounds

- G. G. KHAMIS, A. M. M. ALI, M. G. TOPUZOVA. Studying the Chemical Reactivity of 2-Methylsulphanyl[1,3]benzotellurazole 356

Synthesis of materials for organic electronics

- S. SIVARAMAKRISHNAN, R. AGILESH SARAVANAN, M. THIYAGESAN, VINJAMURI VENKATA KAMESH, J. SUMATHI, NARENDRA KUMAR, S. KIRUBA. Hexabenzocoronene: a Material with Great Potential for Eco-friendly, High-performance Optoelectronic Devices 365

- A. SARAVANAN, D. LAKSHMI, K. SASIREKHA, M. J. SUGANYA, S. N. PADHI, SAMPURNA PANDA, N. RAJESHA. Examining the Eco-friendly Possibilities of Hexabenzocoronene as a Component in Field-effect Transistors and Solar Cells in Organic Electronics 376

- K. RAJALAKSHMI, R. SAKTHIVEL, C. THANGAMANI, J. SIVAPRIYA, D. RAVINDRAN, SULTANA RAZIA, R. B. R. PRAKASH. Exploring the Environmental Benefits of Utilising Dibenzotetrabenzoporphyrin and Metal Nanoparticles in Organic Solar Cells..... 386

ISSN 0209-4541

see b.c.

oranj black

oranj black

Vol. 46 No 2 2023

Vol. 46 No 2 2023



OXIDATION
COMMUNICATIONS

Founding Editor

D. GAL, Hungary

Editor-in-Chief

S. K. IVANOV, Bulgaria

Deputy Editor-in-Chief

Zh. D. KALITCHIN, Bulgaria

Editor

M. I. BONEVA, Bulgaria

Editorial Board

R. L. AUGUSTINE, USA

S. W. BENSON, USA

S. BOURBIGOT, France

D. BRADLEY, UK

A. M. BRAUN, Germany

J. DAHLMANN, Germany

A. D'AMORE, Italy

S. DOBE, Hungary

Sh. GAO, China

N. GETOFF, Austria

V. K. GUPTA, India

J. HAPPEL, USA

J. A. HOWARD, Canada

M. F. R. MULCAHY, Australia

A. NEMETH, Hungary

E. NIKI, Japan

K. OHKUBO, Japan

P. St. PETKOV, Bulgaria

A. RAJARAM, India

R. A. SHELDON, The Netherlands

R. SILAMBARASAM, India

J. J. ZIOLKOWSKI, Poland

ZUWEI Xi, China

Scientific
Bulgarian
communications
Ltd., Co.

Oxidation Communications

oranj black

oranj black



OXIDATION COMMUNICATIONS is the only scientific journal which, in its new form, is devoted to the global oxy-reduction interactions proceeding in nature.

OXIDATION COMMUNICATIONS acts as an international focus for modern fundamental and technological research in this rapidly developing field of science of exclusive importance for life.

The main topics and examples of specific areas of interest to the Journal are:

- (a) gas-, liquid-, and solid-phase oxidation processes; oxygen and hydrogen peroxide as oxidisers, ion-pair interactions, dipole-dipole interactions, π - π interactions, hydrogen bonding;
- (b) mechanism of action of inhibitors during the stabilisation of fuels, lubricating oils and greases, polymer materials and foodstuffs;
- (c) oxidation reactions in the presence of homogeneous, heterogeneous, immobilised and enzymic catalytic systems, combustion, fires, explosions;
- (d) regulators of selective oxidation synthesis of valuable oxygen-containing compounds; highly enantioselective oxidation; stereoselective synthesis;
- (e) biological and biochemical oxidation processes; medical image processing; neural networks and fuzzy logic techniques;
- (f) photochemical oxy-reduction interactions;
- (g) methods for prognosis and evaluation of performance terms or/and storage terms of the materials, as mentioned above in (b);
- (h) tribochemical interactions on metal and other surfaces in the presence of molecular oxygen or its activated species;
- (i) methods for solar and chemical energy accumulation, methods of high performance liquid chromatography (HPLC), gas chromatography (GC), capillary electrophoresis (CE), IR and NMR spectroscopy and other spectroscopic and radio-spectroscopic investigations of oxy-reduction processes;
- (j) studies on the nature of intermediates, generated in the course of oxidation;
- (k) technological aspects of oxidation processes of special interest to the petrochemistry, pharmaceuticals, food and agrochemical industries; engineering applications and information sciences; civil and mechanical engineering;
- (l) combustion and gasification of natural, synthetic and biological fuels, fires, etc.
- (m) ecologically friendly oxidative processes for the atmosphere, soils and water, environmental contaminants;
- (n) educational problems of environmental protection associated with oxidation processes;
- (o) European legislation related to oxy-reduction and synthesis processes; management.

The term 'oxidation' is interpreted widely, including both organic and inorganic compounds as oxidising agents during the oxidation processes of organic and inorganic substrates and other electronic transfer reactions.

The interdisciplinary nature of the Journal ensures appeal to direct exchange of ideas and contributions among academic and industrial scientists.

The Journal is publishing refereed authoritative review articles, papers covering original studies and also short communications, including preliminary results of a particular interest.

Designed to: chemists, researchers in the field of physical chemistry, petrochemistry, organic synthesis, catalysis, molecular biology and biochemistry, tribochemistry, polymer and engineering chemistry.

The Journal is covered and evaluated in Elsevier products – Reaxys, Scopus, SJR, SCImago Journal & Country Rank and in China National Knowledge Infrastructure (CNKI). The Journal is included in EBSCOhost: Academic Search Complete, Academic Search Ultimate, The Belt and Road Initiative Reference Source. The SJR of the journal for 2021 is 0.216. The h-index is 22 (2021) and the Impact score (Scopus) is 0.527 (2021).

Compiled by S. K. Ivanov, Zh. D. Kalitchin, M. I. Boneva, J. P. Ivanova, N. Evtimova and V. Shishkova. Volume 46 (2023) will be published in four separate books, issued at 31.03.2023, 30.06.2023, 30.09.2023 and 31.12.2023. The editorial board can publish special issues in case of over-accumulation of manuscripts submitted for publication. Regular subscription rate: EUR 800/Europe and US\$ 942/rest, 10% agency discount, plus extra postage charges: for Europe EUR 30 without priority mail and EUR 60 with priority mail; for all other countries US\$ 50 without priority mail and US\$ 90 with priority mail, including packing, handling and postage. Bank charges are shared according to the banks requirements. Prices are subject to change without notice, in accordance with market fluctuations. Advertising and announcements are accepted and published in the Journal at US\$ 3 per 1 cm².

Subscription orders should be sent to:

Prof. Dr. Slavi K. Ivanov – SciBulCom Ltd., P. O. Box 249, 7 Nezabavka Str., 1113 Sofia, Bulgaria, www.scibulcom.net; e-mail: scibulcom2@abv.bg.

ISSN 0209-4541

Synthesis of nanopolymers

- R. VEZHAVENDHAN, M. HARINATHA REDDY, NAGESWARA RAO MEDIKONDU, N. PADMAMALINI, Y. ISMAIL, AMOS GAMALEAL DAVID, G. PUTHILIBAI. Dithieno[3,2-b:2',3'-d] Pyrrole: a Green Alternative for Stabilising the Synthesis of Nano Polymers 395

Atomic absorption spectroscopy analysis

- ANDI ARTININGSIH, SRI WIDODO. Heavy Metals Landfill Effect on People and Organism Health Risk: Atomic Absorption Spectroscopy Approach..... 405

Spectral analysis and characterisation

- S. S. HAYRAPETYAN, D. A. MARTIROSYAN, A. A. GRIGORYAN, L. S. HAYRAPETYAN, A. R. SUKIASYAN. Identification of Phytosterols in *Coriandrum sativum* L. Seeds as Function of Their Germination 413

Microwave drying modelling of rainbow trout

- L. DOSPATLIEV, N. KATRANZHEV, M. IVANOVA, M. VARBANOV. Microwave Drying Modelling of Rainbow Trout 420

Risk assessment of mining activity on water quality

- L. A. MARGARYAN, A. M. KHACATRYAN, A. G. SIMONYAN, G. S. SIMONYAN. Environmental Risk Assessment of Kapan Community 429

Technological aspects of oxidation processes of special interest to petrochemistry, pharmaceuticals, food and agrochemical industries; engineering applications and information sciences; civil and mechanical engineering

- Y. TASHEVA, E. DIMITROV, L. KUNCHEV. Exploration Comparison of Contemporary Diesel Blends under Effective Performance of Engine 436

Electrochemical behaviour of aluminum alloy

- K. KAMARSKA. Electrochemical Behaviour of Aluminum Alloy EN AW-2024 in Citric Acid..... 445

Ceramic material and physicochemical characterisation

- A. A. GEORGIEVA, F. S. YOVKOVA, M. A. GEORGIEVA, K. Z. PANAYOTOVA. Obtainment of Barium Titanate Ceramic Samples and Determination of Their Basic Physicochemical Properties 450

Sanitary microbiology studies

- N. VALCHEVA, Y. KOLEVA. Characteristics of Three Bacterial Strains Isolated from Water Sources in Sofia Region, Bulgaria 459

Environmentally friendly processes

- C. RAGAVI, R. RAVINA DEVI, S. ARUN, P. SAKTHISELVAN, S. IVO ROMAULD, K. RAJAKUMARI, VIVEK PAZHAMALAI. Removal of Malachite Green Dye Using Activated Carbon Obtained from *Prosopis juliflora* 465

- E. M. ANJU, G. R. VIJAY SHANKAR. Changes of Geotechnical Properties of Heavy Metal Contaminated Soil after Solidification 477

- G. VINOTH KANNA, A. VIJAYALAKSHMI, R. JOTHILAKSHMI, M. JOGENDRA KUMAR, J. SUMATHI, M. RAJKUMAR, ASHOK KUMAR KOSHARIYA. Enhancing Rice Yields in Red Soil through Azotobacter Cultivation..... 489

- R. VEZHAVENDHAN, PRAKASH BABU KANAKAVALLI, SRI HARSHA ARIGELA, A. CHANDRASHEKHAR, R. SATHISH, SANJEEV KUMAR JOSHI, J. SAMSON ISAAC. Nanofluids in Production of Sustainable Biofuels and Bioplastics..... 499

- M. PETROV, Z. NIKOLAEVA, AL. DIMITROV. Cyclical Aspect of the Complex Factors of Anthropogenic Activity on the Global Ecosystem..... 509

- M. PETROV, Z. NIKOLAEVA, AL. DIMITROV. Spontaneous Global Wildfires as the Consequence of Climate Variability 519

HEAVY METALS LANDFILL EFFECT ON PEOPLE AND ORGANISM HEALTH RISK: ATOMIC ABSORPTION SPECTROSCOPY APPROACH

ANDI ARTININGSIH^{a*}, SRI WIDODO^b

^a*Chemical Engineering Department, Faculty of Industrial Engineering, Universitas Muslim Indonesia, South Sulawesi, Indonesia*

E-mail: andi.artiningsih@umi.ac.id

^b*Mining Engineering Department, Faculty of Engineering, Hasanuddin University, South Sulawesi, Indonesia*

ABSTRACT

Some urban city faced to soils around landfill. This study investigates the heavy metals contamination around the Tamangapa Antang Final Disposal Site (FDS). The samples were divided into 4 points consisting of 2 points that align with the groundwater flow (Point A and Point C), and 2 points that are vertical to the groundwater flow (Point B and Point D) in the Antang landfill. Atomic Absorption Spectroscopy (AAS) was used to examine heavy metal around sample observation. The results of the analysis show that the soils of landfill have spread across the topsoil vertically with the highest concentration, reflecting the vast amount of trash dumped to the FDS. The second highest heavy metal concentration was at its highest at Point A and Point D exceeding the limit, but remaining under the limit recommended by the Indonesian Food and Drug Authority at Point B and Point C. Meanwhile, Point A, B, and C heavy metal concentrations were lower. The findings showed that in general, FDS has been contaminated, hence there is a need to concern for regional disaster management, specifically for people which live around the Antang landfill. The construction of regional centres for recycling and reuse activities, and consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities.

Keywords: landfill, soils, vertical, AAS, heavy metals.

AIMS AND BACKGROUND

The main issue in developing countries is how to select a suitable area for a final disposal site (FDS). Because of the rapid population growth and urbanization, the

* For correspondence.

demand for disposal products has increased by a factor of two¹⁻³. It caused a greater need for landfill space. Hence, an integration approach for waste disposal and waste management plays an important role in reducing the impact of solid waste and heavy metal on the environment and the people who live in and around landfill areas. People who reside nearby landfills are exposed to a number of harmful chemicals that are created during the landfilling process^{4,5}. Litter, dust, excessive rodents, unforeseen landfill fires are some additional contaminants linked to the deposition of trash on landfills. The type and quantity of garbage placed, the age of the landfill, and the climatic conditions of the landfill sites are all factors that affect the by-product or emissions from landfills. Heavy metals, particulate debris, and a number of gaseous contaminants (including dioxins and polycyclic aromatic hydrocarbons) are regularly formed in the landfill as a result of intricate chemical and microbiological processes^{6,7}.

Urbanization is a key issue in developing nations like Indonesia, which has faced various environmental challenges, including the percentage of solid waste, due to social issues including socioeconomic, economic, and population inequality. Traditional solid waste management methods used in Indonesian towns have not been designed to advance environmental protection, public health, or renewable energy sources^{8,9}. These methods of managing solid waste are now widely recognised as the primary contributors to contamination of surface and groundwater as well as a degradation in the cleanliness of cities and towns^{10,11}. Makassar is one of the places, and it deals with concerns including how to manage heavy metal problems in final disposal sites and poorly managed solid waste operations (FDS). Makassar City has been undergoing a significantly rapid development which unfortunately contributes to the increasing amount of waste produced. Waste is an undeniably serious issue which grows with the population in Makassar City.

The Final Disposal Site (FDS) is a vital necessity to create a convenient environment for the community in big cities. If waste from all areas of Big City is not transferred to the FDS even for one day, the city will be full of waste, and it annoys the convenience of the community. It is an undeniable truth that the current conditions of FDS in some big cities have not met the standards of environmental safety. In fact, some FDSs are shut down because they are located too close to people residence and the resulting air pollution from waste deterioration has aggravated the community¹². Poorly managed FDS is critically harmful to human health and potentially damaging the environment. The human activities have caused serious damage to the environment, and significantly increased biological toxicity of heavy metals in soils. Some heavy metals in excessive concentration may harm the organisms around landfill. The heavy metal contamination also damages soils and animals^{13,14}.

Soils may be contaminated with heavy metals that are typically without colour or odour, thus are difficult to be controlled. Heavy metal contamination may not instantly be damaging to environment; however, it has been reported^{15,16} that when the concentration of heavy metals has exceeded the tolerance level of the environment, they will be activated and trigger ecological damage. Previous researchers have investigated

heavy metal concentration in the soils landfill in Slovakia¹⁷. Similarly, heavy metals in landfill may be removed by modified membrane¹⁸.

EXPERIMENTAL

Research area. The administrative area of the research encompassed the regional Bangkala Sub-district at Manggala District, Makassar City, South Sulawesi Province (Fig. 1).




Fig. 1. Location map of Antang landfill, Makassar city, South Sulawesi, Indonesia

Sampling point. The samples used in this study were the topsoil collected from the area of Antang Final Disposal Site, Makassar City, South Sulawesi. There were four sampling areas: two areas (A and C) aligned with the groundwater flow, and areas B and D were vertical to the groundwater flow. All points were measured vertically at the depth of 0–10, 10–20, and 20–30 cm depth.

RESULTS AND DISCUSSION

The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area A at the depth of 0–10, 10–20, and 20–30 are presented in Fig. 2.




Fig. 2. Results of analysis of heavy metals concentrations of area A

Area A shows that the concentration of Cu at the depth of 0–10 cm is 138 ppm, then increases to 168 and 177 ppm at the depth of 10–20 and 20–30 cm, respectively. The concentration of Pb at 0–10 cm depth is around 104 ppm, then increases to 11 ppm at 10–20 cm depth, and to 105 ppm at 20–30 cm depth. Meanwhile, Zn concentration at 0–10 cm was 128 ppm which decreases to 126 and 118 ppm at the depth of 10–20 and 20–30 cm, respectively. The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area B at the depth of 0–10, 10–20, and 20–30 cm are presented in Fig. 3.




Fig. 3. Results of analysis of heavy metals concentrations of area B

Area B shows that the concentration of Cu at the depth of 0–10 cm is 67 ppm, then increases to 72 ppm (10–20 cm) and decreases to 64 ppm (20–30 cm). Pb concentration at 0–10 cm depth is around 52 ppm, then decreases to 43 ppm (10–20 cm) and increases to 63 ppm (20–30 cm). Meanwhile, Zn concentration at 0–10 cm was 61 ppm, then increases to 64 ppm (10–20 cm), and decreases back to 61 ppm (20–30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area C at the depth of 0–10, 10–20, and 20–30 cm are presented in Fig. 4.




Fig. 4. Results of analysis of heavy metals concentrations of area C

Area C shows that the concentration of Cu at the depth of 0–10 cm is 65 ppm, then increases to 84 ppm (10–20 cm) and decreases to 43 ppm (20–30 cm). Pb concentration at 0–10 cm depth is around 61 ppm, then increases to 91 ppm (10–20 cm) and decreases to 31 ppm (20–30 cm). Meanwhile, Zn concentration at 0–10 cm was 50 ppm, then increases to 58 ppm (10–20 cm), and decreases to 4 ppm (20–30 cm). The results of analysis of heavy metals concentrations of Cu, Pb, and Zn at area D at the depth of 0–10, 10–20, and 20–30 cm are presented in Fig. 5.




Fig. 5. Results of analysis of heavy metals concentrations of area D

Area D shows that the concentration of Cu at the depth of 0–10 cm is 158 ppm, then increases to 161 ppm (10–20 cm) and decreases to 149 ppm (20–30 cm). Pb concentration at 0–10 cm depth is around 73 ppm, then decreases to 61 ppm (10–20 cm) and back up again to 73 ppm (20–30 cm). Meanwhile, Zn concentrations at 0–10, 10–20, and 20–30 cm are around 132, 142 and 142 ppm, respectively.

Heavy metals, such as Cu, Pb, and Zn in Antang Final Disposal Site, Makassar City, South Sulawesi have vertically distribution across the topsoil with a variety of concentrations at each depth which has exceeded the limit of contaminant in soil environment, which is acidic. The recommended limit of heavy metal concentrations by the Government of the Republic of Indonesia and Food and Drug Authority is 100 ppm of Cu, 100 ppm of Pb, and 70 ppm of Zn.

Heavy metal Cu is strongly associated with organic matters, iron and manganese oxidation, silica clay and a slight amount of mineral. Cu is particularly absorbed and is slightly mobile in the soils. Organic matter is the dominant factor that controls the

retention of Cu. Most Cu is soluble in the soil surface in a complex form of Cu^{2+} . It means that, if Pb heavy metal in the soils functions as Pb^{2+} oxidiser. If pH increases, the Pb^{2+} ion becomes less soluble oxidiser. A higher pH will form a complex form which allows the organic matters settle as carbonate, hydroxide, and phosphate. Lead is one of heavy metals that poses significant harms to living organisms because it is carcinogenic. The recent study found varying levels of Zn and the average pH was indicative of acidity. The oxidation of Zn in the soils shows Zn^{2+} under oxidised acid conditions. Zn^{2+} is one of the heavy metals that is the most soluble and mobile in the soil, hydrocarbon oxide, and organic matters. Zn solubility in the soils is highly potential at acidic pH (Ref. 19).

The concentrations of heavy metals in this study which also have surpassed the limit are tolerance. High levels of Cu in Final Disposal Site reflect the high quantity of waste dumped to the FDS (Ref. 20). The enrichment of heavy metal in the Antang landfill has been highly accumulated because of the anthropogenic activities or human activities, leachate, and soil characteristics and supported by a very acidic pH, the mobility of acid in heavy metals in the soils is relatively high²¹. Heavy metals will be carried by leachate and deposited in the soils, and as a result, heavy metals like Cu, Pb, and Zn will remain deposited in the soils.

Relatively acid soils have an increasing solubility and mobility of nutrients, thus increasing the concentration of heavy metals in the soils²². If the soil solution is too acidic, plants can not utilise and other nutrients they need to grow. Meanwhile, acidic soils (low pH) are dominated by ion Al, Fe, and Mn which will bind the nutrients vital for the plants, especially P, K S, and Mg. As a result, plants can not absorb their food well despite the richness of nutrients in the soil. In addition, acidic soils contain high level of micro nutrients, which are very dangerous to people's health²³, particularly those living around the landfill because it causes cancer and lung damage in addition to IQ drop among children, and potential death to human in general.

CONCLUSIONS

The increased manufacturing and widespread availability of different consumer items have an impact on the continuous rise in garbage output. Municipal waste management and the effects of heavy metals on the environment and community health in Indonesia and developing countries can be significantly improved by even just coordinated work and integrated activities of experts, researchers, government and municipal officials, as well as citizens and other stakeholders. The expansion of the municipal trash collection system, the construction of regional centres for recycling and reuse activities, and finding practical and affordable alternatives to locate consolidated sanitary landfills for the disposal of municipal solid waste are the main strategic priorities. In addition to keeping people and the environment safe, this can offer financial payback and the chance to partially recoup maintenance costs for waste management systems. As it gathers useful informational bases required for the development of environmental

protection strategies, city development strategies, and information for spatial planning in the Indonesia region, this research can be applied by the city's official agencies for environmental management and protection as well as spatial planning.

REFERENCES

1. N. J. MOKOELE: Complexities of Contemporary Urban Planning in Local Government in the City of Polokwane, Limpopo Province. *Jamba J Disaster Risk Stud*, **15** (1), a132 (2023). <https://doi.org/10.4102/jamba.v15i1.1326>.
2. P. O. NJOKU, J. N. EDOKPAYI, J. O. ODIYO: Health and Environmental Risks of Residents Living Close to a Landfill: a Case Study of Thohoyandou Landfill, Limpopo Province, South Africa. *International Journal of Environmental Research and Public Health (IJERPH)*, **16**, 1 (2019). <https://doi.org/10.3390/ijerph16122125>.
3. Y. H. WELDEYOHANIS, A. B. ANESEYEE, T. H. SODANGO: Evaluation of Current Solid Waste Disposal Site Based on Socio-economic and Geospatial Data: a Case Study of Wolkite Town, Ethiopia. *Geo J*, **87**, 585 (2022). <https://doi.org/10.1007/s10708-020-10268-3>.
4. S. SASAKI, K. WATANABE, N. WIDYANINGSIH, T. ARAKI: Collecting and Dealing of Recyclables in a Final Disposal Site and Surrounding Slum Residence: the Case of Bantar Gebang, Indonesia. *J Mater Cycles Waste Manag*, **21**, (2019). <https://doi.org/10.1007/s10163-018-0798-2>.
5. M. PAN, H. LI, J. WU: Study on Catalysed Treatment of Coking Wastewater by TiO_2 . *Oxid Commun*, **39** (4), 3457 (2016).
6. I. S. ABDELLI, M. ASNOUNE, Z. ARAB, F. ABDELMALEK, A. ADDOU: Management of Household Waste in Sanitary Landfill of Mostaganem District (Western Algeria). *J Mater Cycles Waste Manag*, **19**, 265 (2017). <https://doi.org/10.1007/s10163-015-0415-6>.
7. W. XIAOYUN, Z. YOUFEI, L. SHENGTAO, L. KESI: Research Status of Methane Emission in Landfill Sites. *Oxid Commun*, **39** (4), 3090 (2016).
8. M. M. HOQUE, M. T. U. RAHMAN: Landfill Area Estimation Based on Solid Waste Collection Prediction Using NN Model and Final Waste Disposal Options. *J Clean Prod*, **256**, 1 (2020). <https://doi.org/10.1016/j.jclepro.2020.120387>.
9. T. MESTER, G. SZABÓ, Z. SAJTOS, E. BARANYAI, G. SZABÓ, D. BALLA: Environmental Hazards of a Recultivated Liquid Waste Disposal Site on Soil and Groundwater. *Water*, **14**, 1 (2022). <https://doi.org/10.3390/w14020226>.
10. L. MORAIS, V. NASCIMENTO, S. SIMÕES, J. OMETTO: Regional Distance Routes Estimation for Municipal Solid Waste Disposal, Case Study São Paulo State, Brazil. *Energies*, **14**, 1 (2021). <https://doi.org/10.3390/en14133964>.
11. R. POBLETE, N. K. SALIHOGLU, G. SALIHOGLU: Landfill Leachate Treatment Using Advanced Oxidation Processes. *Oxid Commun*, **39** (1), 493 (2016).
12. B. MANIMEKALAI, P. VIJAYALAKSHMI: Analysis of Leachate Contamination Potential of a Municipal Landfill Using Leachate Pollution Index. *IOSR J Environ Sci Toxicol Food Technol*, **2** (1), 16 (2014). <https://doi.org/10.9790/2402-0211639>.
13. G. U. CHIBUIKE, S. C. OBIORA: Heavy Metal Polluted Soil: Effect on Plants and Bioremediation Methods. *Hindawi Pub Corp; Applied and Environmental Soil Science*, 2014, 1-13. <https://doi.org/10.1155/2014/752708>.
14. V. TSHELEZA, S. NDHLEVE, H. M. KABITI, C. M. MUSAMPA, M. D. V. NAKIN: Vulnerability of Growing Cities to Solid Waste-related Environmental Hazards: the Case of Mthatha, South Africa. *Jamba J Disaster Risk Stud*, **11** (1), a632 (2019). <https://doi.org/10.4102/jamba.v11i1.632>.
15. R. NAWAZ, M. ASLAM, I. NASIM, M. A. IRSHAD, S. AHMAD, M. LATIF, F. HUSSAIN: Air Pollution Tolerance Index and Heavy Metals Accumulation of Tree Species for Sustainable Environmental Management in Megacity of Lahore. *Air*, **1** (1), 55 (2023). <https://doi.org/10.3390/air1010004>.

16. P. LIU, C. WANG, M. HAO, Z. XU, Y. LI: Treatment of Oily Wastewater by the Fenton Reagent and Wet Hydrogen Peroxide Oxidation. *Oxid Commun*, **39**, 2666 (2016).
17. P. ANDRAS, L. E. TURISOVA, S. KHARBISH, J. KRNAC, L. CMIELOVA: Environmental Risk due to Heavy Metal Contamination Caused by Old Copper Mining Activity at Lubietova Deposit, Slovakia. *Energy Environ Res*, **3** (2), 1 (2016). <https://doi.org/10.5539/eer.v3n2p182>.
18. A. R. ESFAHANI, L. ZHAI, A. H. M. A. SADMANI: Removing Heavy Metals from Landfill Leachate Using Electrospun Polyelectrolyte Fiber Mat-laminated Ultrafiltration Membrane. *J Environ Chem Eng*, **9** (4), 105355 (2021). <https://doi.org/10.1016/j.jece.2021.105355>.
19. F. F. B. FABORODE: Gender Assessment of Poultry Waste Utilization among Small-scale Poultry Farmers in Osun State, Nigeria: Exploring the Untapped Potentials. *Afr J Food, Agric Nutr Develop*, **22** (4), 20259 (2022). <https://doi.org/10.18697/ajfand.109.19780>.
20. M. B. SULAIMAN, A. S. MAIGARI, S. DANLADI: Impact of Municipal Solid Waste Dumps on Surrounding Soil and Groundwater in Gombe, Nigeria. *Int J Sci Environ* **5** (5), 3059 (2016).
21. L. BITYUKOVA, A. SHOGENOVA, M. BIRKE: Urban Geochemistry: a Study of Element Distribution in the Soil of Tallin (Estonia). *Environ Geochem Health*, **22**, 173 (2000). <https://doi.org/10.1023/A:1006754326260>.
22. I. SAZYKIN, L. KHMELEVTSOVA, T. AZHOGINA, M. SAZYKINA: Heavy Metals Influence on the Bacterial Community of Soils: a Review. *Agriculture*, **13**, 3, 653 (2023). <https://doi.org/10.3390/agriculture13030653>.
23. K. D. SHARMA, S. JAIN: Municipal solid Waste Generation, Composition, and Management: the Global Scenario. *Soc Responsib J*, **16** (6), 917 (2020). 10.1108/SRJ-06-2019-0210.

Received 29 April 2023

Revised 18 May 2023