

RIWAYAT KORESPONDENSI


1. Account created in GMC (15 November 2022)
2. Submitted manuscript (16 November 2022)
3. Review Round 1 from reviewer, minor revision (24 November 2022)
4. Revised from author (27 November 2022)
5. Accepted (29 November 2022)
6. Galley Prof from editor (10 Desember 2022)
7. Online publishing in volume 130 number 4 2022 (17 December 2022)

KORESPONDENSI


Title : Heavy Metal Levels of Lead and Cadmium in Baronang Fish from South Sulawesi, Indonesia

Link : [h=p://saber.ucv.ve/ojs/index.php/rev_gmc/arGcle/view/25319](http://saber.ucv.ve/ojs/index.php/rev_gmc/arGcle/view/25319)

ACCOUNT CREATED (15 November 2022)



SUBMISSION (16 November 2022)



REVIEW (24 November 2022)

99+

Mail

Chat

Meet

Compose

Inbox1,023

Starred

Snoozed

Sent

Drafts26

More

Labels+

Heavy Metal Levels

Active

8 of 8

Anita Stern-astern88@gmail.com>

to muhammad.ikhtiar, ENRIQUE, me

Thu, Nov 24, 2022, 4:22 AM

Star

Reply

More

Distinguished Drs. Muhammad Ikhtiar and Haeril Amir:

We hereby inform you about the status of your manuscript entitled: **Study of Heavy Metal Levels of Lead (Pb) and Cadmium (Cd) Inside Baronang Fish from South Sulawesi, Indonesia**

We have made a pre-review of the manuscript and it is observed that it requires profound corrections before it is sent to referees revision. In the attachment we are making important comments, suggestions, and recommendations that should be considered and that could significantly improve the manuscript.

We are sending you in attachment the aforementioned pre-revised manuscript for the purposes that, if you wish, correct it according to the observations and suggestions, and forward it for reconsideration for publication in the Gaceta Medica de Caracas. Attached are also the Authors Instructions of our Journal.

We encourage you to take into account the comments and suggestions of the referees.

We appreciate sending the individualized response separately; and indicate in the text of the revised manuscript version that we are sending you in attachment, in red letters, the corrections or modifications made.

Best regards,

Dr. Enrique López-Loyo
Editor-in-Chief


Dr. Anita Stern Israel
Senior Editor

Gaceta Médica de Caracas


2 Attachments • Scanned by Gmail

The screenshot shows an email client interface. On the left is a sidebar with navigation options: Compose, Mail, Chat, and Meet. The main area displays an email from Haeril Amir (haeril.amir@uml.ac.id) to Anita. The email body includes a PDF attachment titled "(ok) Study of Heavy Metal Levels of Lead (Pb) and Cadmium (Cd)" and a text message that says: "Dr. Haeril Amir. Again, Before considering again the manuscript revised, the list of references **MUST be carefully corrected** according to the style of the *Gaceta Médica de Caracas*. As follows: **References** Will be submitted in accordance with ICMJE Recommendations. You are encouraged to follow the recommendations in *Samples of Formatted References for Authors of Journal Articles* [15]. Indicate them with Arabic numbers in parentheses in a correlative way and in the order in which they appear for the first time in the text, tables, and figure legends. In the citations of publications with more than six authors, only the first 6 authors of the work should be included, followed by *et al.*"

ACCEPTED (29 November 2022)



GALLEY PROOF (10 Desember 2022)



LAMPIRAN FILE REVIEW

COMMENT:

1. The title is suggested to be changed
 2. It was corrected in the Abstract and added in Spanish
 3. Thought out the text there were made several corrections. There are many sentences in Indonesian that were translated. Check that the manuscript does not present sentences in the Indonesian language
 4. The units of the heavy metals should be carefully corrected.
 5. The references should be carefully corrected as indicated in the author's instructions of the Gaceta Medica de Caracas
 6. The references should be translated into the English language, not into Indonesian
- These are all form corrections, the referee requires this correction to correct the scientific content of the manuscript

**Heavy Metal Levels of Lead and Cadmium in Baronang Fish from South Sulawesi,
Indonesia**

**Niveles de metales pesados de plomo y cadmio en el pez Baronang del sur de Sulawesi,
Indonesia**

*Muhammad Ikhtiar¹, Nur Ulmy Mahmud ¹, Rezky Aulia Yusuf ¹, Haeril Amir²

¹Public Health Department, Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia

²Nursing Department, Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia

*Corresponding author: Muhammad Ikhtiar

Email: muhammad.ikhtiar@umi.ac.id

Adress: Jalan. Urip Sumeharjo No.05, Kel. Panaikang, Kec. Panakkukang, Makassar City, South Sulawesi, Indonesia, 90231, Telp: +6285343723079

RECIBIDO: 16 DE NOVIEMBRE 2022

ACEPTADO: 24 DE NOVIEMBRE 2022

SUMMARY

Objective: The purpose of this study was to determine the content of the heavy metal lead (Pb) and cadmium (Cd) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

Methods: This was an observational descriptive approach research. The accidental sampling technique was used at two locations, Kayu Bangkoa pier and Kera-Kera pier. Baronang fish samples were taken using fishing rods or nets and each bait was examined and analyzed in the laboratory.

Result: The results of an examination of Baronang fish samples from each sample at three different points around the Bangkoa Wood Pier, indicated a lead (Pb) parameter for sample 1 of 0.14 µg/g, for sample 2 <0.001 µg/g, while sample 3 was <0.001. For the parameter of cadmium (Cd) the sample 1 yield was 10.087 mg/g, sample 2 was 0.085 mg/g and sample 3 0.086 mg/g. The results of the examination of the Baronang fish sample indicate that lead (Pb) and cadmium (Cd) parameters were below standard SNI 7387: 2009, namely lead (Pb) 0.3 and cadmium (Cd) 0.1. While the results of the examination of Baronang fish samples from each sample at three different points around the Kera-Kera Pier indicated a lead (Pb) parameters for sample 1 which was <0.001 µg/g, sample 2 <0.001 µg/g and sample 3 <0.001.

Conclusion: The Baronang fish samples at three points of Bangkoa Wood Pier presented Pb parameter below the SNI 7387: 2009 standard, namely lead (Pb) 0.3, while the results of the cadmium (Cd) examination of the three samples all had passed the maximum limit of the SNI standard 7387: 2009, that is 0.1.

Keywords: Baronang fish, cadmium (Cd), environmental, lead (Pb)

RESUMEN

Objetivo: El propósito de este estudio fue determinar el contenido del metal pesado plomo (Pb) y cadmio (Cd) en el pez Baronang en el muelle de Kayu Bangkoa y el muelle de Kera-Kera en la ciudad de Makassar, South Sulawesi, Indonesia en 2020.

Métodos: Se utilizó un enfoque observacional descriptivo. La técnica de muestreo accidental se utilizó en dos lugares, el muelle de Kayu Bangkoa y el muelle de Kera-Kera. Se tomaron muestras de peces Baronang utilizando cañas de pescar o redes y luego cada cebo se examinó y analizó en el laboratorio.

Resultado: Los resultados del examen de las muestras de pescado Baronang en tres puntos diferentes alrededor del muelle de madera de Bangkoa indicaron parámetros de plomo (Pb) de la muestra 1 de 0,14 µg/g, la muestra 2 <0,001 µg/g mientras que la muestra 3 fue <0.001. Para el parámetro de cadmio (Cd) la muestra de 1 fue de 0.087 mg/g, la muestra 2 de 0.085 mg/g y la muestra 3 de 0.086 mg/g. Lo que indican parámetros de plomo (Pb) y cadmio (Cd) para el pescado Baronang con valores por debajo del estándar SNI 7387: 2009, a saber, plomo (Pb) 0,3 y cadmio (Cd) 0,1. Mientras que los resultados del examen de muestras de pescado Baronang de tres puntos diferentes alrededor del muelle de Kera-Kera indicaron para parámetros de plomo (Pb) que la muestra 1 era <0,001 µg/g, la muestra 2 <0,001 µg/g y la muestra 3 <0.001.

Conclusión: Las muestras de pescado Baronang tomadas en tres puntos indicaron parámetros de plomo (Pb) por debajo del estándar SNI 7387: 2009, mientras que los de cadmio (Cd) superaron el límite máximo de la norma SNI 7387: 2009.

Palabras clave: pez Baronang, cadmio (Cd), ambiental, plomo (Pb)

Introduction

Heavy metal contamination of aquatic ecosystems has been a problem in environmental health for decades. Heavy metals are highly hazardous contaminants due to their persistence,

toxicity, and bioaccumulation in various segments of the environment, such as water, sediments, air, and biota (1, 2). Heavy metal concentrations lower than the required amounts may cause deficiency effects, while high concentrations of these metals lead to health problems (3). Toxic metals including lead (Pb) and cadmium (Cd) are hazardous because they can cause severe health problems in minute quantities (4, 5).

One of the most vulnerable environmental aspects of heavy metal contamination is water (6). Coastal ecosystems are very sensitive to heavy metal contaminants; sediments and marine organisms can accumulate pollutants, and fish can get enriched with metals that are subsequently transferred to man through human consumption (7).

Recently, heavy metal pollution in an aquatic ecosystem is becoming a critical issue. Based on the results of provenance analyses of heavy metals suggest that geogenic sources were the main source of water contamination (8, 9). Nonetheless, the industry activity, and human activities have increased concentrations of heavy metals above normal levels in the environment contributing to serious and widespread environmental issues leading to chronic toxicity (10-13).

In line with this, in Indonesia based on research conducted by the Oceanography Center of LIPI (The Indonesian Institute of Science), reported that high levels of metals are generally detected at the stations located close to the mainland, indicating that heavy metals are enrichment from terrestrial anthropogenic activities (14).

In a study about the incidence of Pb, and Cd in seafood from Africans on the Indian and the Red Sea coasts, it was found that some seafood from African Indian and the Red Sea coasts such as fishes have presented Cd and Pb concentrations higher than permitted limit by FAOUN/EU regions, indicating a possible threat to public health (15).

Based on a study on the bioaccumulation of Cadmium (Cd) and Chromium (Cr) contained in water and fish carried out in the Tallo Makassar river using quantitative and observational methods, it was found that the cadmium content in 3 research points was $<0.003 \text{ mg / L}$ (13). This cadmium content fulfills the requirements based on the determination of heavy metal content which refers to the government regulation PP number 82 of 2001 concerning Water Quality Management and Water Pollution Control with the designation of water with class II criteria, where the maximum cadmium (Cd) content is 0.01 mg/L . The content of heavy metal cadmium in this fish is not only caused by industrial activities but also due to the densely populated environmental conditions and the behavior of the people who always dispose of domestic waste in the area around the river (13).

Based on research conducted in the waters of Benoa Bay, Bali, it was found that the content of heavy metals Pb and Cd in the water had exceeded the quality standards for dissolved metals in water for marine biota as stated in the Bali Governor Regulation No. 16 of 2016. For the baronang fish itself, the content of heavy metal Cd has exceeded the maximum limit of metal contamination, which is 0.1 mg / Kg , while the Pb content in some samples has exceeded the maximum limit of heavy metal contamination of SNI 7387: 2009 (16). Cd bpom 0.10, pb 0.20 NAV.

Fish are a significant bioindicator of heavy metal contamination in the aquatic ecosystem, as they are at the top of the food web and can accumulate considerable amounts of heavy metal in their tissues (13, 17-19). Generally, higher concentrations of metals were found in the liver and gills than in muscles (20). A study using a dry ashing-acid digestion method in fish collected from

Sri Murni Lake and Kepong Metropolitan Lake around Kepong district, Kuala Lumpur, found that the highest concentration of Pb was detected in the gills ($0.151 \pm 0.12\text{mg/g}$) while Cd, was highly accumulated in the bones ($1.750 \pm 3.43 \text{ mg/g}$) (21).

Based on the studies above, water and the ecosystem in it are very susceptible to being contaminated by heavy metals which have a broad impact on long-term public health conditions. Therefore, Periodic research and monitoring of water and fish quality should be given attention so that the quality of the lake ecosystem, especially in relation to heavy metal elements, is under control which makes activities involving water contact safe (10). Therefore, the purpose of this study was to determine the content of the heavy metal lead (Pb) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

Methods

The method used in this study was sampling using the incidental sampling method, at two locations of the Bangkoa Wood Pier and the Kera-Kera Pier. Baronang fish samples are taken using fishing rods or nets and each bait is then examined and analyzed in the laboratory. Primary data collection is done utilizing field observations or reviewing the research location by taking samples of Baronang fish then examined and analyzed in the laboratory.

Baronang fish examination samples were taken at six points at two locations of the bangkoa wood pier and the Kera-Kera pier. Then measured using Automic Absorption Spectrophotometer (AAS) equipment by laboratory staff, namely the South Sulawesi Provincial Health Office Laboratory.

Results

This research was conducted in two different places with sampling on 20 November 2020 at the Kayu Bangkoa pier in the Bulogading Village Ujung Pandang sub-district and on March 20, 2016, at Kera-Kera Pier in Tamalanrea Indah Village, Tamalanrea District, then the clean water samples were taken to the Great Hall Makassar Health Laboratory, South Sulawesi Province to find out the results of examining clean water samples.

Based on the results of the examination Great Hall Makassar Health Laboratory, South Sulawesi Province, with samples of Baronang Fish, to determine water quality based on SNI 7387: 2009. The Maximum Limit of Heavy Metal Contamination in Food is as follows: that categorize “Qualified (Q) = not reach the maximum limit)” and “Unqualified Qualified (UQ) = reach the maximum limit);

Table 1. Results of Sampling of Baronang Fish at the Bangko Wood Pier Bulogading Village Kec. Ujung Pandang

No.		Sample	Checkup result	Level
-----	--	--------	----------------	-------

	Sample Code		Lead (Pb) (µg g)	Cadmium (Cd) (Ug / g)	Pb	Cd
1	Sample 1	Baronang fish	0.14	0.087	Q	Q
2	Sample 2	Baronang fish	<0.01	0.085	Q	Q
3	Sample 3	Baronang fish	<0.01	0.086	Q	Q

Source: Primary Data, 2020

*Q: Qualified

Table 1 shows that the samples of Baronang Fish originating from the Bangkoa Wood Pier were taken at the first point for lead (Pb), namely with sample code 1 0.14 (µg / g), sample 2 <0.01 (µg / g), and sample 3 <0.01 (µg/g). As for the results of the cadmium (Cd) sample, the sample code is 1 0.087 (µg/g), sample 2 0.085 (µg/g), and sample 3 0.086 (µg/g). From the results of the sample examination on Baronang Fish, all of them are still below the SNI 7387: 2009 standard, namely 0.03 mg/kg for lead (Pb), and cadmium also meets the requirements, namely 0.1 mg/kg.

Table 2. Results of Sampling of Baronang Fish at Kera-Kera Pier Kelurahan Tamalanrea Indah Kec. Tamalanrea

No.	Sample Code	Sample	Checkup result		Level	
			Lead (Pb) (µg / g)	Cadmium (Cd) (µg/g)	Pb	Cd
1	Until 1	Baronang fish	<0.01	0.170	Q	UQ
2	Sample 2	Baronang fish	<0.01	0.301	Q	UQ
3	Sample 3	Baronang fish	<0.01	0.386	Q	UQ

Source: Primary Data, 2020

*UQ: Unqualified

Table 2 shows that the samples of Baronang Fish originating from the Kera-Kera Pier were taken at the first point for lead (Pb), namely with sample code 1 <0.01 (µg / g), sample 2 <0.01 (µg / g), and sample 3 <0.01 (µg / g) is still below the standard of SNI 7387: 2009, namely 0.01 mg/kg. As for the results of the cadmium (Cd) sample with a sample code of 1 0.087 (µg / g), sample 2 0.085 (µg / g), and sample 3 0.086 (µg / g). From the results of the sample examination on Baronang Fish, all of them did not meet the requirements of the SNI 7387: 2009 standard, namely 0.01 mg/kg for cadmium (Cd) while the lead (Pb) met the requirements, namely 0.3 mg/kg.

Discussions

Fish has many health benefits due to its nutritional content. However, heavy metals contamination such as lead (Pb) and cadmium (Cd) in aquatic ecosystems, especially in fish, which is the daily consumption of the community, is a major threat to the quality of public health due to its negative impact due to long-term consumption (22).

The metals that pose the highest risks to human health are **one of them is** cadmium and lead which cause important complications in the nervous system, kidneys, bones, lungs, and cardiovascular system due to their toxicity and possible carcinogenic effect (23). Bahkan kontaminasi heavy metals have been linked to the development of mental retardation and even death in instances of very high exposure in the human body (20). Some of them contribute to the possible adverse effects, particularly in the fetus and young children (9).

Cd toxicity by consumption of contaminated fish can cause the following signs and symptoms: increased salivation, choking or vomiting, abdominal pain, vertigo, loss of consciousness, painful spasms of the anal sphincter, and impairment of renal function for severe toxicity (24, 25).

Various systems are affected due to Pb toxic exposure in fish, such as disruption of neurotransmitter function occurs due to Pb accumulation causing neurotoxicity and interfering with the immune system response (26). In humans, Pb can cause defined specific changes in gene and protein expression in response to lead challenges and determine the injurious effects of exposure to lead on a cellular level (27).

Acute Pb toxicity after ingestion of contaminated seafood usually occurs in the brain and kidney, and its absorption in the gastrointestinal tract is influenced by nutritional calcium and iron status and age (children adsorb more, and consequently, are more vulnerable than adults (28).

Exposure pathways (dietary and waterborne), environmental factors (salt-water or freshwater), and Pb binding capacity with protein decide the accumulation pattern of heavy metals exposure. The main cause of heavy metal contamination in the African countries bordering the Indian Ocean and the Red Sea port activity, tourism, and petroleum activities (15). In Pakistan, particularly in Swat River, heavy metal contamination might be due to geogenic (mafic and ultramafic rocks) and anthropogenic activities including industrialization, municipal water, pesticides application, wastewater from farm fields sewage sludge, automobile workshops, mining effluents, hotels effluents and urban districts including (Charsadda Malakand Agency and Swat) located near Swat River (29).

Similar to other countries, in Indonesia, especially in Makassar, heavy metal contamination of cadmium (Cd) and Lead (Pb) in waters and fish occurs due to various activities such as ports, tourism, and fisheries activities (30). Besides that, the density of human settlements which causes a large amount of waste and industrial wastewater is also a major factor in pollution in water areas (30).

Even though the concentration of heavy metals in the waters, especially in fish, is still below the threshold, the control and monitoring of heavy metals in fish are most important to minimize associated fish poisoning (15). The monitoring of THM concentrations in river water,

sediments, and fish muscles is, therefore, essential to make sure acquiescence with food security guidelines and resulting end-user safety (29).

Levels of lead (Pb) and Cadmium (Cd) Heavy Metals in Baronang Fish at Bangkoa Wood Pier

Water conditions contaminated by various metals will significantly affect the aquatic ecosystem in both inland and marine waters. Lead (Pb) is a metal that is widely used by humans, so this metal also causes contamination impacts on the environment. Pb metal pollution can also be caused by the entry of industrial waste containing Pb metal into water bodies. Furthermore, with the presence of the biomagnification process in the waters, the concentration of Pb metal will continue to increase. Heavy metals enter the fish's body tissues through several ways, namely: the respiratory tract (gills), the digestive tract (intestines, liver, kidneys) through the process of the food chain, and penetration through the skin (flesh muscle) (31).

The results of the examination at the Baronang Fish Testing Laboratory at the Makassar Health Laboratory Center, South Sulawesi Province that the samples of baronang fish originating from two different pier areas, in Bulogading and Tamalanrea Indah sub-districts were taken at 6 different points every 3 points per pier there were several comparisons of the number of levels obtained in sampling at two different piers.

The results of the examination of baronang fish samples at 3 points taken around the Bangkoa Wood Pier with lead (Pb) parameters stated that sample 1 with a result of $0.14\mu\text{g/g}$ at sample 1 was found, for sample 2 with a result $<0.001\text{ }\mu\text{g/g}$ while sample 3 resulted <0.001 and for the parameter of cadmium (Cd) with a sample result of $1\text{ }0.087\text{ mg/g}$, for sample 2 results of 0.085 mg/g while for sample 3 with a result of 0.086 mg/g from the results of an examination of the Baronang Fish sample at three points with the parameters of lead (Pb) and cadmium (Cd) are still below the standard of SNI 7389: 2009. Meanwhile, the results of research based on Budiman's research at the Wangisgara station in the upstream Citarum river showed the cadmium (Cd) content in Betok fish was 11 ppb with a weight of 22.9 gr, for 92 ppb catfish with a weight of 36.9 gr and a high level of cadmium (Cd) in Betok fish. Cadmium (Cd) in Tilapia 105 ppb with a weight of 20.2 grams. The three types of fish have exceeded the quality standards set by SNI 7387: 2009 concerning the limit of metal contamination in fish meat and its processed products (32).

At point 2 there is the Losari beach platform around Losari Beach. There are houses, Stella Maris Hospital, Siloam Hospital, Arya Duta Makassar Hotel, Aerotel Smile Makassar, D'bugis Ocean, Anging Mammiri Hotel and community settlements that can produce liquid waste and solid which can cause heavy metal pollution. This also shows from the research that heavy metals originating from human activities can be industrial or household waste materials and are also caused by sludge from dirty drains, recycling, lead-containing manufactured products (gasoline, paint, printing ink, main water pipes, tin solder cans, and battery casings). The solubility of heavy metals in water bodies is controlled by the degree of acidity of the water, the type, and concentration of metals and the state of mineral components (33).

Whereas at point 3 it is near the container port where around the container the entry and exit of cargo ships around the area can provide pollution such as Oil Spills, Hotel Pantai Gapura, Hotel

Makassar Golden and Cafe Balliriate, and Zona Cafe. Borate metal was not identified because it is still influenced by sea characteristics such as tides, sea breezes, and saltwater infiltration.

Levels of Heavy Metal Lead (Pb) and Cadmium (Cd) in Baronang Fish at Kera-Kera Pier

Cadmium (Cd) enters the environment because of human activities. Cadmium (Cd) in water bodies can come from atmospheric deposits, dust, sewage processing water, and industrial wastewater. Cadmium enters the water more due to human activities such as industry where the waste products from the factory are discharged directly into the water which will accumulate at the bottom of the waters and form sediment. Cadmium (Cd) can also enter organisms that live in water where Cadmium (Cd) can enter via oral, inhalation, or dermal. Cd enters the body of an organism, for example, as fish.

From the results of the examination at the Baronang Fish Testing Laboratory at the Makassar Health Laboratory Center, South Sulawesi Province that the samples of Baronang Fish originating from two different pier areas, in Bulogading and Tamalanrea Indah Villages, were taken at 6 different points every 3 points per pier there were several comparisons of the number of levels obtained in sampling at two different piers.

The results of the examination of Baronang fish samples at 3 points taken around the Kera-Kera Pier with lead (Pb) parameters stated that sample 1 with a result <0.001 ug / g, for sample 2 with a result <0.001 ug / g while sampling 3 with a result < 0.001 and for the parameter of cadmium (Cd) with a sample yield of 1 0.170 mg / g, for sample 2 results of 0.301 mg / g while for sample 3 with a result of 0.386 mg / g from the results of an examination of the Baronang fish samples at three points with lead parameters (Pb) and cadmium (Cd) are still below the SNI 7387: 2009 standard, namely lead (Pb) 0.3, while the results of the examination of cadmium (Cd) of the three samples all have passed the maximum limit of the SNI 7387: 2009 0.1 standards.

This shows from the results of the research that the results of the examination of Baronang fish samples on the heavy metal Cadmium (Cd) were identified at point 1 in the vicinity of the Kera-Kera Pier where the area is adjacent to the pond, residential areas that have an impact on waste disposal and plastic waste disposal, and a stopover for passenger and sea boats where the boat uses fuel engines. Cadmium fuel contains up to 0.5 ppm. Based on Latif's research (2012), shows that the Tallo River has several settlements, the PLTU industry, the plywood factory industry, aquaculture, and agriculture (34).

Whereas at point 3 of the results of Setiawan's research demonstrated that the waters of the Tallo River are experiencing high pressure due to the presence of settlements, PLTU, aquaculture, and agriculture. And point 3 is also not far from the Pampang River, causing pollution and a lot of getting into the water (35). Mahmuda's study stated that the Pampang River is flanked by the Tallo River (70 km from the length of the main river) to the north there are several settlements, the alcohol industry, the chocolate processing industry, and hospitals. so that Cadmium (Cd) that enters the body of an organism such as fish will accumulate in the kidneys and liver because these two organs are very specific to fight toxins that enter the body (36).

Conclusion

The results of an examination of Baronang fish samples from each sample at three different points around the Bangkoa Wood Pier found that the lead (Pb) parameters stated that of the sample examination on Baronang Fish, all the samples were still below the SNI 7387: 2009 standard, namely 0.03 mg/kg for lead (Pb) and cadmium also meet the requirements, 0.1 mg/kg.

The results of the examination of Baronang fish samples from each sample at three different points around the Kera-Kera Pier with lead (Pb) parameters stated that originating from the Kera-Kera Pier were taken at the first point for lead (Pb), found all samples still below the standard of SNI 7387: 2009, namely 0.01 mg/kg. As for the results of the cadmium (Cd) from all the samples above the maximum limit of the requirements of the SNI 7387: 2009 standard.

Suggestions

Suggestions to the public and industry who live and have activities in the coastal areas and streams around the Bnagkoa Timber Pier to reduce the risk of pollution such as direct disposal of garbage and waste into rivers/sea.

Near the Kera-Kera Pier, it is hoped that the community will maintain the sanitation of the industrial environment around the Tallo River basin which uses heavy metal materials such as Cadmium (Cd) to carry out better treatment of their waste before being discharged into the environment to reduce the pollutant load that enters the waters of the Tallo River, Makassar City.

Acknowledgments

The author is grateful to the Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia, for numerous supports of this study.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Authors' Contributions

All contributors contributed significantly to this study and all authors agree with the content of the manuscript.

Conflict of Interest

The author(s) declared no potential conflicts of interest with respect to the research authorship, and/or publication of this article.

Availability of Data and Materials

All data generated or analyzed during this study are included in this published article.

Ethical Approval

This article received ethical clearance from *Ethic Commission, Universitas Muslim Indonesia*. Registration Number: UMI0320015671.

Significance for public health

Indonesia is one of the largest consumers of fish in the world. Heavy metal contamination has been a problem in environmental health that can affect human health. This study describes the content of the heavy metal lead (Pb) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

REFERENCES

**COMMENT: THE REFERENCES SHOULDS BE CORRECTED CAREFULLY
ACCORDING TO THE AUTHORS INSTRUCTION OF GACETA MEDICA DE
CARACAS**

1. Gashkina NA, Moiseenko TI, Kudryavtseva LP. Fish response of metal bioaccumulation to reduced toxic load on long-term contaminated lake Imandra. *Ecotoxicology and environmental safety*, 2020. **191**: p. 110205.
2. Xie, Q., et al., Assessment of long-term effects from cage culture practices on heavy metal accumulation in sediment and fish. *Ecotoxicol Environ Saf*, 2020. **194**: p. 110433.
3. Alves, R.N., et al., Oral bioaccessibility of toxic and essential elements in raw and cooked commercial seafood species available in European markets. *Food Chemistry*, 2018. **267**: p. 15-27.
4. Jadoon, S., et al., Potentially toxic elements in drinking water and associated health risk assessment in Abbottabad city, northern Pakistan. *Desalination and water treatment*, 2019. **2019**: p. 392-402.
5. Qian, Y., et al., Assessment of metal mobility in sediment, commercial fish accumulation and impact on human health risk in a large shallow plateau lake in southwest of China. *Ecotoxicol Environ Saf*, 2020. **194**: p. 110346.
6. Ciazela, J., M. Siepak, and P. Wojtowicz, Tracking heavy metal contamination in a complex river-oxbow lake system: Middle Odra Valley, Germany/Poland. *The Science of the total environment*, 2018. **616-617**: p. 996-1006.
7. Bat, L., A. ÖZtekİN, and F. Sahin, Heavy Metal Detection in *Scorpaena Porcus* Linnaeus, 1758 from Sinop Coast of the Black Sea and Potential Risks to Human Health. *Current Agriculture Research Journal*, 2018. **6**.
8. Muhammad, S. and K. Ahmad, Heavy metal contamination in water and fish of the Hunza River and its tributaries in Gilgit–Baltistan: Evaluation of potential risks and provenance. *Environmental Technology & Innovation*, 2020. **20**: p. 101159.
9. Basim, Y. and Z. Khoshnood, Target hazard quotient evaluation of cadmium and lead in fish from Caspian Sea. *Toxicol Ind Health*, 2016. **32**(2): p. 215-20.
10. Hashim, Z., et al., An Analysis of Heavy Metals in Lakes of Former Tin Mining Sites in the City of Ipoh, Perak, Malaysia. *International Journal of Academic Research in Business and Social Sciences*, 2018. **8**.
11. Kusin, F.M., et al., Distribution of heavy metals and metalloid in surface sediments of heavily-mined area for bauxite ore in Pengerang, Malaysia and associated risk assessment. *CATENA*, 2018. **165**: p. 454-464.
12. Sakai, N. and M. Yoneda, Potential Health Risk of Heavy Metals in Malaysia. 2018. p. 19-32.
13. Jais, N.J., et al., Bioakumulasi Logam Berat Kadmium (Cd) dan Kromium (Cr) yang Terdapat dalam Air dan Ikan di Sungai Tallo Makassar. *Window of Public Health Journal*, 2020. **1**(3): p. 261-274.

14. Harmesa, H., L. Lestari, and F. Budiyanoto, Distribusi Logam Berat Dalam Air Laut Dan Sedimen Di Perairan Cimanuk, Jawa Barat, Indonesia. *OLDI (Oseanologi dan Limnologi di Indonesia)*, 2020. **5**(1): p. 19-32.
15. Tamele I.J, Loureiro, PV. Lead, mercury and cadmium in fish and shellfish from the Indian Ocean and Red Sea (African Countries): Public health challenges. *Journal of Marine Science and Engineering*, 2020. **8**(5): p. 344.
16. Mardani, N.P.S., I.W. Restu, and A.H.W. Sari, Kandungan Logam Berat Timbal (Pb) dan Kadmium (Cd) Pada Badan Air dan Ikan di Perairan Teluk Benoa, Bali. *Current Trends in Aquatic Science*, 2018. **1**(1): p. 106-113.
17. Maury-Brachet, R., et al., Mercury contamination levels in the bioindicator piscivorous fish *Hoplias aimara* in French Guiana rivers: mapping for risk assessment. *Environmental Science and Pollution Research*, 2020. **27**(4): p. 3624-3636.
18. Zhong, W., et al., Health risk assessment of heavy metals in freshwater fish in the central and eastern North China. *Ecotoxicology and environmental safety*, 2018. **157**: p. 343-349.
19. Arkianti, N., N.K. Dewi, and N.K.T. Martuti, Kandungan logam berat timbal (Pb) pada ikan di Sungai Lamat Kabupaten Magelang. *Life Science*, 2019. **8**(1): p. 54-63.
20. Maurya, P.K. and D. Malik, Bioaccumulation of heavy metals in tissues of selected fish species from Ganga river, India, and risk assessment for human health. *Human and Ecological Risk Assessment: An International Journal*, 2019. **25**(4): p. 905-923.
21. Ishak, A.R., M.S.M. Zuhdi, and M.Y. Aziz, Determination of lead and cadmium in tilapia fish (*Oreochromis niloticus*) from selected areas in Kuala Lumpur. *The Egyptian Journal of Aquatic Research*, 2020. **46**(3): p. 221-225.
22. Olmedo, P., et al., Determination of toxic elements (mercury, cadmium, lead, tin and arsenic) in fish and shellfish samples. Risk assessment for the consumers. *Environment International*, 2013. **59**: p. 63-72.
23. Zuloaga, J., S. Gallego, and C. Ramírez, Contenido de Hg, Cd, Pb y As en especies de peces: revisión. *Vitae*, 2015. **22**: p. 148-159.
24. Johri, N., G. Jacquillet, and R. Unwin, Heavy metal poisoning: the effects of cadmium on the kidney. *Biometals*, 2010. **23**(5): p. 783-792.
25. Nordberg, G.F., Historical perspectives on cadmium toxicology. *Toxicology and applied pharmacology*, 2009. **238**(3): p. 192-200.
26. Ishaque, A., et al., Toxic effects of lead on fish and human. *Biological and Clinical Sciences Research Journal*, 2020. **2020**(1): p. e045-e045.
27. Gillis, B.S., Z. Arbieva, and I.M. Gavin, Analysis of lead toxicity in human cells. *BMC genomics*, 2012. **13**(1): p. 1-12.
28. Flegal, A.R., Lead in tropical marine systems: a review. *Science of the Total Environment*, 1986. **58**(1-2): p. 1-8.
29. Liu, M., et al., Contamination features, geo-accumulation, enrichments and human health risks of toxic heavy metal (oids) from fish consumption collected along Swat river, Pakistan. *Environmental Technology & Innovation*, 2020. **17**: p. 100554.
30. Noor, R.J., A. Kabangnga, and F. Fathuddin, Distribusi Spasial dan Faktor Kontaminasi Logam Berat di Pesisir Kota Makassar. *Jurnal Kelautan Tropis*, 2021. **24**(1): p. 93-101.

31. Ma'ruf, M., Analysis of Heavy Metal Concentration in Baronang Fish (*Siganus* sp) and Aquatic Environment for Management of Bontang Coastal Areas. 2017, Thesis (unpublished). Mulawarman University.
32. Budiman, B.T.P., Y. Dhahiyat, and I. Rustikawati, Bioakumulasi logam berat pb (timbal) dan cd (kadmium) pada daging ikan yang tertangkap di sungai citarum hulu. *Jurnal Perikanan Kelautan*, 2012. **3**(4).
33. Hardinawati, H., Analisis Kandungan Logam Berat Timbal (Pb) pada Hati, Daging dan Kulit Ikan Baronang (*Siganus Gutattus*) di Pulau Lae-lae. 2017, Universitas Islam Negeri Alauddin Makassar.
34. Ali, A., Studi kuantitas dan kualitas air Sungai Tallo sebagai sumber air baku. Jurusan Teknik Sipil, Universitas Hassanuddin. Makasar, 2012.
35. Setiawan, H., Akumulasi dan distribusi logam berat pada vegetasi mangrove di pesisir Sulawesi Selatan. *Jurnal Ilmu Kehutanan*, 2015. **7**(1): p. 12-24.
36. Mahmuda AF. Studi Analisis Kualitas dan Kuantitas Air Sungai Pampang Kota madya Makassar. Makassar: Universitas Hassanudin, 2007.

LAMPIRAN REVISI

COMMENT:

1. The title is suggested to be changed
 2. It was corrected in the Abstract and added in Spanish
 3. Thought out the text there were made several corrections. There are many sentences in Indonesian that were translated. Check that the manuscript does not present sentences in the Indonesian language
 4. The units of the heavy metals should be carefully corrected.
 5. The references should be carefully corrected as indicated in the author's instructions of the Gaceta Medica de Caracas
 6. The references should be translated into the English language, not into Indonesian
- These are all form corrections, the referee requires this correction to correct the scientific content of the manuscript

RESPONSE

All the comments above done by author. Thank you so much.

**Heavy Metal Levels of Lead and Cadmium in Baronang Fish from South Sulawesi,
Indonesia**

**Niveles de metales pesados de plomo y cadmio en el pez Baronang del sur de Sulawesi,
Indonesia**

*Muhammad Ikhtiar¹, Nur Ulmy Mahmud ¹, Rezky Aulia Yusuf ¹, Haeril Amir²

¹Public Health Department, Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia

²Nursing Department, Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia

ORCID :0000-0001-9598-5428¹

ORCID : 0000-0001-8592-0220¹

ORCID : 0000-0002-2350-0150²

*Corresponding author: Muhammad Ikhtiar

Email: muhammad.ikhtiar@umi.ac.id

Adress: Jalan. Urip Sumeharjo No.05, Kel. Panaikang, Kec. Panakkukang, Makassar City, South Sulawesi, Indonesia, 90231, Telp: +6285343723079

RECIBIDO: 16 DE NOVIEMBRE 2022

ACEPTADO: 24 DE NOVIEMBRE 2022

SUMMARY

Objective: The purpose of this study was to determine the content of the heavy metal lead (Pb) and cadmium (Cd) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

Methods: This was an observational descriptive approach research. The accidental sampling technique was used at two locations, Kayu Bangkoa pier and Kera-Kera pier. Baronang fish samples were taken using fishing rods or nets and each bait was examined and analyzed in the laboratory.

Result: The results of an examination of Baronang fish samples from each sample at three different points around the Bangkoa Wood Pier, indicated a lead (Pb) parameter for sample 1 of 0.14 µg/g, for sample 2 <0.001 µg/g, while sample 3 was <0.001. For the parameter of cadmium (Cd) the sample 1 yield was 10.087 mg/g, sample 2 was 0.085 mg/g and sample 3 0.086 mg/g. The results of the examination of the Baronang fish sample indicate that lead (Pb) and cadmium (Cd) parameters were below standard SNI 7387: 2009, namely lead (Pb) 0.3 and cadmium (Cd) 0.1. While the results of the examination of Baronang fish samples from each sample at three different points around the Kera-Kera Pier indicated a lead (Pb) parameters for sample 1 which was <0.001 µg/g, sample 2 <0.001 µg/g and sample 3 <0.001.

Conclusion: The Baronang fish samples at three points of Bangkoa Wood Pier presented Pb parameter below the SNI 7387: 2009 standard, namely lead (Pb) 0.3, while the results of the cadmium (Cd) examination of the three samples all had passed the maximum limit of the SNI standard 7387: 2009, that is 0.1.

Keywords: Baronang fish, cadmium (Cd), environmental, lead (Pb)

RESUMEN

Objetivo: El propósito de este estudio fue determinar el contenido del metal pesado plomo (Pb) y cadmio (Cd) en el pez Baronang en el muelle de Kayu Bangkoa y el muelle de Kera-Kera en la ciudad de Makassar, South Sulawesi, Indonesia en 2020.

Métodos: Se utilizó un enfoque observacional descriptivo. La técnica de muestreo accidental se utilizó en dos lugares, el muelle de Kayu Bangkoa y el muelle de Kera-Kera. Se tomaron muestras de peces Baronang utilizando cañas de pescar o redes y luego cada cebo se examinó y analizó en el laboratorio.

Resultado: Los resultados del examen de las muestras de pescado Baronang en tres puntos diferentes alrededor del muelle de madera de Bangkoa indicaron parámetros de plomo (Pb) de la muestra 1 de 0,14 µg/g, la muestra 2 <0,001 µg/g mientras que la muestra 3 fue <0.001. Para el parámetro de cadmio (Cd) la muestra de 1 fue de 0.087 mg/g, la muestra 2 de 0.085 mg/g y la muestra 3 de 0.086 mg/g. Lo que indican parámetros de plomo (Pb) y cadmio (Cd) para el pescado Baronang con valores por debajo del estándar SNI 7387: 2009, a saber, plomo (Pb) 0,3 y cadmio (Cd) 0,1. Mientras que los resultados del examen de muestras de pescado Baronang de tres puntos diferentes alrededor del muelle de Kera-Kera indicaron para parámetros de plomo (Pb) que la muestra 1 era <0,001 µg/g, la muestra 2 <0,001 µg/g y la muestra 3 <0.001.

Conclusión: Las muestras de pescado Baronang tomadas en tres puntos indicaron parámetros de plomo (Pb) por debajo del estándar SNI 7387: 2009, mientras que los de cadmio (Cd) superaron el límite máximo de la norma SNI 7387: 2009.

Introduction

Heavy metal contamination of aquatic ecosystems has been a problem in environmental health for decades. Heavy metals are highly hazardous contaminants due to their persistence, toxicity, and bioaccumulation in various segments of the environment, such as water, sediments, air, and biota (1, 2). Heavy metal concentrations lower than the required amounts may cause deficiency effects, while high concentrations of these metals lead to health problems (3). Toxic metals including lead (Pb) and cadmium (Cd) are hazardous because they can cause severe health problems in minute quantities (4, 5).

One of the most vulnerable environmental aspects of heavy metal contamination is water (6). Coastal ecosystems are very sensitive to heavy metal contaminants; sediments and marine organisms can accumulate pollutants, and fish can get enriched with metals that are subsequently transferred to man through human consumption (7).

Recently, heavy metal pollution in an aquatic ecosystem is becoming a critical issue. Based on the results of provenance analyses of heavy metals suggest that geogenic sources were the main source of water contamination (8, 9). Nonetheless, the industry activity, and human activities have increased concentrations of heavy metals above normal levels in the environment contributing to serious and widespread environmental issues leading to chronic toxicity (10-13).

In line with this, in Indonesia based on research conducted by the Oceanography Center of LIPI (The Indonesian Institute of Science), reported that high levels of metals are generally detected at the stations located close to the mainland, indicating that heavy metals are enrichment from terrestrial anthropogenic activities (14).

In a study about the incidence of Pb, and Cd in seafood from Africans on the Indian and the Red Sea coasts, it was found that some seafood from African Indian and the Red Sea coasts such as fishes have presented Cd and Pb concentrations higher than permitted limit by FAOUN/EU regions, indicating a possible threat to public health (15).

Based on a study on the bioaccumulation of Cadmium (Cd) and Chromium (Cr) contained in water and fish carried out in the Tallo Makassar river using quantitative and observational methods, it was found that the cadmium content in 3 research points was $<0.003 \text{ mg / L}$ (13). This cadmium content fulfills the requirements based on the determination of heavy metal content which refers to the government regulation PP number 82 of 2001 concerning Water Quality Management and Water Pollution Control with the designation of water with class II criteria, where the maximum cadmium (Cd) content is 0.01 mg/L . The content of heavy metal cadmium in this fish is not only caused by industrial activities but also due to the densely populated environmental conditions and the behavior of the people who always dispose of domestic waste in the area around the river (13).

Based on research conducted in the waters of Benoa Bay, Bali, it was found that the content of heavy metals Pb and Cd in the water had exceeded the quality standards for dissolved metals in water for marine biota as stated in the Bali Governor Regulation No. 16 of 2016. For the baronang

fish itself, the content of heavy metal Cd has exceeded the maximum limit of metal contamination, which is 0.1 mg / Kg, while the Pb content in some samples has exceeded the maximum limit of heavy metal contamination of SNI 7387: 2009 (16). Cd drug and supervisory food body (BPOM) 0.10, pb 0.20 NAV.

Fish are a significant bioindicator of heavy metal contamination in the aquatic ecosystem, as they are at the top of the food web and can accumulate considerable amounts of heavy metal in their tissues (13, 17-19). Generally, higher concentrations of metals were found in the liver and gills than in muscles (20). A study using a dry ashing-acid digestion method in fish collected from Sri Murni Lake and Kepong Metropolitan Lake around Kepong district, Kuala Lumpur, found that the highest concentration of Pb was detected in the gills ($0.151 \pm 0.12\text{mg/g}$) while Cd, was highly accumulated in the bones ($1.750 \pm 3.43\text{ mg/g}$) (21).

Based on the studies above, water and the ecosystem in it are very susceptible to being contaminated by heavy metals which have a broad impact on long-term public health conditions. Therefore, Periodic research and monitoring of water and fish quality should be given attention so that the quality of the lake ecosystem, especially in relation to heavy metal elements, is under control which makes activities involving water contact safe (10). Therefore, the purpose of this study was to determine the content of the heavy metal lead (Pb) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

Methods

The method used in this study was sampling using the incidental sampling method, at two locations of the Bangkoa Wood Pier and the Kera-Kera Pier. Baronang fish samples are taken using fishing rods or nets and each bait is then examined and analyzed in the laboratory. Primary data collection is done utilizing field observations or reviewing the research location by taking samples of Baronang fish then examined and analyzed in the laboratory.

Baronang fish examination samples were taken at six points at two locations of the bangkoa wood pier and the Kera-Kera pier. Then measured using Automic Absorption Spectrophotometer (AAS) equipment by laboratory staff, namely the South Sulawesi Provincial Health Office Laboratory.

Results

This research was conducted in two different places with sampling on 20 November 2020 at the Kayu Bangkoa pier in the Bulogading Village Ujung Pandang sub-district and on March 20, 2016, at Kera-Kera Pier in Tamalanrea Indah Village, Tamalanrea District, then the clean water samples were taken to the Great Hall Makassar Health Laboratory, South Sulawesi Province to find out the results of examining clean water samples.

Based on the results of the examination Great Hall Makassar Health Laboratory, South Sulawesi Province, with samples of Baronang Fish, to determine water quality based on SNI 7387: 2009. The Maximum Limit of Heavy Metal Contamination in Food is as follows: that categorize “Qualified (Q) = not reach the maximum limit)” and “Unqualified Qualified (UQ) = reach the maximum limit);

**Table 1. Results of Sampling of Baronang Fish at the Bangko Wood Pier Bulogading Village
Kec. Ujung Pandang**

No.	Sample Code	Sample	Checkup result		Level	
			Lead (Pb) (µg g)	Cadmium (Cd) (Ug / g)	Pb	Cd
1	Sample 1	Baronang fish	0.14	0.087	Q	Q
2	Sample 2	Baronang fish	<0.01	0.085	Q	Q
3	Sample 3	Baronang fish	<0.01	0.086	Q	Q

Source: Primary Data, 2020

*Q: Qualified

Table 1 shows that the samples of Baronang Fish originating from the Bangkoa Wood Pier were taken at the first point for lead (Pb), namely with sample code 1 0.14 (µg / g), sample 2 <0.01 (µg / g), and sample 3 <0.01 (µg/g). As for the results of the cadmium (Cd) sample, the sample code is 1 0.087 (µg/g), sample 2 0.085 (µg/g), and sample 3 0.086 (µg/g). From the results of the sample examination on Baronang Fish, all of them are still below the SNI 7387: 2009 standard, namely 0.03 mg/kg for lead (Pb), and cadmium also meets the requirements, namely 0.1 mg/kg.

Table 2. Results of Sampling of Baronang Fish at Kera-Kera Pier Kelurahan Tamalanrea Indah Kec. Tamalanrea

No.	Sample Code	Sample	Checkup result		Level	
			Lead (Pb) (µg / g)	Cadmium (Cd) (µg/g)	Pb	Cd
1	Until 1	Baronang fish	<0.01	0.170	Q	UQ
2	Sample 2	Baronang fish	<0.01	0.301	Q	UQ
3	Sample 3	Baronang fish	<0.01	0.386	Q	UQ

Source: Primary Data, 2020

*UQ: Unqualified

Table 2 shows that the samples of Baronang Fish originating from the Kera-Kera Pier were taken at the first point for lead (Pb), namely with sample code 1 <0.01 ($\mu\text{g} / \text{g}$), sample 2 <0.01 ($\mu\text{g} / \text{g}$), and sample 3 <0.01 ($\mu\text{g} / \text{g}$) is still below the standard of SNI 7387: 2009, namely 0.01 mg/kg . As for the results of the cadmium (Cd) sample with a sample code of 1 0.087 ($\mu\text{g} / \text{g}$), sample 2 0.085 ($\mu\text{g} / \text{g}$), and sample 3 0.086 ($\mu\text{g} / \text{g}$). From the results of the sample examination on Baronang Fish, all of them did not meet the requirements of the SNI 7387: 2009 standard, namely 0.01 mg/kg for cadmium (Cd) while the lead (Pb) met the requirements, namely 0.3 mg/kg .

Discussions

Fish has many health benefits due to its nutritional content. However, heavy metals contamination such as lead (Pb) and cadmium (Cd) in aquatic ecosystems, especially in fish, which is the daily consumption of the community, is a major threat to the quality of public health due to its negative impact due to long-term consumption (22).

The metals that pose the highest risks to human health are one of them is cadmium and lead which cause important complications in the nervous system, kidneys, bones, lungs, and cardiovascular system due to their toxicity and possible carcinogenic effect (23). Bahkan kontaminasi heavy metals have been linked to the development of mental retardation and even death in instances of very high exposure in the human body (20). Some of them contribute to the possible adverse effects, particularly in the fetus and young children (9).

Cd toxicity by consumption of contaminated fish can cause the following signs and symptoms: increased salivation, choking or vomiting, abdominal pain, vertigo, loss of consciousness, painful spasms of the anal sphincter, and impairment of renal function for severe toxicity (24, 25).

Various systems are affected due to Pb toxic exposure in fish, such as disruption of neurotransmitter function occurs due to Pb accumulation causing neurotoxicity and interfering with the immune system response (26). In humans, Pb can cause defined specific changes in gene and protein expression in response to lead challenges and determine the injurious effects of exposure to lead on a cellular level (27).

Acute Pb toxicity after ingestion of contaminated seafood usually occurs in the brain and kidney, and its absorption in the gastrointestinal tract is influenced by nutritional calcium and iron status and age (children adsorb more, and consequently, are more vulnerable than adults (28).

Exposure pathways (dietary and waterborne), environmental factors (salt-water or freshwater), and Pb binding capacity with protein decide the accumulation pattern of heavy metals exposure. The main cause of heavy metal contamination in the African countries bordering the Indian Ocean and the Red Sea port activity, tourism, and petroleum activities (15). In Pakistan, particularly in Swat River, heavy metal contamination might be due to geogenic (mafic and ultramafic rocks) and anthropogenic activities including industrialization, municipal water, pesticides application, wastewater from farm fields sewage sludge, automobile workshops, mining effluents, hotels effluents and urban districts including (Charsadda Malakand Agency and Swat) located near Swat River (29).

Similar to other countries, in Indonesia, especially in Makassar, heavy metal contamination of cadmium (Cd) and Lead (Pb) in waters and fish occurs due to various activities such as ports,

tourism, and fisheries activities (30). Besides that, the density of human settlements which causes a large amount of waste and industrial wastewater is also a major factor in pollution in water areas (30).

Even though the concentration of heavy metals in the waters, especially in fish, is still below the threshold, the control and monitoring of heavy metals in fish are most important to minimize associated fish poisoning (15). The monitoring of THM concentrations in river water, sediments, and fish muscles is, therefore, essential to make sure acquiescence with food security guidelines and resulting end-user safety (29).

Levels of lead (Pb) and Cadmium (Cd) Heavy Metals in Baronang Fish at Bangkoa Wood Pier

Water conditions contaminated by various metals will significantly affect the aquatic ecosystem in both inland and marine waters. Lead (Pb) is a metal that is widely used by humans, so this metal also causes contamination impacts on the environment. Pb metal pollution can also be caused by the entry of industrial waste containing Pb metal into water bodies. Furthermore, with the presence of the biomagnification process in the waters, the concentration of Pb metal will continue to increase. Heavy metals enter the fish's body tissues through several ways, namely: the respiratory tract (gills), the digestive tract (intestines, liver, kidneys) through the process of the food chain, and penetration through the skin (flesh muscle) (31).

The results of the examination at the Baronang Fish Testing Laboratory at the Makassar Health Laboratory Center, South Sulawesi Province that the samples of baronang fish originating from two different pier areas, in Bulogading and Tamalanrea Indah sub-districts were taken at 6 different points every 3 points per pier there were several comparisons of the number of levels obtained in sampling at two different piers.

The results of the examination of baronang fish samples at 3 points taken around the Bangkoa Wood Pier with lead (Pb) parameters stated that sample 1 with a result of $0.14\mu\text{g/g}$ at sample 1 was found, for sample 2 with a result $<0.001\text{ }\mu\text{g/g}$ while sample 3 resulted <0.001 and for the parameter of cadmium (Cd) with a sample result of 1 0.087 mg/g , for sample 2 results of 0.085 mg/g while for sample 3 with a result of 0.086 mg/g from the results of an examination of the Baronang Fish sample at three points with the parameters of lead (Pb) and cadmium (Cd) are still below the standard of SNI 7389: 2009. Meanwhile, the results of research based on Budiman's research at the Wangisgara station in the upstream Citarum river showed the cadmium (Cd) content in Betok fish was 11 ppb with a weight of 22.9 gr, for 92 ppb catfish with a weight of 36.9 gr and a high level of cadmium (Cd) in Betok fish. Cadmium (Cd) in Tilapia 105 ppb with a weight of 20.2 grams. The three types of fish have exceeded the quality standards set by SNI 7387: 2009 concerning the limit of metal contamination in fish meat and its processed products (32).

At point 2 there is the Losari beach platform around Losari Beach. There are houses, Stella Maris Hospital, Siloam Hospital, Arya Duta Makassar Hotel, Aerotel Smile Makassar, D'bugis Ocean, Anging Mammiri Hotel and community settlements that can produce liquid waste and solid which can cause heavy metal pollution. This also shows from the research that heavy metals originating from human activities can be industrial or household waste materials and are also caused by sludge from dirty drains, recycling, lead-containing manufactured products (gasoline,

paint, printing ink, main water pipes, tin solder cans, and battery casings. The solubility of heavy metals in water bodies is controlled by the degree of acidity of the water, the type, and concentration of metals and the state of mineral components (33).

Whereas at point 3 it is near the container port where around the container the entry and exit of cargo ships around the area can provide pollution such as Oil Spills, Hotel Pantai Gapura, Hotel Makassar Golden and Cafe Balliriate, and Zona Cafe. Borate metal was not identified because it is still influenced by sea characteristics such as tides, sea breezes, and saltwater infiltration.

Levels of Heavy Metal Lead (Pb) and Cadmium (Cd) in Baronang Fish at Kera-Kera Pier

Cadmium (Cd) enters the environment because of human activities. Cadmium (Cd) in water bodies can come from atmospheric deposits, dust, sewage processing water, and industrial wastewater. Cadmium enters the water more due to human activities such as industry where the waste products from the factory are discharged directly into the water which will accumulate at the bottom of the waters and form sediment. Cadmium (Cd) can also enter organisms that live in water where Cadmium (Cd) can enter via oral, inhalation, or dermal. Cd enters the body of an organism, for example, as fish.

From the results of the examination at the Baronang Fish Testing Laboratory at the Makassar Health Laboratory Center, South Sulawesi Province that the samples of Baronang Fish originating from two different pier areas, in Bulogading and Tamalanrea Indah Villages, were taken at 6 different points every 3 points per pier there were several comparisons of the number of levels obtained in sampling at two different piers.

The results of the examination of Baronang fish samples at 3 points taken around the Kera-Kera Pier with lead (Pb) parameters stated that sample 1 with a result <0.001 ug / g, for sample 2 with a result <0.001 ug / g while sampling 3 with a result < 0.001 and for the parameter of cadmium (Cd) with a sample yield of 1 0.170 mg / g, for sample 2 results of 0.301 mg / g while for sample 3 with a result of 0.386 mg / g from the results of an examination of the Baronang fish samples at three points with lead parameters (Pb) and cadmium (Cd) are still below the SNI 7387: 2009 standard, namely lead (Pb) 0.3, while the results of the examination of cadmium (Cd) of the three samples all have passed the maximum limit of the SNI 7387: 2009 0.1 standards.

This shows from the results of the research that the results of the examination of Baronang fish samples on the heavy metal Cadmium (Cd) were identified at point 1 in the vicinity of the Kera-Kera Pier where the area is adjacent to the pond, residential areas that have an impact on waste disposal and plastic waste disposal, and a stopover for passenger and sea boats where the boat uses fuel engines. Cadmium fuel contains up to 0.5 ppm. Based on Latif's research (2012), shows that the Tallo River has several settlements, the PLTU industry, the plywood factory industry, aquaculture, and agriculture (34).

Whereas at point 3 of the results of Setiawan's research demonstrated that the waters of the Tallo River are experiencing high pressure due to the presence of settlements, PLTU, aquaculture, and agriculture. And point 3 is also not far from the Pampang River, causing pollution and a lot of getting into the water (35). Mahmuda's study stated that the Pampang River is flanked by the Tallo River (70 km from the length of the main river) to the north there are several settlements, the

alcohol industry, the chocolate processing industry, and hospitals. so that Cadmium (Cd) that enters the body of an organism such as fish will accumulate in the kidneys and liver because these two organs are very specific to fight toxins that enter the body (36).

Conclusion

The results of an examination of Baronang fish samples from each sample at three different points around the Bangkoa Wood Pier found that the lead (Pb) parameters stated that of the sample examination on Baronang Fish, all the samples were still below the SNI 7387: 2009 standard, namely 0.03 mg/kg for lead (Pb) and cadmium also meet the requirements, 0.1 mg/kg.

The results of the examination of Baronang fish samples from each sample at three different points around the Kera-Kera Pier with lead (Pb) parameters stated that originating from the Kera-Kera Pier were taken at the first point for lead (Pb), found all samples still below the standard of SNI 7387: 2009, namely 0.01 mg/kg. As for the results of the cadmium (Cd) from all the samples above the maximum limit of the requirements of the SNI 7387: 2009 standard.

Suggestions

Suggestions to the public and industry who live and have activities in the coastal areas and streams around the Bnagkoa Timber Pier to reduce the risk of pollution such as direct disposal of garbage and waste into rivers/sea.

Near the Kera-Kera Pier, it is hoped that the community will maintain the sanitation of the industrial environment around the Tallo River basin which uses heavy metal materials such as Cadmium (Cd) to carry out better treatment of their waste before being discharged into the environment to reduce the pollutant load that enters the waters of the Tallo River, Makassar City.

Acknowledgments

The author is grateful to the Faculty of Public Health, Universitas Muslim Indonesia, Makassar, Indonesia, for numerous supports of this study.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Authors' Contributions

All contributors contributed significantly to this study and all authors agree with the content of the manuscript.

Conflict of Interest

The author(s) declared no potential conflicts of interest with respect to the research authorship, and/or publication of this article.

Availability of Data and Materials

All data generated or analyzed during this study are included in this published article.

Ethical Approval

This article received ethical clearance from *Ethic Commission, Universitas Muslim Indonesia*. Registration Number: UMI0320015671.

Significance for public health

Indonesia is one of the largest consumers of fish in the world. Heavy metal contamination has been a problem in environmental health that can affect human health. This study describes the content of the heavy metal lead (Pb) in the Baronang fish at Kayu Bangkoa pier and the Kera-Kera Pier in the city of Makassar, South Sulawesi, Indonesia in 2020.

REFERENCES

**COMMENT: THE REFERENCES SHOULDS BE CORRECTED CAREFULLY
ACCORDING TO THE AUTHORS INSTRUCTION OF GACETA MEDICA DE
CARACAS**

1. Gashkina NA, Moiseenko TI, Kudryavtseva LP. Fish response of metal bioaccumulation to reduced toxic load on long-term contaminated lake Imandra. *Ecotoxicology and environmental safety*, 2020. **191**: p. 110205.
2. Xie, Q., et al., Assessment of long-term effects from cage culture practices on heavy metal accumulation in sediment and fish. *Ecotoxicol Environ Saf*, 2020. **194**: p. 110433.
3. Alves, R.N., et al., Oral bioaccessibility of toxic and essential elements in raw and cooked commercial seafood species available in European markets. *Food Chemistry*, 2018. **267**: p. 15-27.
4. Jadoon, S., et al., Potentially toxic elements in drinking water and associated health risk assessment in Abbottabad city, northern Pakistan. *Desalination and water treatment*, 2019. **2019**: p. 392-402.
5. Qian, Y., et al., Assessment of metal mobility in sediment, commercial fish accumulation and impact on human health risk in a large shallow plateau lake in southwest of China. *Ecotoxicol Environ Saf*, 2020. **194**: p. 110346.
6. Ciazela, J., M. Siepak, and P. Wojtowicz, Tracking heavy metal contamination in a complex river-oxbow lake system: Middle Odra Valley, Germany/Poland. *The Science of the total environment*, 2018. **616-617**: p. 996-1006.
7. Bat, L., A. ÖZtekİN, and F. Sahin, Heavy Metal Detection in *Scorpaena Porcus* Linnaeus, 1758 from Sinop Coast of the Black Sea and Potential Risks to Human Health. *Current Agriculture Research Journal*, 2018. **6**.
8. Muhammad, S. and K. Ahmad, Heavy metal contamination in water and fish of the Hunza River and its tributaries in Gilgit–Baltistan: Evaluation of potential risks and provenance. *Environmental Technology & Innovation*, 2020. **20**: p. 101159.
9. Basim, Y. and Z. Khoshnood, Target hazard quotient evaluation of cadmium and lead in fish from Caspian Sea. *Toxicol Ind Health*, 2016. **32**(2): p. 215-20.
10. Hashim, Z., et al., An Analysis of Heavy Metals in Lakes of Former Tin Mining Sites in the City of Ipoh, Perak, Malaysia. *International Journal of Academic Research in Business and Social Sciences*, 2018. **8**.
11. Kusin, F.M., et al., Distribution of heavy metals and metalloid in surface sediments of heavily-mined area for bauxite ore in Pengerang, Malaysia and associated risk assessment. *CATENA*, 2018. **165**: p. 454-464.

12. Sakai, N. and M. Yoneda, Potential Health Risk of Heavy Metals in Malaysia. 2018. p. 19-32.
13. Jais, N.J., et al., Bioaccumulation of Cadmium (Cd) and Chromium (Cr) Heavy Metals in Water and Fish in the Tallo River Makassar. *Window of Public Health Journal*, 2020. 1(3): p. 261-274..
14. Harmesa, H., L. Lestari, and F. Budiyo, Distribution of Heavy Metals in Seawater and Sediments in Cimanuk Waters, West Java, Indonesia. *OLDI (Oceanology and Limnology in Indonesia)*, 2020. 5(1): p. 19-32.
15. Tamele I.J, Loureiro, P.V. Lead, mercury and cadmium in fish and shellfish from the Indian Ocean and Red Sea (African Countries): Public health challenges. *Journal of Marine Science and Engineering*, 2020. 8(5): p. 344.
16. Mardani, N.P.S., I.W. Restu, and A.H.W. Sari, Heavy Metal Lead (Pb) and Cadmium (Cd) Content in Water and Fish Bodies in Benoa Bay Waters, Bali. *Current Trends in Aquatic Science*, 2018. 1(1): p. 106-113.
17. Maury-Brachet, R., et al., Mercury contamination levels in the bioindicator piscivorous fish *Hoplias aimara* in French Guiana rivers: mapping for risk assessment. *Environmental Science and Pollution Research*, 2020. 27(4): p. 3624-3636.
18. Zhong, W., et al., Health risk assessment of heavy metals in freshwater fish in the central and eastern North China. *Ecotoxicology and environmental safety*, 2018. 157: p. 343-349.
19. Arkianti, N., N.K. Dewi, and N.K.T. Martuti, Kandungan logam berat timbal (Pb) pada ikan di Sungai Lemat Kabupaten Magelang. *Life Science*, 2019. 8(1): p. 54-63.
20. Maurya, P.K. and D. Malik, Bioaccumulation of heavy metals in tissues of selected fish species from Ganga river, India, and risk assessment for human health. *Human and Ecological Risk Assessment: An International Journal*, 2019. 25(4): p. 905-923.
21. Ishak, A.R., M.S.M. Zuhdi, and M.Y. Aziz, Determination of lead and cadmium in tilapia fish (*Oreochromis niloticus*) from selected areas in Kuala Lumpur. *The Egyptian Journal of Aquatic Research*, 2020. 46(3): p. 221-225.
22. Olmedo, P., et al., Determination of toxic elements (mercury, cadmium, lead, tin and arsenic) in fish and shellfish samples. Risk assessment for the consumers. *Environment International*, 2013. 59: p. 63-72.
23. Zuloaga, J., S. Gallego, and C. Ramírez, Contenido de Hg, Cd, Pb y As en especies de peces: revisión. *Vitae*, 2015. 22: p. 148-159.
24. Johri, N., G. Jacquillet, and R. Unwin, Heavy metal poisoning: the effects of cadmium on the kidney. *Biometals*, 2010. 23(5): p. 783-792.
25. Nordberg, G.F., Historical perspectives on cadmium toxicology. *Toxicology and applied pharmacology*, 2009. 238(3): p. 192-200.
26. Ishaque, A., et al., Toxic effects of lead on fish and human. *Biological and Clinical Sciences Research Journal*, 2020. 2020(1): p. e045-e045.
27. Gillis, B.S., Z. Arbieva, and I.M. Gavin, Analysis of lead toxicity in human cells. *BMC genomics*, 2012. 13(1): p. 1-12.
28. Flegal, A.R., Lead in tropical marine systems: a review. *Science of the Total Environment*, 1986. 58(1-2): p. 1-8.

29. Liu, M., et al., Contamination features, geo-accumulation, enrichments and human health risks of toxic heavy metal (oids) from fish consumption collected along Swat river, Pakistan. *Environmental Technology & Innovation*, 2020. **17**: p. 100554.
30. Noor, R.J., A. Kabangnga, and F. Fathuddin, Spatial Distribution and Contamination Factors of Heavy Metals on the Coastal City of Makassar. *Journal of Tropical Oceanology*, 2021. 24(1): p. 93-101.
31. Ma'ruf, M., Analysis of Heavy Metal Concentration in Baronang Fish (*Siganus* sp) and Aquatic Environment for Management of Bontang Coastal Areas. 2017, Thesis (unpublished). Mulawarman University.
32. Budiman, B.T.P., Y. Dhahiyat, and I. Rustikawati, Bioaccumulation of heavy metals pb (lead) and cd (cadmium) in fish meat caught in the upstream Citarum river. *Marine Fisheries Journal*, 2012. 3(4).
33. Hardinawati, H., Analysis of Lead (Pb) Heavy Metal Content in the Liver, Meat and Skin of Baronang Fish (*Siganus Gutattus*) on Lae-lae Island. 2017, Alauddin State Islamic University Makassar.
34. Ali, A., Study of the quantity and quality of Tallo River water as a source of raw water. Department of Civil Engineering, Hassanuddin University. Makassar, 2012
35. Setiawan, H., Accumulation and distribution of heavy metals in mangrove vegetation on the coast of South Sulawesi. *Journal of Forestry Sciences*, 2015. 7(1): p. 12-24.
36. Mahmuda AF. Analysis Study of Water Quality and Quantity of Pampang River, Makassar Municipality. Makassar: Hassanudin University, 2007.

LAMPIRAN LOA



Gaceta Médica de Caracas

Órgano oficial de la Academia Nacional de Medicina
y del Congreso Venezolano de Ciencias Médicas

Fundada el 13 de marzo de 1893

Caracas, November 28, 2022

MANUSCRIPT ACCEPTANCE

Distinguished Dr. Muhammad Ikhtiar and Haeril Amir:

We are pleased to inform you that your manuscript entitled: **Heavy Metal Levels of Lead and Cadmium in Baronang Fish from South Sulawesi, Indonesia**, has been accepted for publication in volume 130 (4) 2022 of the Gaceta Médica de Caracas.

Dr. Enrique López-Loyo

Editor-in-Chief

Dr. Anita Stern Israel

Senior Editor

Gaceta Médica de Caracas