

BUKTI KORESPONDENSI
ARTIKEL JURNAL NASIONAL TERAKREDITASI SINTA 1

Judul Artikel : Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

Jurnal : Journal of Public Health and Pharmacy

Penulis : **Andi Surahman Batara**, Idhar Darlis

No	Perihal	Tanggal
1	Bukti konfirmasi submit artikel dan artikel yang disubmit	29 Mei 2024
2	Peer Review by Editor in chief	17 Juli 2024
2	Bukti konfirmasi review danl revisi	6 Okt 2024 & Revisi 12 Okt 2024
3	Letter of Acceptance	23 Oktober 2024
4	Naskah di Publish	5 November 2024

1. Bukti Submit Artikel (29 May 2024)

Journal of Public Health and Pharmacy

Tasks5

EnglishView Site

surahman

Submission LibraryView Metadata

Submissions

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

Andi Surahman Batara, Idhar Darlis

SubmissionReviewCopyeditingProduction

Submission Files

20839-2

surahman, 35. Template-journal-of-Public-Health-and-Pharmacy-New (1).docx (2)

May 29, 2024

Manuscript Submission

23762-1

ayanix, 5422-Manuscript Revised Author-22597-1-18-20240727.docx

September 10, 2024

Manuscript Revised Author

Download All Files

Pre-Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
Notification Regarding Your Article	indraafrianto	-	0	☐
revision	surahman	surahman	2	☐
Hasil turnitin	surahman	-	0	☐

2. Peer review By Editor in chief

Download All

Pre-Review Discussions

Add discu

Name	From	Last Reply	Replies
Notification Regarding Your Article	indraafrianto	-	0
revision	surahman	surahman	2
Hasil turnitin	surahman	-	0

3.

3. Review By Reviewer (06 Okt 2024)

Journal of Public Health and Pharmacy

ISSN: 2775-4952

Website: <https://jurnal.unismuhpalu.ac.id/index.php/jphp>

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

ARTICLE INFO

Received: filled in by the editor

Accepted: filled in by the editor

Volume: filled in by the editor

Issue: filled in by the editor

DOI: 10.56338/jphp

KEYWORDS

Stunting; Parenting; 0-59
Months Old Infant;

ABSTRACT (250-300 Words)

Introduction: Parenting is a factor most significantly affect and contribute to the incidence of stunting in infants aged 0-59 months. Parenting has recently attracted many attention since parents are the one who have the most interaction with their own children. Hence, current research was carried out to know the correlation between parenting and the incidence of stunting in infants aged 0-59 months.

Methods: This research was carried out through quantitative method with *cross sectional approach*. In this case, the samples involved are 169 infants. The data obtained were further analyzed through logistic regression with the assistance of SPSS Version 26.

Results: There is a relationship between parenting and the incidence of stunting in infants aged 0-59 months old with the logistic regression P-value of <0.05 .

Conclusion: Parenting on the incidence of stunting in infants aged 0-59 months old is very important. In this case, knowledge and nutritional intake are necessary to support the growth and development of infants aged 0-59 months.

Publisher: Pusat Pengembangan Teknologi Informasi dan Jurnal Universitas Muhammadiyah Palu

INTRODUCTION

Stunting becomes an issue for countries with middle and lower income (1-3). The stunting rate recorded worldwide is around 152 million infants or 22.5% of children worldwide (4). In Asia, the rate reaches 81.7 million and more than one third of it is in Africa (4). In this case, the three Asian countries with the highest stunting incidence are India, Pakistan, and Bangladesh (5, 6). It is also confirmed by Indonesia whose stunting incidence increases annually, in which it reached 20.2% in 2017 and became 22.7% in 2018 (7).

Stunting is defined as a condition where a child experiences low growth and being retarded due to bad diet so they are at risk of death (8, 9). It can also be defined as low height that is indicated by height lower than -2 of the standard deviation of WHO. Many factors affecting the incidence of stunting. They are particularly divided into two: those are direct factor and indirect factor. In this case, the indirect factor includes food nutrition, disease, gender, family characteristics, breastfeeding pattern, and healthcare service (10). The lack of food nutrition contributes to at least 50% of baby mortality (11).

Family is the forefront factor affecting the incidence of stunting on children. This can be seen based on the facts in the field, where many children experiencing stunting come from family with low purchasing ability, dirty environment, and the unavailability of clean water (8). A research project that was carried out in Sambas Indonesia reported the correlation between the incidence of stunting and preterm birth, low birthweight, diarrhea, and immunization (12). In the research, among 559 children analyzed, 20.8 % of them experienced stunting at the age of 0-59 months old, indicating that the age range is the most susceptible to stunting incident in postpartum period.

Current research focuses on infants aged 0-59 months in Luwu Regency, considering that the prevalence of North Luwu Regency has stunting incidence reaching 29.8% in 2022 (7). According to the data reported electronically in Community-Based Nutrition Record and Report, the number of stunting incidence in February 2023 in Pattimang Village reaches 11.1%.

Dikomentari [Au1]: The abstract provides a concise summary of the study, presenting the research's objective, methodology, results, and conclusion. It clearly outlines the correlation between parenting and the incidence of stunting in infants aged 0-59 months. However, the abstract could benefit from a clearer explanation of the implications of the findings and their contribution to the broader field of study.

Dikomentari [Au2]: The introduction effectively establishes the global and regional context of stunting, emphasizing its prevalence and significance. It highlights the primary objective of investigating the effect of parenting on stunting incidence. However, the introduction could be improved by providing a more detailed conceptual framework that links parenting styles directly to child nutrition and health outcomes. Additionally, the statistics provided could be more updated or better contextualized to strengthen the background.

METHOD

The method applied in this research is quantitative with cross-sectional approach, aiming to know the effect of parenting on the incidence of stunting in infants aged 0-59 months old in Pattimang Village, Malangke District, North Luwu Regency. The samples involved are 169 infants aged 0-59 months old. The sampling was carried out through purposive sampling with the criteria that the infants aged 0-59 months old, lived in Pattimang, Village, and the mother or the parents of the infant can communicate well. The data collected were then processed through editing, coding, processing, and cleaning. Meanwhile, the data analysis was done using SPSS Software. In order to answer the research questions, inferential statistic was conducted through logistic regression. Data that has been analyzed were then described to obtain the results. This research has received ethical approval Number: RK. 155/KEPK/STIK/VII/2023.

RESULTS

Table 1. Distribution of Respondents Based on Education and Occupation in Pattimang Village, Malangke District, North Luwu Regency

Characteristics	Number	Percentage (%)
Education		
None	9	5.3
Elementary School	18	10.7
Secondary School	81	47.9
High School	55	32.5
D3/S1/S2	6	3.6
Total	169	100
Occupation		
Civil Servant	3	1.8
Private Employee	2	1.2
Housewife	109	64.5
Farmer	33	19.5
Laborer	22	13.0
Total	169	100

Table 1 above shows that the participants are mostly secondary school graduates by 81 people (47.9%), while the least is D3/S1/S2 by 6 people (3.6%). Furthermore, the respondents mostly worked as housewife by 109 people (64.5%), while the least is private employee by 2 people (1.2%).

Table 2. Distribution of parenting on the respondents in Pattimang Village Malangke District North Luwu Regency

Parenting	Total	Percentage (%)
Poor	101	59.8
Good	68	40.2
Total	169	100

Based on table 2 above, the distribution of poor parenting on the respondents is 101 children (59.8%), while those received good parenting is 68 children (40.2%).

Dikomentari [Au3]: The methods section outlines a quantitative, cross-sectional approach and details the study sample, criteria, and statistical analysis (logistic regression with SPSS). While the approach seems appropriate for the research objectives, the methodology lacks detail in areas such as the specifics of the data collection process, measurement of parenting practices, and the handling of confounding variables. Including these elements would provide a clearer understanding of the study's rigor and replicability.

Dikomentari [Au4]: The results are presented clearly, with tables detailing the characteristics of respondents, distribution of parenting practices, and stunting incidence. The relationship between poor parenting and stunting is statistically significant (P-value < 0.05), with an Exp(B) value suggesting a strong association. However, the results section could be strengthened by providing more in-depth interpretation and discussion of the data beyond the descriptive statistics.

Dikomentari [A5]: The following is an example of a table, including the table name and font size, that I have adjusted according to the template. Please apply similar adjustments to the other tables

Table 3. Distribution of Stunting Incidence on Infants aged 0-59 months old in Pattimang Village
Malangke District North Luwu Regency

Stunting	Total	Percentage (%)
Yes	33	19.5
No	136	80.5
Total	169	100

Based on table 3, it shows that the incidence of stunting on infants aged 0-59 months old occurred on 33 infants (19.5%), while those who were not stunting are 136 infants (80.5%).

Table 4. Test Results of Parenting on the Incidence of Stunting on Infants Aged 0-59 months old in Pattimang Village
Malangke District North Luwu Regency

Variable	B	S.E	Wald	Df	Sig	Exp(B)
Parenting	1.406	0.572	6.036	1	0.014	4.079

Based on the results presented on Table 4 of the logistic regression analysis, it shows that parenting affect the incidence of stunting with p-value below 0.05, that it 0.014 with Exp (B) value of 4.079.

DISCUSSION

Parenting and the Incidence of Stunting

The results of this research indicate that parenting affects the incidence of stunting in Pattimang Village of Luwu. In this case, among 169 respondents involved, 101 or 59.8% of them received poor parenting, showing that the rate is quite high. Furthermore, based on the identification results on the level of education of the respondents, most of the mothers are high school graduates by 81 people or 47.9%. Related to this fact, previous study has stated that the low level of education is an inhibiting factor of receiving knowledge leading to the incidence of stunting on children (13). In this case, the knowledge refers to the those applied when feeding the children with food containing rich vitamin and nutrition (14). Such cases frequently occur in lower and middle income countries (3).

Another previous research (15) also reported that parents' role and parenting contribute to the incidence of stunting, such as the research carried out by Pertiwi that reported that negative parenting can increase the risk of stunting by three times (16). Parenting is a part of the complexity of socio-economy, environment, and demography (17). In addition, parenting has also become the most significant part to handle stunting in current era, thus certain policy from the government is needed to handle the incidence of stunting, especially those related to stunting.

The Incidence of Stunting on Infants Aged 0-59 months Old

The results of this research also show that 33 infants or 19.5% of the respondents involved at the age of 0-59 months old experienced stunting. This age is surely susceptible to the incidence of stunting as reported by the research conducted by Patriota that the age range of 50-59 months old is susceptible and needs special attention (18). In addition, the results of this study are also supported by the survey carried out by Adhikari in Nepal (19) and Research by Khan in Pakistan (11).

In the context of Indonesia, Sari (15) has conducted relevant research on the main cause of stunting on 0-59 months old infants in Indonesia, in which the imbalance between the income of family living in the rural area

Dikomentari [Au6]: The discussion appropriately links the results to existing literature, acknowledging the importance of parenting in preventing stunting. The section effectively discusses the socioeconomic and educational factors associated with poor parenting practices. However, it could benefit from a deeper critical analysis, exploring possible mechanisms behind the association between parenting and stunting and comparing the findings with those of similar studies. Furthermore, it should address any limitations in the data or potential biases in more detail.

and urban area becomes one of the factors contributing to the rate of stunting. This reason is assured by the fact that most of the mothers in this research working as housewife by 109 people or 64.5%, followed by farmers by 33 people or 19.5%.

Infants at the age of 0-59 months old must also have special attention and concern from the parents in order to prevent the risk of stunting on the children. The involvement of family and support from the government are significant in preventing the incidence of stunting. Therefore, all parties need to take their own respective roles.

Effect of Parenting on the Incidence of Stunting on Infants at The Age of 0-59 months old

Based on the statistical analysis of this research, there is a correlation between the variables with P value below 0.05, that is 0.014 with Exp (B) value of 4.079. The results of this research are also supported by the previous research carried out by Mulyaningsih (2021) in Indonesia which found that the economy status of household and the education level of parents are the significant covariate of higher stunting risk (1).

Parenting is also significantly correlated with the characteristics of family household, that includes individual factors (gender, diarrhea history and birth place), family factor (head of family, mother, socio-cultural orientation, and family system factor) (10, 20).

Parenting takes an essential role in preventing stunting. Currently it has become a concern for many people since parents are the closest ones who have the most interaction with their children (21-24).

This research has limitations in samples that have the same location, in the future we recommend that research reach a wider population. From the method there needs to be further interview data to explore the mother's experience so it is recommended to use a mixed method. Further research is expected to see wider subjects, such as environmenta, economy factor, etc.

CONCLUSION

Based on the results of this research, it can be summed up that there is a significant correlation between the parenting and incidence of stunting on infants aged 0-59 months old. All authors contributed to this research starting from compiling the research, writing the manuscript and processing the research results.

Conflict of interest

No Conflict of interest

Funding

No Funding

Acknowledgment

Thanks all for Thank you to all parties who contributed to this research, especially the villagers of Pattimang Village, Malangke District, North Luwu Regency.

BIBLIOGRAPHY

1. Mulyaningsih T, Mohanty I, Widyaningsih V, Gebremedhin TA, Miranti R, Wiyono VH. Beyond personal factors: Multilevel determinants of childhood stunting in Indonesia. PLoS One. 2021;16(11):e0260265.
2. Victora CG, Christian P, Vdaletti LP, Gatica-Domínguez G, Menon P, Black RE. Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. Lancet. 2021;397(10282):1388-99.

Dikomentari [Au7]: The conclusion succinctly summarizes the study's key findings, emphasizing the significant correlation between parenting and stunting. It also briefly mentions contributions and implications, though these could be expanded to enhance the discussion of practical applications and recommendations for further research. A more explicit call for policy or community interventions based on the study's findings would add depth to this section.

3. Li Z, Kim R, Vollmer S, Subramanian SV. Factors Associated With Child Stunting, Wasting, and Underweight in 35 Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3(4):e203386.
4. UNICEF. Joint child malnutrition estimates: World Health Organization; 2023 [Available from: <https://www.who.int/data/gho/data/themes/topics/joint-child-malnutrition-estimates-unicef-who-wb>].
5. Headley D, Hoddinott J, Park S. Drivers of nutritional change in four South Asian countries: a dynamic observational analysis. *Matern Child Nutr*. 2016;12 Suppl 1(Suppl 1):210-8.
6. Gusnedi G, Nindrea RD, Purnakarya I, Umar HB, Andrafikar, Syafrawati, et al. Risk factors associated with childhood stunting in Indonesia: A systematic review and meta-analysis. *Asia Pac J Clin Nutr*. 2023;32(2):184-95.
7. RISKESDAS. Laporan Hasil Riset Kesehatan Dasar Tahun 2018 2018 [Available from: https://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-risikesdas-2018_1274.pdf].
8. WHO. Stunting in a nutshell 2020 [Available from: <https://www.who.int/news/item/19-11-2015-stunting-in-a-nutshell#:~:text=Stunting%20is%20the%20impaired%20growth.infection%2C%20and%20inadequate%20psychosocial%20stimulation>].
9. Rahmawaty S, Meyer BJ. Stunting is a recognized problem: Evidence for the potential benefits of ω -3 long-chain polyunsaturated fatty acids. *Nutrition*. 2020;73:110564.
10. Yani DI, Rahayuwati L, Sari CWM, Komariah M, Fauziah SR. Family Household Characteristics and Stunting: An Update Scoping Review. *Nutrients*. 2023;15(1).
11. Khan S, Zaheer S, Safdar NF. Determinants of stunting, underweight and wasting among children < 5 years of age: evidence from 2012-2013 Pakistan demographic and health survey. *BMC Public Health*. 2019;19(1):358.
12. Sartika AN, Khoirunnisa M, Meiyetrian E, Ermayani E, Pramesthi IL, Nur Ananda AJ. Prenatal and postnatal determinants of stunting at age 0-11 months: A cross-sectional study in Indonesia. *PLoS One*. 2021;16(7):e0254662.
13. Vaivada T, Akseer N, Akseer S, Somaskandan A, Stefopoulos M, Bhutta ZA. Stunting in childhood: an overview of global burden, trends, determinants, and drivers of decline. *Am J Clin Nutr*. 2020;112(Suppl 2):777s-91s.
14. Pradana Putri A, Rong JR. Parenting functioning in stunting management: A concept analysis. *J Public Health Res*. 2021;10(2).
15. Sari K, Sartika RAD. The Effect of the Physical Factors of Parents and Children on Stunting at Birth Among Newborns in Indonesia. *J Prev Med Public Health*. 2021;54(5):309-16.
16. Nabilah Z, Arman A, Patimah S. Analysis of Specific Nutrition Intervention Programs for Breastfeeding Mothers on Stunting Incidents in Children Aged 25-36 Months in the Work Area of the Antang Health Center, Makassar City. *An Idea Health Journal*. 2021;1(3).
17. Hukom EH, Hutomo WMP, Prakoso RJ. Relationship Between Complementary Foods and Family Income on Stunting in Toddlers. *An Idea Health Journal*. 2023;3(1).
18. Patriota É SO, Abrantes LCS, Figueiredo A, Pizato N, Buccini G, Gonçalves VSS. Association between household food insecurity and stunting in children aged 0-59 months: Systematic review and meta-analysis of cohort studies. *Matern Child Nutr*. 2024;20(2):e13609.
19. Adhikari RP, Shrestha ML, Acharya A, Upadhaya N. Determinants of stunting among children aged 0-59 months in Nepal: findings from Nepal Demographic and health Survey, 2006, 2011, and 2016. *BMC Nutr*. 2019;5:37.
20. Chen L, Vivekananda K, Guan L, Reupert A. Parenting and family experiences of Chinese fathers with mental illness. *J Psychiatr Ment Health Nurs*. 2023;30(2):267-77.
21. Abdulaziz R, Suryanti N, Setiawan AS. A Review on Maternal Parenting, Child's Growth Stunting, and Oral Health. *Eur J Dent*. 2024;18(1):26-40.
22. Lin YH, Gau SS. Association between morningness-eveningness and the severity of compulsive Internet use: the moderating role of gender and parenting style. *Sleep Med*. 2013;14(12):1398-404.
23. Amir H, Permatananda P, Cahyani DD, Langelo W, Rosita R, Sajodin S, et al. Enhancing skill conceptualization, critical thinking, and nursing knowledge through reflective case discussions: a systematic review. *J Med Life*. 2023;16(6):851-5.
24. Anto A, Telaumbanua f, Fuadah N. The Relationship of Family Role With Smoking Behaviour in Adolescent

Authors

. An Idea Nursing Journal. 2024;3(1).

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

ARTICLE INFO	ABSTRACT (250-300 Words)
Received: filled in by the editor Accepted: filled in by the editor Volume: filled in by the editor Issue: filled in by the editor DOI: 10.56338/jphp KEYWORDS Stunting; Parenting; 0-59 Months Old Infant;	Introduction: Parenting is a factor most significantly affect and contribute to the incidence of stunting in infants aged 0-59 months. Parenting has recently attracted many attention since parents are the one who have the most interaction with their own children. Hence, current research was carried out to know the correlation between parenting and the incidence of stunting in infants aged 0-59 months. Methods: This research was carried out through quantitative method with <i>cross sectional approach</i>. In this case, the samples involved are 169 infants. The data obtained were further analyzed through logistic regression with the assistance of SPSS Version 26. Results: There is a relationship between parenting and the incidence of stunting in infants aged 0-59 months old with the logistic regression P-value of <0.05.59,8% respondent of them received poor parenting and his education is only senior high school and This finding found that the higher the knowledge of parents, the more it will influence the provision of food to their children with good nutritional value. Conclusion: Parenting on the incidence of stunting in infants aged 0-59 months old is very important. In this case, knowledge and nutritional intake are necessary to support the growth and development of infants aged 0-59 months.
Publisher: Pusat Pengembangan Teknologi Informasi dan Jurnal Universitas Muhammadiyah Palu	

INTRODUCTION

Stunting becomes an issue for countries with middle and lower income (1-3). The stunting rate recorded worldwide is around 152 million infants or 22.5% of children worldwide (4). In Asia, the rate reaches 81.7 million and more than one third of it is in Africa (4). In this case, the three Asian countries with the highest stunting incidence are India, Pakistan, and Bangladesh (5, 6). It is also confirmed by Indonesia whose stunting incidence increases annually, in which it reached 20.2% in 2017 and became 22.7% in 2018 (7).

Stunting is defined as a condition where a child experiences low growth and being retarded due to bad diet so they are at risk of death (8, 9). It can also be defined as low height that is indicated by height lower than -2 of the standard deviation of WHO. Many factors affecting the incidence of stunting. They are particularly divided into two: those are direct factor and indirect factor. In this case, the indirect factor includes food nutrition, disease, gender, family characteristics, breastfeeding pattern, and healthcare service (10). The lack of food nutrition contributes to at least 50% of baby mortality (11).

Family is the forefront factor affecting the incidence of stunting on children. This can be seen based on the facts in the field, where many children experiencing stunting come from family with low purchasing ability, dirty environment, and the unavailability of clean water (8). A research project that was carried out in Sambas Indonesia reported the correlation between the incidence of stunting and preterm birth, low birthweight, diarrhea, and immunization (12). In the research, among 559 children analyzed, 20.8 % of them experienced stunting at the age of 0-59 months old, indicating that the age range is the most susceptible to stunting incident in postpartum period.

Current research focuses on infants aged 0-59 months in Luwu Regency, considering that the prevalence of North Luwu Regency has stunting incidence reaching 29.8% in 2022 (7). According to the data reported

Dikomentari [Au1]: The abstract is clear in terms of objectives, methods, results, and conclusion. However, consider emphasizing key findings such as the magnitude of the relationship between parenting and stunting. Clarify how "knowledge and nutritional intake" specifically support infant growth, as this is a crucial point in your conclusion.

Dikomentari [Au2]: The introduction provides global and national perspectives on stunting, but it could benefit from more context on why parenting, specifically in the local context of Pattimang Village, is a critical factor. Consider streamlining some sections for conciseness, especially where data on stunting rates are repeated. Discuss how cultural practices or socioeconomic factors in the study area influence parenting styles and consequently stunting.

electronically in Community-Based Nutrition Record and Report, the number of stunting incidence in February 2023 in Pattimang Village reaches 11.1%. The culture and geographical location of Pattimang village is far from Luwu city, which means that the reach of health workers is often hampered, especially since there are no hospitals or health centers that are complete in terms of infrastructure.

METHOD

The method applied in this research is quantitative with cross-sectional approach, This method is suitable for research that looks at the direct relationship between independent and dependent variables. aiming to know the effect of parenting on the incidence of stunting in infants aged 0-59 months old in Pattimang Village, Malangke District, North Luwu Regency. The samples involved are 169 infants aged 0-59 months old. The sampling was carried out through purposive sampling with the criteria that the infants aged 0-59 months old, lived in Pattimang, Village, and the mother or the parents of the infant can communicate well. The data collected were then processed through editing, coding, processing, and cleaning. Meanwhile, the data analysis was done using SPSS Software. In order to answer the research questions, inferential statistic was conducted through logistic regression. Data that has been analyzed were then described to obtain the results. This research has received ethical approval Number: RK. 155/KEPK/STIK/VII/2023. So, there are several confounding variables such as access to health and economics, but all of these are triggers from parenting patterns.

RESULTS

Table 1. Distribution of Respondents Based on Education and Occupation in Pattimang Village, Malangke District, North Luwu Regency

Characteristics	Number	Percentage (%)
Education		
None	9	5.3
Elementary School	18	10.7
Secondary School	81	47.9
High School	55	32.5
D3/S1/S2	6	3.6
Total	169	100
Occupation		
Civil Servant	3	1.8
Private Employee	2	1.2
Housewife	109	64.5
Farmer	33	19.5
Laborer	22	13.0
Total	169	100

Table 1 above shows that the education and occupation. Education is last school level and occupation is daily activities. participants are mostly secondary school graduates by 81 people (47.9%), while the least is D3/S1/S2 by 6 people (3.6%). Furthermore, the respondents mostly worked as housewife by 109 people (64.5%), while the least is private employee by 2 people (1.2%).

Table 2. Distribution of parenting on the respondents in Pattimang Village Malangke District North Luwu Regency

Parenting	Total	Percentage (%)
Poor	101	59.8
Good	68	40.2

Dikomentari [Au3]: The cross-sectional approach is suitable, but a brief justification for the choice of the sample size (169 infants) would add strength to the methodology. Details on how "poor" and "good" parenting were defined or measured should be included to ensure clarity in the analysis. Mention if and how potential confounding factors (e.g., economic status, health access) were controlled.

Dikomentari [Au4]: The tables provide a comprehensive breakdown of education, occupation, parenting, and stunting incidence. However, it would be helpful to briefly interpret the findings within the text immediately after each table for better clarity. Table 4's logistic regression results are critical. Discuss the significance of the Exp(B) value (4.079) to emphasize the strength of the association between parenting and stunting.

Dikomentari [A5]: The following is an example of a table, including the table name and font size, that I have adjusted according to the template. Please apply similar adjustments to the other tables

Total	169	100
--------------	------------	------------

Based on table 2 above **parenting is methods in guiding**, the distribution of poor parenting on the respondents is 101 children (59.8%), while those received good parenting is 68 children (40.2%).

Table 3. Distribution of Stunting Incidence on Infants aged 0-59 months old in Pattimang Village
Malangke District North Luwu Regency

Stunting	Total	Percentage (%)
Yes	33	19.5
No	136	80.5
Total	169	100

Based on table 3, **stunting is growth condition is not in line with age it** shows that the incidence of stunting on infants aged 0-59 months old occurred on 33 infants (19.5%), while those who were not stunting are 136 infants (80.5%).

Table 4. Test Results of Parenting on the Incidence of Stunting on Infants Aged 0-59 months old in Pattimang Village
Malangke District North Luwu Regency

Variable	B	S.E	Wald	Df	Sig	Exp(B)
Parenting	1.406	0.572	6.036	1	0.014	4.079

Based on the results presented on Table 4 of the logistic regression analysis, it shows that parenting affect the incidence of stunting with p-value below 0.05, that it 0.014 with Exp (B) value of 4.079.

DISCUSSION

Parenting and the Incidence of Stunting

The results of this research indicate that parenting affects the incidence of stunting in Pattimang Village of Luwu. In this case, among 169 respondents involved, 101 or 59.8% of them received poor parenting, showing that the rate is quite high. Furthermore, based on the identification results on the level of education of the respondents, most of the mothers are high school graduates by 81 people or 47.9%. Related to this fact, previous study has stated that the low level of education is an inhibiting factor of receiving knowledge leading to the incidence of stunting on children (13). In this case, the knowledge refers to the those applied when feeding the children with food containing rich vitamin and nutrition (14). Such cases frequently occur in lower and middle income countries (3).

Another previous research (15) also reported that parents' role and parenting contribute to the incidence of stunting, such as the research carried out by Pertiwi that reported that negative parenting can increase the risk of stunting by three times (16). Parenting is a part of the complexity of socio-economy, environment, and demography (17). In addition, parenting has also become the most significant part to handle stunting in current era, thus certain policy from the government is needed to handle the incidence of stunting, especially those related to stunting.

Dikomentari [Au6]: The discussion effectively ties findings to prior research but would benefit from further elaboration on how the local context influences the outcomes. There are suggestions for policy implications, but these could be developed further to propose specific interventions to improve parenting practices. Address the limitations more comprehensively, such as potential biases in self-reported data or the cross-sectional nature limiting causal inferences.

The Incidence of Stunting on Infants Aged 0-59 months Old

The results of this research also show that 33 infants or 19.5% of the respondents involved at the age of 0-59 months old experienced stunting. This age is surely susceptible to the incidence of stunting as reported by the research conducted by Patriota that the age range of 50-59 months old is susceptible and needs special attention (18). In addition, the results of this study are also supported by the survey carried out by Adhikari in Nepal (19) and Research by Khan in Pakistan (11).

In the context of Indonesia, Sari (15) has conducted relevant research on the main cause of stunting on 0-59 months old infants in Indonesia, in which the imbalance between the income of family living in the rural area and urban area becomes one of the factors contributing to the rate of stunting. This reason is assured by the fact that most of the mothers in this research working as housewife by 109 people or 64.5%, followed by farmers by 33 people or 19.5%.

Infants at the age of 0-59 months old must also have special attention and concern from the parents in order to prevent the risk of stunting on the children. The involvement of family and support from the government are significant in preventing the incidence of stunting. Therefore, all parties need to take their own respective roles.

Effect of Parenting on the Incidence of Stunting on Infants at The Age of 0-59 months old

Based on the statistical analysis of this research, there is a correlation between the variables with P value below 0.05, that is 0.014 with Exp (B) value of 4.079. The results of this research are also supported by the previous research carried out by Mulyaningsih (2021) in Indonesia which found that the economy status of household and the education level of parents are the significant covariate of higher stunting risk (1).

Parenting is also significantly correlated with the characteristics of family household, that includes individual factors (gender, diarrhea history and birth place), family factor (head of family, mother, socio-cultural orientation, and family system factor) (10, 20).

Parenting takes an essential role in preventing stunting. Currently it has become a concern for many people since parents are the closest ones who have the most interaction with their children (21-24).

This research has limitations in samples that have the same location, in the future we recommend that research reach a wider population. From the method there needs to be further interview data to explore the mother's experience so it is recommended to use a mixed method. Further research is expected to see wider subjects, such as environmenta, economy factor, etc.

CONCLUSION

Based on the results of this research, it can be summed up that there is a significant correlation between the parenting and incidence of stunting on infants aged 0-59 months old. All authors contributed to this research starting from compiling the research, writing the manuscript and processing the research results. **The researcher also realized that there were shortcomings and obstacles that could certainly influence the research results and there was another element, namely culture, which we suggested to be continued with a qualitative ethnographic design.**

Conflict of interest

No Conflict of interest

Funding

No Funding

Acknowledgment

Thanks all for Thank you to all parties who contributed to this research, especially the villagers of Pattimang Village, Malangke District, North Luwu Regency.

Dikomentari [Au7]: The conclusion restates the findings clearly but can be enhanced by providing actionable recommendations based on the results. Suggest practical steps for stakeholders (e.g., health workers, parents) to improve parenting practices to reduce stunting.

BIBLIOGRAPHY

1. Mulyaningsih T, Mohanty I, Widyarningsih V, Gebremedhin TA, Miranti R, Wiyono VH. Beyond personal factors: Multilevel determinants of childhood stunting in Indonesia. *PLoS One*. 2021;16(11):e0260265.
2. Victora CG, Christian P, Vidaletti LP, Gatica-Domínguez G, Menon P, Black RE. Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. *Lancet*. 2021;397(10282):1388-99.
3. Li Z, Kim R, Vollmer S, Subramanian SV. Factors Associated With Child Stunting, Wasting, and Underweight in 35 Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3(4):e203386.
4. UNICEF. Joint child malnutrition estimates: World Health Organization; 2023 [Available from: <https://www.who.int/data/gho/data/themes/topics/joint-child-malnutrition-estimates-unicef-who-wb>].
5. Headey D, Hoddinott J, Park S. Drivers of nutritional change in four South Asian countries: a dynamic observational analysis. *Matern Child Nutr*. 2016;12 Suppl 1(Suppl 1):210-8.
6. Gusnedi G, Nindrea RD, Purnakarya I, Umar HB, Andrafikar, Syafrawati, et al. Risk factors associated with childhood stunting in Indonesia: A systematic review and meta-analysis. *Asia Pac J Clin Nutr*. 2023;32(2):184-95.
7. RISKESDAS. Laporan Hasil Riset Kesehatan Dasar Tahun 2018 2018 [Available from: https://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-riskesdas-2018_1274.pdf].
8. WHO. Stunting in a nutshell 2020 [Available from: <https://www.who.int/news/item/19-11-2015-stunting-in-a-nutshell#:~:text=Stunting%20is%20the%20impaired%20growth,infection%2C%20and%20inadequate%20psychosocial%20stimulation>].
9. Rahmawaty S, Meyer BJ. Stunting is a recognized problem: Evidence for the potential benefits of ω -3 long-chain polyunsaturated fatty acids. *Nutrition*. 2020;73:110564.
10. Yani DI, Rahayuwati L, Sari CWM, Komariah M, Fauziah SR. Family Household Characteristics and Stunting: An Update Scoping Review. *Nutrients*. 2023;15(1).
11. Khan S, Zaheer S, Safdar NF. Determinants of stunting, underweight and wasting among children < 5 years of age: evidence from 2012-2013 Pakistan demographic and health survey. *BMC Public Health*. 2019;19(1):358.
12. Sartika AN, Khoirunnisa M, Meiyetrian E, Ermayani E, Pramesthi IL, Nur Ananda AJ. Prenatal and postnatal determinants of stunting at age 0-11 months: A cross-sectional study in Indonesia. *PLoS One*. 2021;16(7):e0254662.
13. Vaivada T, Akseer N, Akseer S, Somaskandan A, Stefopoulos M, Bhutta ZA. Stunting in childhood: an overview of global burden, trends, determinants, and drivers of decline. *Am J Clin Nutr*. 2020;112(Suppl 2):777s-91s.
14. Pradana Putri A, Rong JR. Parenting functioning in stunting management: A concept analysis. *J Public Health Res*. 2021;10(2).
15. Sari K, Sartika RAD. The Effect of the Physical Factors of Parents and Children on Stunting at Birth Among Newborns in Indonesia. *J Prev Med Public Health*. 2021;54(5):309-16.
16. Nabilah Z, Arman A, Patimah S. Analysis of Specific Nutrition Intervention Programs for Breastfeeding Mothers on Stunting Incidents in Children Aged 25-36 Months in the Work Area of the Antang Health Center, Makassar City. *An Idea Health Journal*. 2021;1(3).
17. Hukom EH, Hutomo WMP, Prakoso RJ. Relationship Between Complementary Foods and Family Income on Stunting in Toddlers. *An Idea Health Journal*. 2023;3(1).
18. Patriota É SO, Abrantes LCS, Figueiredo A, Pizato N, Buccini G, Gonçalves VSS. Association between household food insecurity and stunting in children aged 0-59 months: Systematic review and meta-analysis of cohort studies. *Matern Child Nutr*. 2024;20(2):e13609.
19. Adhikari RP, Shrestha ML, Acharya A, Upadhaya N. Determinants of stunting among children aged 0-59 months in Nepal: findings from Nepal Demographic and health Survey, 2006, 2011, and 2016. *BMC Nutr*. 2019;5:37.

20. Chen L, Vivekananda K, Guan L, Reupert A. Parenting and family experiences of Chinese fathers with mental illness. *J Psychiatr Ment Health Nurs*. 2023;30(2):267-77.
21. Abdulaziz R, Suryanti N, Setiawan AS. A Review on Maternal Parenting, Child's Growth Stunting, and Oral Health. *Eur J Dent*. 2024;18(1):26-40.
22. Lin YH, Gau SS. Association between morningness-eveningness and the severity of compulsive Internet use: the moderating role of gender and parenting style. *Sleep Med*. 2013;14(12):1398-404.
23. Amir H, Permatananda P, Cahyani DD, Langelo W, Rosita R, Sajodin S, et al. Enhancing skill conceptualization, critical thinking, and nursing knowledge through reflective case discussions: a systematic review. *J Med Life*. 2023;16(6):851-5.
24. Anto A, Telaumbanua f, Fuadah N. The Relationship of Family Role With Smoking Behaviour in Adolescent Authors. *An Idea Nursing Journal*. 2024;3(1).

Accepted (19-oktober 2024)

Journal of Public Health and Pharmacy

Task

English

View Site

surahman

Round 1 Submission

Notification

Reviewer's

Revisions

Notifications

[jphp] Editor Decision

2024-10-19 10:13 AM

Andi Surahman Batara, Idhar Darlis:

We have reached a decision regarding your submission to Journal of Public Health and Pharmacy, "Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old".

Our decision is to: **Accept Submission**

Dr. Ahmad Yani, S.K.M, M.Kes
Faculty of Public Health Universitas Muhammadiyah Palu
ahmadyani@unismuhpalu.ac.id

Journal of Public Health and Pharmacy

<https://jurnal.unismuhpalu.ac.id/index.php/jphp>
jpublichealpharm@unismuhpalu.ac.id

2024-10-06 02:09 PM

2024-10-19 10:13 AM

2024-10-23 02:18 AM

2024-10-24 11:04 AM

2024-10-24 11:08 AM

Q Search

Q Search

Upload File

24512-1

Manuscript Revised Author

5427-24146-1-S-20241002.docx

October 12

Manuscript Revised Author

Letter of Acceptance

5422/JPHP-UMP/X/2024

After a meticulous review process, we are pleased to inform you that your manuscript has been accepted for publication in Journal of Public Health and Pharmacy. The essential details for your accepted manuscript are as follows:

Manuscript ID : 5422
Title : **“Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old”**
Author : **Andi Surahman Batara^{1*}, Idhar Darlis²**
Affiliation : ^{1*}Faculty of Public Health, Universitas Muslim Indonesia, Sulawesi Selatan, Indonesia, andisurahman.batara@umi.ac.id
²Faculty of Public Health, Universitas Pejuang Republik Indonesia, Sulawesi Selatan, Indonesia, idhar17a1@gmail.com

Corresponding
Email* : andisurahman.batara@umi.ac.id

Your article is scheduled for publication in **Vol. 4 No. 3: NOVEMBER 2024 of Journal of Public Health and Pharmacy**, providing you with a glimpse of the anticipated publication timeline. Should you have any inquiries or require further assistance, please do not hesitate to contact our editorial team at ahmadyani@unismuhpalu.ac.id or jpublichealpharm@unismuhpalu.ac.id

Once again, thank you for choosing Journal of Public Health and Pharmacy. We eagerly anticipate the successful publication of your valuable contribution.

October 23rd, 2024
Best regards,



Dr. Ahmad Yani

Editor in Chief, Journal of Public Health and Pharmacy

CERTIFICATE OF PUBLICATION

This is to certify that

Andi Surahman Batara, Idhar Darlis

has successfully published the article titled

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

in the Journal of Public Health and Pharmacy, Volume 4, Number 3, (2024)

doi : 10.56338/jphp.v4i3.5422

This publication contributes to advancing knowledge in the fields of public health and pharmacy, showcasing valuable insights from experts across various disciplines.

Date of Publication: 5 Nov 2024

Editor in Chief



JPHP
Journal of Public Health and Pharmacy
UNIVERSITAS
MUHAMMADIYAH PALU

Dr. Ahmad Yani, S.K.M., M.Kes

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

Andi Surahman Batara^{1*}, Idhar Darlis²

¹Faculty of Public Health, Universitas Muslim Indonesia, Sulawesi Selatan, Indonesia, andisurahman.batara@umi.ac.id

²Faculty of Public Health, Universitas Pejuang Republik Indonesia, Sulawesi Selatan, Indonesia, idhar17a1@gmail.com

*Corresponding Author: E-mail: andisurahman.batara@umi.ac.id

ARTICLE INFO

Manuscript Received: 29 May, 2024

Revised: 12 Oct, 2024

Accepted: 23 Oct, 2024

Date of Publication: 05 Nov, 2024

Volume: 4

Issue: 3

DOI: [10.56338/jphp.v4i3.5422](https://doi.org/10.56338/jphp.v4i3.5422)

KEYWORDS

Stunting;
Parenting;
0-59 Months Old Infant

ABSTRACT

Introduction: Parenting is a factor most significantly affect and contribute to the incidence of stunting in infants aged 0-59 months. Parenting has recently attracted many attentions since parents are the one who have the most interaction with their own children. Hence, current research was carried out to know the correlation between parenting and the incidence of stunting in infants aged 0-59 months.

Methods: This research was carried out through quantitative method with cross sectional approach. In this case, the samples involved are 169 infants. The data obtained were further analyzed through logistic regression with the assistance of SPSS Version 26.

Results: There is a relationship between parenting and the incidence of stunting in infants aged 0-59 months old with the logistic regression P-value of <0.05. 59,8% respondent of them received poor parenting and his education is only senior high school and This finding found that the higher the knowledge of parents, the more it will influence the provision of food to their children with good nutritional value.

Conclusion: Parenting on the incidence of stunting in infants aged 0-59 months old is very important. In this case, knowledge and nutritional intake are necessary to support the growth and development of infants aged 0-59 months.

Publisher: Pusat Pengembangan Teknologi Informasi dan Jurnal Universitas Muhammadiyah Palu

INTRODUCTION

Stunting becomes an issue for countries with middle and lower income (1-3). The stunting rate recorded worldwide is around 152 million infants or 22.5% of children worldwide (4). In Asia, the rate reaches 81.7 million and more than one third of it is in Africa (4). In this case, the three Asian countries with the highest stunting incidence are India, Pakistan, and Bangladesh (5, 6). It is also confirmed by Indonesia whose stunting incidence increases annually, in which it reached 20.2% in 2017 and became 22.7% in 2018 (7).

Stunting is defined as a condition where a child experiences low growth and being retarded due to bad diet so they are at risk of death (8, 9). It can also be defined as low height that is indicated by height lower than -2 of the standard deviation of WHO. Many factors affecting the incidence of stunting. They are particularly divided into two: those are direct factor and indirect factor. In this case, the indirect factor includes food nutrition, disease, gender, family characteristics, breastfeeding pattern, and healthcare service (10). The lack of food nutrition contributes to at least 50% of baby mortality (11).

Family is the forefront factor affecting the incidence of stunting on children. This can be seen based on the facts in the field, where many children experiencing stunting come from family with low purchasing ability, dirty environment, and the unavailability of clean water (8). A research project that was carried out in Sambas Indonesia reported the correlation between the incidence of stunting and preterm birth, low birthweight, diarrhea, and

immunization (12). In the research, among 559 children analyzed, 20.8 % of them experienced stunting at the age of 0-59 months old, indicating that the age range is the most susceptible to stunting incident in postpartum period.

Current research focuses on infants aged 0-59 months in Luwu Regency, considering that the prevalence of North Luwu Regency has stunting incidence reaching 29.8% in 2022 (7). According to the data reported electronically in Community-Based Nutrition Record and Report, the number of stunting incidences in February 2023 in Pattimang Village reaches 11.1%. The culture and geographical location of Pattimang village is far from Luwu city, which means that the reach of health workers is often hampered, especially since there are no hospitals or health centers that are complete in terms of infrastructure.

METHOD

The method applied in this research is quantitative with cross-sectional approach, this method is suitable for research that looks at the direct relationship between independent and dependent variables. aiming to know the effect of parenting on the incidence of stunting in infants aged 0-59 months old in Pattimang Village, Malangke District, North Luwu Regency. The samples involved are 169 infants aged 0-59 months old. The sampling was carried out through purposive sampling with the criteria that the infants aged 0-59 months old, lived in Pattimang, Village, and the mother or the parents of the infant can communicate well. The data collected were then processed through editing, coding, processing, and cleaning. Meanwhile, the data analysis was done using SPSS Software. In order to answer the research questions, inferential statistic was conducted through logistic regression. Data that has been analyzed were then described to obtain the results. This research has received ethical approval Number: RK. 155/KEPK/STIK/VII/2023. So, there are several confounding variables such as access to health and economics, but all of these are triggers from parenting patterns.

RESULTS

Table 1. Distribution of Respondents Based on Education and Occupation in Pattimang Village, Malangke District, North Luwu Regency

Characteristics	Number	Percentage (%)
Education		
None	9	5.3
Elementary School	18	10.7
Secondary School	81	47.9
High School	55	32.5
D3/S1/S2	6	3.6
Total	169	100
Occupation		
Civil Servant	3	1.8
Private Employee	2	1.2
Housewife	109	64.5
Farmer	33	19.5
Laborer	22	13.0
Total	169	100

Table 1 above shows that the education and occupation. Education is last school level and occupation is daily activities. participants are mostly secondary school graduates by 81 people (47.9%), while the least is D3/S1/S2 by 6 people (3.6%). Furthermore, the respondents mostly worked as housewife by 109 people (64.5%), while the least is private employee by 2 people (1.2%).

Table 2. Distribution of parenting on the respondents in Pattimang Village Malangke District North Luwu Regency

Parenting	Total	Percentage (%)
Poor	101	59.8
Good	68	40.2
Total	169	100

Based on table 2 above parenting is methods in guiding, the distribution of poor parenting on the respondents is 101 children (59.8%), while those received good parenting is 68 children (40.2%).

Table 3. Distribution of Stunting Incidence on Infants aged 0-59 months old in Pattimang Village Malangke District North Luwu Regency

Stunting	Total	Percentage (%)
Yes	33	19.5
No	136	80.5
Total	169	100

Based on table 3, stunting is growth condition is not in line with age it shows that the incidence of stunting on infants aged 0-59 months old occurred on 33 infants (19.5%), while those who were not stunting are 136 infants (80.5%).

Table 4. Test Results of Parenting on the Incidence of Stunting on Infants Aged 0-59 months old in Pattimang Village Malangke District North Luwu Regency

Variable	B	S.E	Wald	Df	Sig	Exp(B)
Parenting	1.406	0.572	6.036	1	0.014	4.079

Based on the results presented on Table 4 of the logistic regression analysis, it shows that parenting affect the incidence of stunting with p-value below 0.05, that it 0.014 with Exp (B) value of 4.079.

DISCUSSION

Parenting and the Incidence of Stunting

The results of this research indicate that parenting affects the incidence of stunting in Pattimang Village of Luwu. In this case, among 169 respondents involved, 101 or 59.8% of them received poor parenting, showing that the rate is quite high. Furthermore, based on the identification results on the level of education of the respondents, most of the mothers are high school graduates by 81 people or 47.9%. Related to this fact, previous study has stated that the low level of education is an inhibiting factor of receiving knowledge leading to the incidence of stunting on children (13). In this case, the knowledge refers to the those applied when feeding the children with food containing rich vitamin and nutrition (14). Such cases frequently occur in lower- and middle-income countries (3).

Another previous research (15) also reported that parents' role and parenting contribute to the incidence of stunting, such as the research carried out by Pertiwi that reported that negative parenting can increase the risk of stunting by three times (16). Parenting is a part of the complexity of socio-economy, environment, and demography (17). In addition, parenting has also become the most significant part to handle stunting in current era, thus certain policy from the government is needed to handle the incidence of stunting, especially those related to stunting.

The Incidence of Stunting on Infants Aged 0-59 months Old

The results of this research also show that 33 infants or 19.5% of the respondents involved at the age of 0-59 months old experienced stunting. This age is surely susceptible to the incidence of stunting as reported by the research conducted by Patriota that the age range of 50-59 months old is susceptible and needs special attention

(18). In addition, the results of this study are also supported by the survey carried out by Adhikari in Nepal (19) and Research by Khan in Pakistan (11).

In the context of Indonesia, Sari (15) has conducted relevant research on the main cause of stunting on 0-59 months old infants in Indonesia, in which the imbalance between the income of family living in the rural area and urban area becomes one of the factors contributing to the rate of stunting. This reason is assured by the fact that most of the mothers in this research working as housewife by 109 people or 64.5%, followed by farmers by 33 people or 19.5%.

Infants at the age of 0-59 months old must also have special attention and concern from the parents in order to prevent the risk of stunting on the children. The involvement of family and support from the government are significant in preventing the incidence of stunting. Therefore, all parties need to take their own respective roles.

Effect of Parenting on the Incidence of Stunting on Infants at the Age of 0-59 months old

Based on the statistical analysis of this research, there is a correlation between the variables with P value below 0.05, that is 0.014 with Exp (B) value of 4.079. The results of this research are also supported by the previous research carried out by Mulyaningsih (2021) in Indonesia which found that the economy status of household and the education level of parents are the significant covariate of higher stunting risk (1).

Parenting is also significantly correlated with the characteristics of family household, that includes individual factors (gender, diarrhea history and birth place), family factor (head of family, mother, socio-cultural orientation, and family system factor) (10, 20).

Parenting takes an essential role in preventing stunting. Currently it has become a concern for many people since parents are the closest ones who have the most interaction with their children (21-24).

This research has limitations in samples that have the same location, in the future we recommend that research reach a wider population. From the method there needs to be further interview data to explore the mother's experience so it is recommended to use a mixed method. Further research is expected to see wider subjects, such as environmenta, economy factor, etc.

CONCLUSION

Based on the results of this research, it can be summed up that there is a significant correlation between the parenting and incidence of stunting on infants aged 0-59 months old. All authors contributed to this research starting from compiling the research, writing the manuscript and processing the research results. The researcher also realized that there were shortcomings and obstacles that could certainly influence the research results and there was another element, namely culture, which we suggested to be continued with a qualitative ethnographic design.

CONFLICT OF INTEREST

No Conflict of interest

FUNDING

No Funding

ACKNOWLEDGMENT

Thanks all for Thank you to all parties who contributed to this research, especially the villagers of Pattimang Village, Malangke District, North Luwu Regency.

BIBLIOGRAPHY

1. Mulyaningsih T, Mohanty I, Widyaningsih V, Gebremedhin TA, Miranti R, Wiyono VH. Beyond personal factors: Multilevel determinants of childhood stunting in Indonesia. *PLoS One*. 2021;16(11):e0260265.

2. Victora CG, Christian P, Vidaletti LP, Gatica-Domínguez G, Menon P, Black RE. Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. *Lancet*. 2021;397(10282):1388-99.
3. Li Z, Kim R, Vollmer S, Subramanian SV. Factors Associated With Child Stunting, Wasting, and Underweight in 35 Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3(4):e203386.
4. UNICEF. Joint child malnutrition estimates: World Health Organization; 2023 [Available from: <https://www.who.int/data/gho/data/themes/topics/joint-child-malnutrition-estimates-unicef-who-wb>].
5. Headey D, Hoddinott J, Park S. Drivers of nutritional change in four South Asian countries: a dynamic observational analysis. *Matern Child Nutr*. 2016;12 Suppl 1(Suppl 1):210-8.
6. Gusnedi G, Nindrea RD, Purnakarya I, Umar HB, Andrafikar, Syafrawati, et al. Risk factors associated with childhood stunting in Indonesia: A systematic review and meta-analysis. *Asia Pac J Clin Nutr*. 2023;32(2):184-95.
7. RISKESDAS. Laporan Hasil Riset Kesehatan Dasar Tahun 2018 2018 [Available from: https://kesmas.kemkes.go.id/assets/upload/dir_519d41d8cd98f00/files/Hasil-risikesdas-2018_1274.pdf].
8. WHO. Stunting in a nutshell 2020 [Available from: <https://www.who.int/news/item/19-11-2015-stunting-in-a-nutshell#:~:text=Stunting%20is%20the%20impaired%20growth,infection%2C%20and%20inadequate%20psychosocial%20stimulation>].
9. Rahmawaty S, Meyer BJ. Stunting is a recognized problem: Evidence for the potential benefits of ω -3 long-chain polyunsaturated fatty acids. *Nutrition*. 2020;73:110564.
10. Yani DI, Rahayuwati L, Sari CWM, Komariah M, Fauziah SR. Family Household Characteristics and Stunting: An Update Scoping Review. *Nutrients*. 2023;15(1).
11. Khan S, Zaheer S, Safdar NF. Determinants of stunting, underweight and wasting among children < 5 years of age: evidence from 2012-2013 Pakistan demographic and health survey. *BMC Public Health*. 2019;19(1):358.
12. Sartika AN, Khoirunnisa M, Meiyetrian E, Ermayani E, Pramesthi IL, Nur Ananda AJ. Prenatal and postnatal determinants of stunting at age 0-11 months: A cross-sectional study in Indonesia. *PLoS One*. 2021;16(7):e0254662.
13. Vaivada T, Akseer N, Akseer S, Somaskandan A, Stefopoulos M, Bhutta ZA. Stunting in childhood: an overview of global burden, trends, determinants, and drivers of decline. *Am J Clin Nutr*. 2020;112(Suppl 2):777s-91s.
14. Pradana Putri A, Rong JR. Parenting functioning in stunting management: A concept analysis. *J Public Health Res*. 2021;10(2).
15. Sari K, Sartika RAD. The Effect of the Physical Factors of Parents and Children on Stunting at Birth Among Newborns in Indonesia. *J Prev Med Public Health*. 2021;54(5):309-16.
16. Nabilah Z, Arman A, Patimah S. Analysis of Specific Nutrition Intervention Programs for Breastfeeding Mothers on Stunting Incidents in Children Aged 25-36 Months in the Work Area of the Antang Health Center, Makassar City. *An Idea Health Journal*. 2021;1(3).
17. Hukom EH, Hutomo WMP, Prakoso RJ. Relationship Between Complementary Foods and Family Income on Stunting in Toddlers. *An Idea Health Journal*. 2023;3(1).
18. Patriota É SO, Abrantes LCS, Figueiredo A, Pizato N, Buccini G, Gonçalves VSS. Association between household food insecurity and stunting in children aged 0-59 months: Systematic review and meta-analysis of cohort studies. *Matern Child Nutr*. 2024;20(2):e13609.
19. Adhikari RP, Shrestha ML, Acharya A, Upadhaya N. Determinants of stunting among children aged 0-59 months in Nepal: findings from Nepal Demographic and health Survey, 2006, 2011, and 2016. *BMC Nutr*. 2019;5:37.
20. Chen L, Vivekananda K, Guan L, Reupert A. Parenting and family experiences of Chinese fathers with mental illness. *J Psychiatr Ment Health Nurs*. 2023;30(2):267-77.
21. Abdulaziz R, Suryanti N, Setiawan AS. A Review on Maternal Parenting, Child's Growth Stunting, and Oral Health. *Eur J Dent*. 2024;18(1):26-40.

22. Lin YH, Gau SS. Association between morningness-eveningness and the severity of compulsive Internet use: the moderating role of gender and parenting style. *Sleep Med.* 2013;14(12):1398-404.
23. Amir H, Permatananda P, Cahyani DD, Langelo W, Rosita R, Sajodin S, et al. Enhancing skill conceptualization, critical thinking, and nursing knowledge through reflective case discussions: a systematic review. *J Med Life.* 2023;16(6):851-5.
24. Anto A, Telaumbanua f, Fuadah N. The Relationship of Family Role With Smoking Behaviour in Adolescent Authors. *An Idea Nursing Journal.* 2024;3(1).

Journal of Public Health and Pharmacy

Tasks5

EnglishView Site

surahman

Submissions

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

Andi Surahman Batara, Idhar'Darlis

Submission

Review

Copyediting

Production

Submission Files

20839-2

surahman, 3S_Template-Journal-of-Public-Health-and-Pharmacy-New 11).docx 12)

May 29, 2024

Manuscript Submission

23762-1

ayanlx, 5422-Manuscript RevisedAuthor-22597-1-18-20240727.docx

September 10, 2024

Manuscript Revised Author

Download All Files

Pre-Review Discussions

Name

indraafrianto

2024-07-17 10:39 PM

Notification Remusfurn

surahman

2024-07-19 05:26 PM

Hasil turaitin

surahman

2024-07-19 05:29 PM

Add discussion

Replies

Closed

https://jurnal.unismuhpalu.ac.id/index.php/PHP/\$\$call\$\$\$/tab/author-dashboard/author-dashboard-tab/fetch-tab?submissionId=5422&stageId=

Rourad 1 Status

Submission accepted.

Notifications

URhIJ	Editor Decision	2024-10-06 02:09 PM
URhRJ	Editor Decision	2024-10-19 10:13 AM
URh12J	Editor Decision	2024-10-23 02:18 AM
URhRJ	Editor Decision	2024-10-24 11:04 AM
URh11J	Editor Decision	2024-10-24 11:08 AM

Reviewer's Attachment5

2.4146-1	99_5422-Manuscript Blind Review-23764-1-4-20240910.docx	October 2, 2024
24291-1	5422-Manuscript BHand Review-23764-1-4-20240910-2.docx	October 6, 2024

Effect of Parenting on Stunting Incidence In Infant Aged 0-59 Months Old

Andi Surahman Batara, [dhar Darlis

Submission Review Copyediting Production

Copyediting Discussions

Name

From

Last Reply

Replies

Closed

ayanix

2024-10-19 10:17 AM

surahman

2024-10-19 06:01 PM

surahman

2024-10-20 05:01 PM

msyukran

2024-10-23 02:16 AM

Add discussion

• R.[22].Q[.lm'.lfln]

LoA Endosed

Copyedited

Q Search

24667-1 ayanix, CP_S422-Manuscript Revised Author-24666-1-6-20241012.docx

October 19, 2024

Manuscript Copy Editing

24970.1 tproduction, Galley Proof..docx

October 24, 2024

Galley Proof Manuscript

Submission Review Copyediting Production

Production Discussions

Name

From

Last Reply

Replies

Closed

ayanix

2024-10-23 06:33 PM

tproduction

2024-10-24 11:08 AM

tproduction

2024-11-02 09:58 AM

tproduction

2024-11-05 09:58 AM

Add discussion

Letter of Acceptance (LMA)

Proof Correction

Final file Production

Article Publication in the JI. I. if vol 4 No. 3 20241

Galleys

Turnitin-Manuscript Submission- 20839-2-2-20240529.docx

Effect of Parenting on Stunting Incidence in Infant Aged 0-59 Months Old

Andi Surahman Batara¹, Idhar Darlis²

¹ Faculty of Public Health, Universitas Muslim Indonesia, Sulawesi Selatan, Indonesia, andisurahman.batara@umi.ac.id

² Faculty of Public Health, Universitas Pejuang Republik Indonesia, Sulawesi Selatan, Indonesia, idhar17a1@gmail.com

•corresponding Author: E-mail: andisurahman.batara@umi.ac.id

ARTICLE INFO

Received: filled in by the editor

Accepted: filled in by the editor

Volume: filled in by the editor

Issue: filled in by the editor

DOI: 10.56338/jphp

KEYWORDS

Stunting, Parenting, 0-59 Months Old Infant

ABSTRACT (250-300 Words)

Introduction: Parenting is a factor most significantly affect and contribute to the incidence of stunting in infants aged 0-59 months. Parenting has recently attracted many attention since parents are the one who have the most interaction with their own children. Hence, current research was carried out to know the correlation between parenting and the incidence of stunting in infants aged 0-59 months.

Methods: This research was carried out through quantitative method with *cross sectional approach*. In this case, the samples involved are 169 infants. The data obtained were further analyzed through logistic regression with the assistance of SPSS Version 26.

Results: There is a relationship between parenting and the incidence of stunting in infants aged 0-59 months old with the logistic regression P-value of <0.05.

Conclusion: Parenting on the incidence of stunting in infants aged 0-59 months old is very important. In this case, knowledge and nutritional intake are necessary to support the growth and development of infants aged 0-59 months.

Publisher: Pusat Pengembangan Teknologi Informasi dan Jurnal Universitas Muhammadiyah Palu

INTRODUCTION

Stunting becomes an issue for countries with middle and lower income (1-3). The stunting rate recorded worldwide is around 152 million infants or 22.5% of children worldwide (4). In Asia, the rate reaches 81.7 million and more than one third of it is in Africa (4). In this case, the three Asian countries with the highest stunting incidence are India, Pakistan, and Bangladesh (5, 6). It is also confirmed by Indonesia whose stunting incidence increases annually, in which it reached 20.2% in 2017 and became 22.7% in 2018 (7).

Stunting is defined as a condition where a child experiences low growth and being retarded due to bad diet so they are at risk of death (8, 9). It can also be defined as low height that is indicated by height lower than -2 of the standard deviation of WHO. Many factors affecting the incidence of stunting. They are particularly divided into two: those are direct factor and indirect factor. In this case, the indirect factor includes *food* nutrition, disease, gender, family characteristics, breastfeeding pattern, and healthcare service (10). The lack of food nutrition contributes to at least 50% of baby mortality (11).

Family is the forefront factor affecting the incidence of stunting on children. This can be seen based on the facts in the field, where many children experiencing stunting come from family with low purchasing ability, dirty environment, and the unavailability of clean water (8). A research project that was carried out in Sambas Indonesia reported the correlation between the incidence of stunting and preterm birth, low birthweight, diarrhea, and immunization (12). In the research, among 559 children analyzed, 20.8 % of them experienced stunting at the age of 0-59 months old, indicating that the age range is the most susceptible to stunting incident in postpartum period.

Current research focuses on infants aged 0-59 months in Luwu Regency, considering that the prevalence of North Luwu Regency has stunting incidence reaching 29.8% in 2022 (7). According to the data reported electronically in Community-Based Nutrition Record and Report, the number of stunting incidence in February 2023 in Pattimang Village reaches 11.1%.

METHOD

The method applied in this research is quantitative with cross-sectional approach, aiming to know the effect of parenting on the incidence of stunting in infants aged 0-59 months old in Pattimang Village, Malangke District, North Luwu Regency. The samples involved are 169 infants aged 0-59 months old. The sampling was carried out through purposive sampling with the criteria that the infants aged 0-59 months old, lived in Pattimang, Village, and the mother or the parents of the infant can communicate well. The data collected were then processed through editing, coding, processing, and cleaning. Meanwhile, the data analysis was done using SPSS Software. In order to answer the research questions, inferential statistic was conducted through logistic regression. Data that has been analyzed were then described to obtain the results. This research has received ethical approval Number: IRK. 155/KEPK/STIK/V11/2023.

RESULTS

Table 1. Distribution of Respondents Based on Education and Occupation in Pattimang Village, Malangke District, North Luwu Regency

Characteristics	Number	Percentage (%)
Education		
None	9	5.3
Elementary School	18	10.7
Secondary School	81	47.9
High School	55	32.5
D3/S1/S2	6	3.6
Total	169	100
Occupation		
Civil Servant	3	1.8
Private Employee	2	1.2
Housewife	109	64.5
Farmer	33	19.5
Laborer	22	13.0
Total	169	100

Table 1 above shows that the participants are mostly secondary school graduates by 81 people (47.9%), while the least is D3/S1/S2 by 6 people (3.6%). Furthermore, the respondents mostly worked as housewife by 109 people (64.5%), while the least is private employee by 2 people (1.2%).

**Table 2. Distribution of parenting on the respondents in Pattimang Village
Malangke District North Luwu Regency**

Parenting	Total	Percentage (%)
Poor	101	59.8
Good	68	40.2
Total	169	100

Based on table 2 above, the distribution of poor parenting on the respondents is 101 children (59.8%), while those received good parenting is 68 children (40.2%).

**Table 3. Distribution of Stunting Incidence on Infants aged 0-59 months old in Pattimang Village
Malangke District North Luwu Regency**

Stunting	Total	Percentage (%)
Yes	33	19.5
No	136	80.5
Total	169	100

Based on table 3, it shows that the incidence of stunting on infants aged 0-59 months old occurred on 33 infants (19.5%), while those who were not stunting are 136 infants (80.5%).

**Table 4. Test Results of Parenting on the Incidence of Stunting on Infants Aged 0-59 months old in
Pattimang Village Malangke District North Luwu Regency**

Variable	B	S.E	Wald	Df	Sig	Exp(B)
Parenting	1.406	0.572	6.036	1	0.014	4.079

Based on the results presented on Table 4 of the logistic regression analysis, it shows that parenting affect the incidence of stunting with p-value below 0.05, that it 0.014 with Exp (B) value of 4.079.

DISCUSSION

Parenting and the Incidence of Stunting

The results of this research indicate that parenting affects the incidence of stunting in Pattimang Village of Luwu. In this case, among 169 respondents involved, 101 or 59.8% of them received poor parenting, showing that the rate is quite high. Furthermore, based on the identification results on the level of education of the respondents, most of the mothers are high school graduates by 81 people or 47.9%. Related to this fact, previous study has stated that the low level of education is an inhibiting factor of receiving knowledge leading to the incidence of stunting on children (13). In this case, the knowledge refers to the those applied when feeding the children with food containing rich vitamin and nutrition (14). Such cases frequently occur in lower and middle income countries (3).

Another previous research (15) also reported that parents' role and parenting contribute to the incidence of stunting, such as the research carried out by Pertiwi that reported that negative parenting can increase the risk of stunting by three times (16). Parenting is a part of the complexity of socio-economy, environment, and demography (17). In addition, parenting has also become the most significant part to handle stunting in current era, thus certain policy from the government is needed to handle the incidence of stunting, especially those related to stunting.

The Incidence of Stunting on Infants Aged 0-59 months Old

The results of this research also show that 33 infants or 19.5% of the respondents involved at the age of 0-59 months old experienced stunting. This age is surely susceptible to the incidence of stunting as reported by the research conducted by Patriota that the age range of 50-59 months old is susceptible and needs special attention (18). In addition, the results of this study are also supported by the survey carried out by Adhikari in Nepal (19) and Research by Khan in Pakistan (11).

In the context of Indonesia, Sari (15) has conducted relevant research on the main cause of stunting on 0-59 months old infants in Indonesia, in which the imbalance between the income of family living in the rural area and urban area becomes one of the factors contributing to the rate of stunting. This reason is assured by the fact that most of the mothers in this research working as housewife by 109 people or 64.5%, followed by farmers by 33 people or 19.5%.

Infants at the age of 0-59 months old must also have special attention and concern from the parents in order to prevent the risk of stunting on the children. The involvement of family and support from the government are significant in preventing the incidence of stunting. Therefore, all parties need to take their own respective roles.

Effect of Parenting on the Incidence of Stunting on Infants at The Age of 0-59 months old

Based on the statistical analysis of this research, there is a correlation between the variables with P value below 0.05, that is 0.014 with Exp (B) value of 4.079. The results of this research are also supported by the previous research carried out by Mulyaningsih (2021) in Indonesia which found that the economy status of household and the education level of parents are the significant covariate of higher stunting risk (1).

Parenting is also significantly correlated with the characteristics of family household, that includes individual factors (gender, diarrhea history and birth place), family factor (head of family, mother, socio-cultural orientation, and family system factor) (10, 20).

Parenting takes an essential role in preventing stunting. Currently it has become a concern for many people since parents are the closest ones who have the most interaction with their children (21-24).

This research has limitations in samples that have the same location, in the future we recommend that research reach a wider population. From the method there needs to be further interview data to explore the mother's experience so it is recommended to use a mixed method. Further research is expected to see wider subjects, such as environmental, economy factor, etc.

CONCLUSION

Based on the results of this research, it can be summed up that there is a significant correlation between the parenting and incidence of stunting on infants aged 0-59 months old. All authors contributed to this research starting from compiling the research, writing the manuscript and processing the research results.

Conflict of interest

No Conflict of interest

Funding

No Funding

Acknowledgment

Thanks all for Thank you to all parties who contributed to this research, especially the villagers of Pattimang Village, Malangke District, North Iwu Regency.

BIBLIOGRAPHY

Salinan 5422-Manuscript Submission-20839-2-2-20240529.docx

ORIGINALITY REPORT

5%

SIMILARITY INDEX

PRIMARY SOURCES

1	jurnal.unismuhpalu.ac.id Internet	63 words — 3%
2	www.businessperspectives.org Internet	13 words — 1%
3	www.coursehero.com Internet	11 words — 1%
4	garuda.kemdikbud.go.id Internet	8 words — < 1%
5	Haeril Amir, Muhammad Yusuf, Yuliana Syam, Andi Masyitha Irwan et al. "Comparison between traditional and disposable bed baths in Intensive Care Unit", European Journal of Clinical and Experimental Medicine, 2023 Crossref	7 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF