

Master Turnitin

Status of Sustainability of the Utilization of the Rabbitfish (*Siganus canaliculatus*) in the Seagrass Ecosystem of the West...

 05 Januari 2026 (02)

 UMI 2025

 Universitas Muslim Indonesia

Document Details

Submission ID

trn:oid:::1:3452757864

Submission Date

Jan 5, 2026, 10:10 AM GMT+8

Download Date

Jan 5, 2026, 10:13 AM GMT+8

File Name

3_ejabf_Issue_48-4011_P_1.pdf

File Size

1.2 MB

16 Pages

4,736 Words

26,123 Characters

6% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text
- Cited Text
- Small Matches (less than 15 words)

Exclusions

- 6 Excluded Sources

Match Groups

-  **10 Not Cited or Quoted 6%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 3%  Internet sources
- 3%  Publications
- 4%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 10 Not Cited or Quoted 6%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 3% Internet sources
- 3% Publications
- 4% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Publication	U R Pradini, H Arief. "Sustainability Status of Baung Fish in Public Waters, Kampar...	2%
2	Internet	www.preprints.org	<1%
3	Internet	faculty.uobasrah.edu.iq	<1%
4	Internet	www.bioflux.com.ro	<1%
5	Student papers	Universitas Terbuka	<1%
6	Internet	f1000research.com	<1%
7	Internet	repository.umi.ac.id	<1%

Egyptian Journal of Aquatic Biology & Fisheries
Zoology Department, Faculty of Science,
Ain Shams University, Cairo, Egypt.
ISSN 1110 – 6131
Vol. 28(3): 807 – 822 (2024)
www.ejabf.journals.ekb.eg



Status of Sustainability of the Utilization of the Rabbitfish (*Siganus canaliculatus*) in the Seagrass Ecosystem of the West Coast of South Sulawesi

Abdul Rauf^{1*}, Muhammad Ikhsan Wamnebo², Kamil Yusuf¹, Muhammad Yunus¹,
Muhammad Ilham Rauf¹

¹Department of Marine Science, Faculty of Fisheries and Marine Sciences, Universitas Muslim Indonesia, Indonesia

²Department of Aquaculture, Faculty of Fisheries and Marine Sciences Universitas Muslim Indonesia, Indonesia

*Corresponding Author: abdul.rauf@umi.ac.id

ARTICLE INFO

Article History:

Received: Feb. 7, 2024

Accepted: May 19, 2024

Online: June 17, 2024

Keywords:

Seagrass,
Rabbitfish,
Sustainability status,
Rabbitfish

ABSTRACT

The rabbitfish (*Siganus canaliculatus*) is a type of demersal fish living near the bottom of the water. This fish is often found in coral reef areas and seagrass beds. Currently, the use of the seagrass ecosystem is limited to fishing activities, which, beside the predators, target the rabbitfish broodstock, causing a decrease in the population of the seedlings. Therefore, it is necessary to develop the rearing activities of the rabbitfish seeds in this ecosystem applying technological innovations and using fixed net cages. However, the sustainability status of this rearing business is under question. Hence, the current research was designed to analyze the sustainability status of the rabbitfish utilization activities in the most extensive seagrass ecosystem on the western coast of South Sulawesi in Pangkep and Barru regencies, which is abundant with the seedlings of *Siganus canaliculatus*. To hit this target, the Rabbitfish method was implemented. Based on the ecological, economic, social, and technological in addition to the institutional and legal dimensions, the sustainability status of the rabbitfish utilization activities at the study site is generally categorized as quite sustainable, with an average ordination value of 58.54.

INTRODUCTION

The rabbitfish (*Siganus canaliculatus*) belongs to the Siganidae family, and it is a type of demersal fish that lives at the bottom or close to the bottom of the waters. This fish is often found in coral reefs and seagrass beds (Duffy, 2006; Feryatunet *al.*, 2012; Ismail *et al.*, 2023), both adult fish and their seeds. Currently, the use of seagrass ecosystems (Rappe, 2010; Firmansyah, 2022) has not been widely implemented and is still limited to fishing activities, hence with no precise restrictions, chances for overfishing are available.

The West Coast of South Sulawesi, especially in Pangkep and Barru regencies, has an extensive seagrass ecosystem, where many rabbitfish seedlings are found. These fish breeds are vulnerable to being targeted by predators and fishermen due to their small size, resulting in low survival rates to reach adulthood. In addition, the fishing of rabbitfish broodstock by fishermen has also caused a decrease in the population of these fish seedlings in seagrass ecosystems. To overcome this problem, it is necessary to develop aquaculture activities by raising the seeds of rabbitfish, which are abundant in this ecosystem, through technological innovation using fixed net cages installed in seagrass ecosystems and their surroundings (Rifai *et al.*, 2014; Martha *et al.*, 2018, Rahman *et al.*, 2019). The results of this cultivation activity are quite significant, it lasts for 3-4 months. Moreover, the size of fish seedlings can reach 4-5cm, with an average weight of 300-400 grams (Rauf *et al.*, 2017).

The sustainability of the cultivation business of rearing the rabbitfish in the seagrass ecosystem (Latuconsina *et al.*, 2013; Syukur, 2016) hasn't been figured out yet. In an effort to overcome this problem, a comprehensive study was needed on the sustainability status of fish farming businesses, so that the potential of the seagrass ecosystem can be preserved and utilized sustainably.

MATERIALS AND METHODS

Research location and time

This research was carried out in Pangkep and Barru regencies (West Coast of South Sulawesi) during May – December 2021. The location selection was based on the presence of seagrass ecosystem resources on the west coast of South Sulawesi.

Research methods

The methodological approach in this research involved using survey and questionnaire methods, along with direct interviews with respondents. Respondents were determined using a "purposive sampling" technique, namely that the selected respondents were expected to be able to understand and comprehend the problems to be studied. The research framework can be seen in Fig.(1).

Based on the research framework illustrated in Fig. (1), the potential of the rabbitfish on the West Coast of South Sulawesi is quite large, especially in seagrass ecosystems. The rabbitfish breeds are found in this ecosystem, therefore efforts are needed to maintain the population of these seedlings, resulting in a high survival until they reach maturity. One method of cultivation that can be used to maintain the survival of these fish is to use net cages in seagrass beds and their surroundings. To assess the sustainability of this aquaculture business, it is necessary to take into account five aspects in Rabbitfish analysis, namely ecological, economic, socio-cultural, technological, and institutional aspects.




Fig.1. Research framework

Data analysis

Analysis of the sustainability of rabbitfish farming businesses was carried out using a multidimensional scaling (MDS) approach called RAP-IB which is a modification of the RAP-FISH method used to assess the sustainability status of capture fisheries (Pitcher & Preikshot, 2001; Kavanagh & Pitcher, 2004; Ali, 2019). Sustainability analysis is expressed in the rabbitfish cultivation sustainability index (IKB-IB). The analysis was carried out through several stages:

- The determination of attributes in the rabbitfish farming business includes five main dimensions: ecological, economic, social, technological, and institutional (Fauzi, 2019). Each dimension has several attributes that serve as indicators of sustainability (Shumka *et al.*, 2023; Yusuf *et al.*, 2023). These attributes are taken from previous studies (Junaidi *et al.*, 2022; Satyawan *et al.*, 2023).
- Each attribute is assessed on an ordination scale based on sustainability criteria on each dimension. This assessment uses field observations and secondary data, along with scientific evaluations from scorers (Tetelepta *et al.*, 2022). Therefore, each

attribute is given a score that reflects the level of sustainability in the rabbitfish farming dimension. The score range is 0 to 3, or adjusted according to the special conditions of each attribute. The scale starts from bad (0) to good (3), where bad values reflect the least favorable conditions for sustainable development (Syamet *al.*, 2023; Youniset *al.*, 2023), while good values reflect the most favorable conditions. Between these two value extremes, there are one or more intermediate values, depending on the number of ranks on each attribute (Fauzi, 2019).

- RAPFISH analysis uses the multi-dimensional scaling (MDS) method to determine a single value point that reflects the relative position of seaweed farming. The success of the analysis can be seen from the low stress value, where the stress value that is considered good is less than 25 (Fauzi, 2019).
- In addition to ordination, RAPFISH has an analysis feature called leveraging. Leveraging helps detect dominant attributes by looking at changes in the ordination position (from bad to good), as they are omitted one by one. The leverage value ranges from 2 to 6%, measured through changes in the root mean square (RMS). If the RMS value has a significant increase, it shows that the attribute has a high sensitivity to sustainability (Fauzi, 2019).
- Monte-Carlo analysis is used to find sources of error in data variations (Fauzi, 2019). Errors in RAPFISH can occur due to several things, such as errors in attribute scoring due to lack of information, misunderstanding of attributes or scoring methods, scoring variations due to differences in opinion or judgment from different researchers, data input errors, or the existence of missing data, and high "stress" values of analysis results.

Preparation of index and sustainability status of rabbitfish cultivation business. The score value of each attribute is analyzed multidimensionally to determine one or several points that reflect the sustainable position of the rabbitfish cultivation business being studied relative to reference points, namely good points and bad points. The resulting index value is an illustration of the current status of the rabbitfish cultivation business. This value is determined by the score value of each attribute in each dimension studied (Fauzi, 2019). Sustainability status categories are presented in Table (1).

Table 1. Sustainability status categories for rabbitfish cultivation businesses based on the index value of the analysis results

Index value	Category
$\leq 24,9$	Unsustainable
25 – 49,9	Less sustainable
50– 74,9	Quite sustainable
>75	Sustainable

Source: Purwaningsih and Santosa (2015) and Soejarwo *et al.* (2019).

RESULTS

The assessment of the sustainability of the rabbitfish farming on the West Coast of South Sulawesi was carried out using the multi dimensionalscaling (MDS) approach using the Rap-Fish Tools, where this method was initially used to assess the sustainability status of capture fisheries (Pitcher & Preikshot, 2001; El Kifafet *et al.*, 2023). The results of this sustainability analysis are expressed in a business sustainability index which reflects the sustainability status of the implementation of the rabbitfish cultivation business development program based on existing conditions. The sustainable index value for each dimension of sustainability (ecological, economic, social, technological and institutional) is determined by providing a scoring value for each dimension which is the result of the assessment and opinion of the expert team (Fauzi, 2019; Hidayah *et al.*, 2020; Hamedet *et al.*, 2023). This sustainability analysis uses the Rapfish technique with the MDS method with five various dimensions, namely: ecological, economical, socio-cultural, technological, legal, and institutional.

The ordination results of the ecological dimension (Fig.2) on the horizontal axis show the difference in values from bad (0%) to good (100%) ordination results for the sustainability of the rabbitfish farming business, and the vertical axis shows the difference from the mix of evaluated attribute scores. The ordination analysis shows that ecological sustainability lies between bad and good. Moreover, the ordination analysis resulted in a quadratic correlation (R^2) of 95.25% and a stress value (S) of 15.57%.




Fig. 2. Results of the ecological dimension ordination analysis

To see each attribute as a leverage factor on the sustainability index value in the ecological dimension, a leverage analysis was carried out (Fig.3). The leverage analysis aims to detect the sensitivity of attributes in a dimension (Alder *et al.*, 2000; Fauzi, 2019). The greater the leverage value, the more sensitive the attribute, meaning that the greater the change in the attribute, the greater the impact on the level of sustainability, hence it needs attention. The leverage value for each attribute is as follows: (1) seed size

= 6.29; (2) distribution = 4.39 ; (3) seagrass meadow conditions = 5.38; (4) number of catches = 10.17; (5) catch size = 4.61; (6) utilization rate = 3.67.




Fig.3. Results of the ecological dimension leverage analysis

The result of the ordination of the economic dimension (Fig.4), resulted in a quadratic correlation (R^2) of 96.25% and a stress value (S) of 14.66%. This value indicates that all these attributes relate to the economic dimension. The stress value reflects the goodness of fit in multi-dimensional scaling (MDS) analysis, which indicates a measure of how precisely the configuration of a point can reflect the original data. A low stress value indicates good fit, while a high stress value indicates a good condition. In multi dimensionalscalling (MDS) analysis, a good stress value is less than 25% (Fauzi&Anna, 2005).




Fig.4. Results of the economic dimension ordination analysis

The leverage value of each attribute is as follows (Fig.5): (1) The rabbitfish business profit = 3.03; (2) GDP contribution of fisheries sector = 3.66; (3) market = 3.70; (4) income relative to UMR = 3.88; (5) other income = 2.79; (6) fisheries business cooperation = 7.51.




Fig.5. Results of the economic dimension leverage analysis

The results of the ordination of the socio-cultural dimension (Fig.6) resulted in a quadratic correlation (R^2) of 92.55% and stress value (S) of 14.36 %. This value indicates that all attributes are related to the socio-cultural dimension.




Fig.6. Results of the social dimension ordination analysis

The leverage value for each attribute (Fig. 7) is as follows: (1) conflict = 1.17; (2) local wisdom = 6.00; (3) knowledge about the environment = 8.26; (4) involvement of family members = 6.79; (5) frequency of meetings between residents = 9.26; (6) education level = 1.81.




Fig.7. Results of the leverage analysis of socio-cultural dimensions

The ordination results from the technological dimensions are presented in Fig. (8). The ordination analysis shows that sustainable technology lies between bad and good. Moreover, the ordination analysis with two iterations yielded a quadratic correlation (R^2) of 93.65% and a stress value (S) of 14.55%. This value indicates that all these attributes relate to the technology dimension.




Fig. 8. Results of the technological dimension ordination analysis

The leverage value for each attribute (Fig. 9) is as follows: (1) processing before sales = 0.96; (2) negative influence of fishing gear = 5.38; (3) nature and type of fishing gear = 7.48; (4) availability and mastery of information = 6.67; (5) technology utilization = 7.59; (6) facilities and infrastructure = 4.07.




Fig. 9. Results of the technological dimension leverage analysis

The ordination analysis (Fig.10) shows that the institutional and legal sustainability lies between bad and good. Moreover, the ordination analysis with 2 (two) iterations resulted in a quadratic correlation (R^2) of 94.28% and a stress value (S) of 14.24%. This value indicates how much the results of this analysis can be trusted. The closer to 100% the R^2 value, the more it is believed that all attributes are related to the legal and institutional dimensions.




Fig.10. Results of the legal and institutional dimension ordination analysis

The leverage value for each attribute is as follows: (1) fishing access restrictions = 6.90; (2) Financial institutions = 7.36; (3) involvement in decision making = 0.56; (4) law enforcement = 7.70; (5) user group = 11.49; (6) formal rules= 7.94 (Fig. 11).




Fig. 11. Results of the leverage analysis of legal and institutional dimensions.

The results of MDS and Monte Carlo analysis are presented in Table (2).

Table 2. Results of the MDS and Monte Carlo analysis

Index status	Status	Result	
		MDS	Monte Carlo
1. Ecology	Sustainable	78.80	77.09
2. Economy	Sustainable	84.44	83.87
3. Social-culture	Less sustainable	26.00	27.52
4. Technology	Quite sustainable	72.75	71.59
5. Law and Institutions	Less sustainable	30.72	31.40
Average amount	Quite sustainable	58.54	58.29

Fig. (12) shows that community activities in terms of ecological, economic and technological dimensions are sufficient to support the sustainability of the rabbitfish cultivation in seagrass ecosystems in the coastal areas of Pangkep and Barru regencies. Meanwhile, the socio-cultural, institutional and legal dimensions need to receive attention from the government, both district/city, provincial and central levels to ensure the sustainability of the rabbitfish cultivation.

The average number in Table (2) shows that the sustainability status of the rabbitfish farming business on the the West Coast of South Sulawesi is quite sustainable, with a MDS value of 58.54.




Fig.12. Diagram of the sustainability levels for the rabbitfish cultivation

DISCUSSION

1. Ecological dimension sustainability index

The RAPFISH analysis confirmed by MONTE CARLO (Fig. 2) shows that the ecological dimension in the rabbitfish farming business obtained a sustainability status value of 78.80. These results illustrate that from an ecological perspective, the effort is included in the sustainable category with a score above 75.

The results of the leverage analysis of ecological dimensions (Fig. 3) illustrate that the three most sensitive attributes are catch count (10.17), seedling size (6.29), and seagrass meadow condition (5.38). That is, in formulating policies to improve the sustainability status of the ecological dimension, it is necessary to pay attention and consider these three attributes.

2. Economic dimension sustainability index

The RAPFISH analysis confirmed by MONTE CARLO (Fig. 4) shows that the economic dimension in the rabbitfish farming business obtained a sustainability status value of 84.44. These results illustrate that from an economic perspective, the business is included in the sustainable category with a score above 75.

The results of the leverage analysis of the economic dimension (Fig.5) illustrate that the three most sensitive attributes are fisheries business cooperation (7.51), income relative to UMR (Regional Minimum Wage) (3.88), and market (3.70). That is, in formulating policies to improve the sustainability status of the economic dimension, it is necessary to pay attention and consider these three attributes.

3. Socio-cultural dimension sustainability index

The RAPFISH analysis confirmed by MONTE CARLO (Fig. 6) shows that the socio-cultural dimension in the rabbitfish farming business obtained a sustainability status value of 26.00. These results illustrate that from a socio-cultural perspective, the business is included in the less sustainable category with a score below 50.

The results of leverage analysis of the socio-cultural dimension (Fig. 7) illustrate that the three most sensitive attributes are the frequency of meetings between citizens (9.26), knowledge about the environment (8.26), and the involvement of family members (6.79). That is, in formulating policies to improve the sustainability status of the socio-cultural dimension, it is necessary to pay attention and consider these three attributes.

4. Technology dimension sustainability index

The RAPFISH analysis confirmed by MONTE CARLO (Fig. 8) shows that the technological dimension in the rabbitfish farming business obtained a sustainability status value of 72.75. These results illustrate that from a technological perspective, the business is included in the category of quite sustainable with a score below 75.

The results of the leverage analysis of the technology dimension (Fig. 9) illustrate that the three most sensitive attributes are utilization technology (7.59), nature and type of fishing gear (7.48), and availability and mastery of information (6.67). That is, in formulating policies to improve the sustainability status of the technological dimension, it is necessary to pay attention and consider these three attributes.

5. Sustainability index for legal and institutional dimensions

The RAPFISH analysis confirmed by MONTE CARLO (Fig.10) shows that the legal and institutional dimensions in the rabbitfish farming business obtained a sustainability status value of 26.00. These results illustrate that from the perspective of legal and institutional dimensions, the business is included in the less sustainable category with a score below 50.

The results of leverage analysis of legal and institutional dimensions (Fig. 11) illustrate that the three most sensitive attributes are the frequency of meetings between citizens (9.26), knowledge of the environment (8.26), and involvement of family members (6.79). That is, in formulating policies to improve the status of sustainability from the legal and institutional dimensions, it is necessary to pay attention and consider these three attributes.

MDS and Monte Carlo analysis

The Monte Carlo simulation aims to see the level of disturbance to the ordination value (Nababan et al., 2017; Fauzi, 2019). By carrying out 30 iterations, the Monte Carlo analysis using a scatter plot shows that the ordination value is quite stable, therefore it is believed that the ordination value has a fairly high level of confidence. Additionally, the scatter plot results, the Monte Carlo and MDS simulations are shown in Table (2) and Fig.(12).

CONCLUSION

Based on the description of the results and previous discussion, it can be concluded that:

- The rabbitfish cultivation (rearing) in the seagrass ecosystem has the potential to be developed as an alternative livelihood for fishermen on the West Coast of South Sulawesi.
- The sustainability status of the rabbitfish cultivation business in the west coastal region of South Sulawesi in terms of the ecological dimension with an ordination value of 78.80 is a good sustainability level category, the economic dimension with an ordination value of 84.44 is a good sustainability level category, the socio-cultural dimension with an ordination value 26.00 is a sufficient level of sustainability category, the technology dimension with an ordination value of 72.75 is a sufficient level of sustainability, the institutional and legal dimensions with an ordination value of 30.72 are a sufficient level of sustainability.
- The sustainability status of the rabbitfish utilization activities in the West Coastal region of South Sulawesi is generally categorized as quite sustainable, with an average ordination value of 58.54.

REFERENCES

- Ali, S. A.** (2019). Rapfish Analysis to Assess the Status of the Sustainability of Capture Fisheries Systems in Bone Bay. *International Journal of Science and Research.*, 6(9):817–826.
<https://doi.org/10.21275/ART2017 6104>
- Duffy, J.E.** (2006). Biodiversity and the functioning of seagrass ecosystems. *Marine ecology progress series*, Vol. 311: 233–250. USA.
<https://doi.org/10.3354/meps311233>
- El Kifaf, A.; Farmayanti, N. and Dewi, T. G.** (2023). Sustainability Of Milk Fish Farm Using Multi-Dimensional Scalling (MDS) Approach. *Societa: Journal of Agribusiness Sciences.*, 12(1): 1-9.
<https://doi.org/10.32502/jsct.v12i1.6305>
- Fauzi, A. and S. Anna.** (2005). *Fisheries and Marine Resources Modeling: for Policy Analysis*. PT. Gramedia Pustaka Utama, Jakarta.
- Fauzi, A.** (2019). *Sustainability Analysis Techniques*. Jakarta: PT Gramedia Pustaka Utama.
- Feryatun, F.; B. Hendrarto., andN. Widyorini.** (2012). Density and Distribution of Seagrass Based on Different Activity Zones in the Waters of Pramuka Island, Seribu Islands. *Journal of Management of Aquatic Resources*. Volume, Number, Year 2012, Pages 44-50.
<https://doi.org/10.14710/marj.v1i1.255>
- Firmansyah, F.; Nurafni, N.; Koroy, K. and Wahab, I.** (2022). Seagrass Ecosystem Status on Dodola Island, Morotai Island Regency. *Aurelia Journal.*, 4(2): 229-238.

<http://dx.doi.org/10.15578/aj.v4i2.11371>

- Hamed, E. S. A. E.; Rashedy, S. H.; Ahmed, H. O. and Al-Amgad, Z.** (2023). Marine Algae Based-Meal as Substitute Ingredient Feeds for the Fighting Conch *Strombus tricornis* L. (Mollusca, Gastropoda): Body Weights and Histomorphological Studies. *Egyptian Journal of Aquatic Biology and Fisheries.*, 27(6): 687–706. <https://doi.org/10.21608/ejabf.2023.329960>
- Hidayah, Z.; Nuzula, N. I. and Wiyanto, D. B.** (2020). Analysis of the Sustainability of Fisheries Resource Management in the Madura Strait, East Java. *GadjahMada University Fisheries Journal.*, 22(2): 101-111. <https://doi.org/10.22146/jfs.53099>
- Ismail, Z.; Kasim, F.; and Panigoro, C.** (2023). Density and Diversity and Current Conditions of the Seagrass Ecosystem in the Waters of Ponelo Island. *The NIKE Journal.*, 11(3): 123-128. <https://doi.org/10.37905/nj.v11i3.21984>
- J. Alder, T.J.; Pitcher, D.; Preikshot, K.; Kaschner, and B. Feriss.** (2000). “How good is good? A Rapid appraisal technique for evaluation of the sustainability status of fisheries of the North Atlantic. In Pauly and Pitcher (eds). *Methods for evaluating the impacts of fisheries on the north Atlantic ecosystem*”, Fisheries Center Research Reports., 8(2): 136 – 182. <http://dx.doi.org/10.14288/1.0348147>
- Junaidi, R.; Efizon, D. and Adriman, A.** (2022). Sustainability Status of Management of Terubuk Fish(*Tenualosamacrura*) In Bengkalis District. *Asian Journal of Aquatic Sciences.*, 5(2): 301-314. <https://doi.org/10.31258/ajoas.5.2.301-314>
- Kavanagh, P. and T.J. Pitchers.** (2004). Implementing Microsoft Excel Software For Rapfish: A Technique for The Rapid Appraisal of Fisheries Status. University of British Columbia, Canada. *Fisheries Center Research Reports.*, 12 (2): 75 p. <https://dx.doi.org/10.14288/1.0074801>
- Latuconsina, H.; Ambo-Rappe, R. and Nessa, M. N.** (2012). Species Composition and Community Structure of Seagrass Fish in the Waters of Tanjung Oyster, Inner Ambon Bay. *Journal of Tropical Marine Science and Technology.*, 4 (1): 35-46. <https://doi.org/10.29244/jitkt.v4i1.7804>
- Latuconsina, H.; Ambo-Rappe, R. and Nessa, M. N.** (2013). Association of Rabbit fish (*Siganus canaliculatus* Park, 1797) in the Seagrass Ecosystem of Inner Ambon Bay Waters. In: Simanjuntak CPH (eds.). *Proceedings of the National Fish Seminar VII*. Indonesian Ichthyology Society. pp. 123-137.
- Martha L. G. M. R.; G. S. P. Julyantoro and H. W. A. Sari.** (2019). Condition and Diversity of Seagrass Types in the Waters of Serang Island, Bali Province. *Journal of Marine Aquatic Science.*, 5(1): 131-141. <https://doi.org/10.24843/jmas.2019.v05.i01.p16>
- Nababan, B. O.; Sari, Y. D. and Hermawan, M.** (2017). Sustainability Analysis of Small-Scale Capture Fisheries in Tegal Regency, Central Java (Rapfish Approach Technique). *Journal of Maritime and Fisheries Socioeconomics.*, 2(2): 137. <https://doi.org/10.15578/jsekp.v2i2.5868>.

-
- Pitcher, T. J. and D. B. Preikshot.** (2001). Rapfish: A Rapid Appraisal Technique to Evaluate the Sustainability Status of Fisheries. *Fisheries Research.*, 49(3): 255-270.
[https://doi.org/10.1016/S0165-7836\(00\)00205-8](https://doi.org/10.1016/S0165-7836(00)00205-8)
- Purwaningsih, R. and Santosa, H.** (2015). Development of Sustainability Assessment Methods for Fisheries Industry Clusters. *Proceedings of the 6th SNST*, 1-6.
- Rappe, R. A.** (2010). Fish Community Structure in Different Seagrass Fields on Barrang Lompo Island. *Journal of Tropical Marine Science and Technology.*, 2 (2): 62-73.
<https://doi.org/10.29244/jitkt.v2i2.7853>
- Rahman, I. and Astriana, B. H.** (2019). Counseling regarding the seagrass ecosystem as an effort to preserve the ecosystem in the waters of Sire Beach, North Lombok Regency. *Abdi Human Journal.*, 6(2): 251-258.
<https://doi.org/10.29303/abdiinsani.v6i2.245>
- Rauf, A.; Asni, A.; Hamsiah, and Asmidar.** (2017). Evaluation of the Suitability of Rabbit fish Cultivation Land in the Seagrass Ecosystem on the West Coast of South Sulawesi., 2(1): 58–63.
<https://doi.org/10.24198/jaki.v2i1.23413>
- Rifai, U.; Hendarto, N. and Gani, A.** (2014). Raising Rabbit fish in Floating Net Karamba. (online) <http://um4rbbl.blogspot.com>. Accessed on November 10 2014 at 22.00 WITA. Makassar.
- Satyawan, N. M.; Tanjov, Y. E.; Purwanto, A.; Jaya, M. M.; Khikmawati, L. T.; Sarasati, W. and Bramana, A.** (2023). Sustainability Status of The Ecological Dimension in The Fisheries Management of Bali Sardine (*Sardinella lemuru* Bleeker 1853) in The Bali Strait. *Journal of Tropical Biology.*, 23(2): 272-281.
<https://doi.org/10.29303/jbt.v23i2.4855>
- Shumka; Spase; Yukio Nagahama; Sarjmir Hoxha, and Koji Asano.** (2023). “Overfishing and Recent Risk for Collapse of Fishery in Coastal Mediterranean Lagoon Ecosystem (Karavasta Lagoon, Southeastern Adriatic Sea).” *Fisheries and Aquatic Sciences.*, 26(4): 294–303.
<https://doi.org/10.47853/FAS.2023.e25>
- Syam, A. R.; Rahman, A.; Taukhid, I.; Rahmia, M. and Putri, A.** (2023). *Eco-environmental assessment in the Sembilan Archipelago, Indonesia: its relation to the abundance of humphead wrasse and coral reef fish composition.*, 26(1): 738–751.
<https://doi.org/10.47853/FAS.2023.e66>
- Syukur, A.** (2016). Seagrass Conservation for Sustainability of Fish Resources in Indonesian Coastal Waters. *Journal of Tropical Biology.*, 16(1): 56-68.
<https://doi.org/10.29303/jbt.v16i1.217>
- Soejarwo, P. A.; Yusuf, R. and Zulham, A.** (2019). Analysis of the Sustainability of Seaweed Cultivation Businesses in East Sumba, East Nusa Tenggara. *Sosek Journal. Center for Maritime and Fisheries Socioeconomic Research.* Vol 14. No 1.
<https://doi.org/10.15578/JSEKP.V14I1.7815>

Tetelepta; Johannes M.S.; Yuliana Natan; Jesaja A. Pattikawa, and Agil S. Bernardus.(2022). “Population Paraeters and Sustainable Status of Lompa Fish Thryssa Baelama (Forsskal, 1775) Manage through Sasi Approach at Haruku Village.” *Fisheries and Aquatic Sciences.*, 25(2): 101–16.

<https://doi.org/10.47853/FAS.2022.e10>

Younis, A. M.; Hanafy, S.; Raafa, M. and El-Nagggar, M. H. (2023). Evaluation of Pollution Status and Risk Assessment Along the Al-Shuaibah Coastline, the Red Sea, Saudi Arabia. *Egyptian Journal of Aquatic Biology and Fisheries.*, 27(6): 923–945. <https://doi.org/10.21608/ejabf.2023.332417>

Yusuf, I.; Ismail, A. andNuva. (2023). Evaluation of the Sustainability Status of Seaweed Cultivation in Pasikolaga District, Muna Regency (Rapfish Approach, Multi Dimensional Scaling). *Journal of Environmental Sciences.*, 21(1):150-158,

<https://doi.org/doi:10.14710/jil.21.1.150-158>