

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/345805912>

The Measurement of Cortisol Levels in the Blood Pregnant Trimester III with ELISA (Enzyme Linked Immunosorbent Assay) Method

Preprint · October 2020

DOI: 10.13140/RG.2.2.30692.40327

CITATIONS

0

READS

45

7 authors, including:



Mochammad Hatta

Universitas Hasanuddin

423 PUBLICATIONS 5,294 CITATIONS

[SEE PROFILE](#)



Alfina Baharuddin

Universitas Muslim Indonesia

35 PUBLICATIONS 2 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Breast Cancer Chemotherapy Response [View project](#)



BIOSCIENCES AND BIOTECHNOLOGY [View project](#)



Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfmt.com



Official Organ of Indian Association of Medico-Legal Experts (Regd.)

Indian Journal of Forensic Medicine & Toxicology

EDITOR in Chief

Prof. R K Sharma

Formerly at All India Institute of Medical Sciences, New Delhi, E-mail: editor.ijfmt@gmail.com

EDITOR

Prof. Dr. Adarsh Kumar

Forensic Medicine & Toxicology, AIIMS, New Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Prof Mete Gulmen** Cukurova University, TURKEY
2. **Prof. Leandro Duarte De Carvalho**, Minas Gerais, Belo Horizonte, Brazil
3. **Prof. Donata Favretto** (Full Professor) Forensic Toxicology at University of Padova, Italy
4. **Prof. Babak Mostafazadeh** Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
5. **Prof Halis Dokgoz**, Mersin University, TURKEY
6. **Prof Jozef Sidlo**, Comenius University, Bratislava, SLOVAKIA
7. **Dr. Rahul Pathak** (Lecturer) Forensic Science, Dept of Life Sciences Anglia Ruskin University, Cambridge, United Kingdom
8. **Dr. Hareesh** (Professor & Head) Forensic Medicine, Ayder Referral Hospital, College of Health Sciences, Mekelle University, Mekelle Ethiopia East Africa
9. **Dr. Mokhtar Ahmed Alhrani** (Specialist) Forensic Medicine & Clinical Toxicology, Director of Forensic Medicine Unit, Attorney General's Office, Sana'a, Yemen
10. **Dr. Sarathchandra** Kodikara (Senior Lecturer) Forensic Medicine, Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka
11. **Dr Noha A. Magdie El Rafie**, Forensic Toxicology, Ain Shams University, Cairo, EGYPT

SCIENTIFIC COMMITTEE

1. **Prof Udai Pratap Singh**, Department of Anthropology Lucknow University Lucknow
2. **Dr Anil Rahule** (Associate Professor) Dept of Anatomy, Govt Medical College Nagpur
3. **Dr Shankar Bakkanwar** (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnatakad
4. **Dr K. Ravikumar** Raksha Shakti University, Ahmedabad, Gujrat.
5. **Dr. Pragnesh Parmar** (Associate Professor) Forensic Medicine, Valsad, Gujrat
6. **Dr Vandana Mudda** (Awati) (Associate Prof) Dept of FMT, M.R. Medical College, Gulbarga, Karnataka,
7. **Dr. Asha Srivastava** (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
8. **Dr. Lav Kesharwani** (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad U.P,
9. **Dr. Anu Sharma** (Associate Prof) Dept of Anatomy, DMCH, Ludhiana (PB)
10. **Dr. Shalini Gupta** (Prof) Oral Pathology and Microbiology, Dental Sciences King George Medical University, Lucknow, UP
11. **Dr Rituja Sharma**, Associate Prof, Law Banasthali Vidyapeeth Jaipur

"Indian Journal of Forensic Medicine & Toxicology" is peer reviewed quarterly journal. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine and environment medicine. It has been assigned International standard serial No. p-0973-9122 and e- 0973-9130. The Journal has been assigned RNI No. DELENG/2008/21789. The journal is indexed with Index Copernicus (Poland) and is covered by EMBASE (Excerpta Medica Database). The journal is also abstracted in Chemical Abstracts (CAS) database (USA). The journal is also covered by EBSCO (USA) database. The Journal is now part of UGC, DST and CSIR Consortia. It is now official publication of Indian Association of Medico-Legal Experts (Regd.).

NATIONAL EDITORIAL ADVISORY BOARD

Chairman

Prof Sudhir K Gupta - Head, Department of Forensic Medicine
All India Institute of Medical Sciences, New Delhi

Members

1. **Prof. SK Dhattarwal**, Forensic Medicine, PGIMS, Rohtak, Haryana
2. **Prof. N K Aggrawal** Forensic Medicine, UCMS, Delhi
3. **Prof Ajay Ghangale** Forensic Medicine Dr DY Patil Medical College, Pune, Maharashtra
4. **Dr. Amar Jyoti** Patwory Professor, Forensic Medicine NEIGRIHMS, Shillong
5. **Dr S. Venkata Raghava** Professor, Forensic Medicine, Bangalore Medical College, Bengaluru
6. **Prof Praveen Arora**, Professor Department of Forensic Medicine & Toxicology, SAIMS, Indore
7. **Dr. Pankaj Datta** (Principal & Head) Department of Prosthodontics, Indraprastha Dental College & Hospital, Ghaziabad
8. **Dr. Mahindra Nagar** (Head) Department of Anatomy, UCMS & GTB Hospital, Delhi
9. **Dr. Virender Kumar Chhoker** Professor Forensic Medicine and Toxicology, Santosh Medical College, Ghaziabad, UP
10. **Dr. Dayanand G Gannur** (Professor) Department of Forensic Medicine & Toxicology, Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
11. **Dr. Alok Kumar** Professor Department of Forensic Medicine & Toxicology, UP Rural Institute of Medical Sciences and Research, Saifai, Etawah, U.P.

Print-ISSN:0973-9122 Electronic - ISSN: 0973-9130

Frequency: Quarterly, © All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Forensic Medicine & Toxicology. Indian Journal of Forensic Medicine & Toxicology does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Website: www.ijfmt.com

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida
- 201 301 (Uttar Pradesh)

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida
- 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida
- 201 301 (Uttar Pradesh)

112.	The Measurement of Cortisol Levels in the Blood Pregnant Trimester III with ELISA (Enzyme Linked Immunosorbent Assay) Method	671
	<i>Ruqaiyah, Nusratuddin Abdullah, Mochammad Hatta, Nasrudin A Mappeware, Ayatullah Harun , Fatmawati Amir, Alfina Baharuddin</i>	
113.	Contamination of Polynuclear Aromatic Hydrocarbons-(PNAH) in Sediments: Identification and Distribution in the River of Shatt Al-Arab-(ROSA).....	677
	<i>Wisam Abdul-Ameer Farid, Wasen Abdul-Ameer Ali, Aseel Nadum Al-Salman</i>	
114.	Serum Androgens in Men with Rheumatoid Arthritis	684
	<i>Abdulsatar J. Mathkhor . Abdalnasser H. Abdullah. Amer S. Khudhairi</i>	
115.	Disposition of Policy Implementers: Study on The Hospitals Payment System Change in The Indonesian National Health Insurance Era.....	689
	<i>Abu Khoiri, Widi Hidayat, Djazuly Chalidyanto, Fendy Suhariadi</i>	
116.	How to Teach Intramuscular Injection Through Virtual Learning Environment in Covid-19 Pandemic Time for Nursing Students	697
	<i>Adam Mahmoud Salameh Khraisat, Hermizi Hapidin, Nik Siti Hanifah Nik Ahmad, Mohammad Dzulsyafiq Mohammad Yusoff, Mohd Said Nurumal, Kasmah Wati Pardi</i>	
117.	A Virtual Learning Environment for Teaching Intramuscular Injection: A Focus Group Discussion Analysis.....	705
	<i>Adam Mahmoud Salameh Khraisat, Hermizi Hapidin, Nik Siti Hanifah Nik Ahmad, Mohammad Dzulsyafiq Mohammad Yusoff, Mohd Said Nurumal, Kasmah Wati Pardi</i>	
118.	Risk Factors of Type 2 Diabetes Mellitus in Hospital of Bengkulu City, Indonesia: Case Control Study	710
	<i>Agung Sutriyawan, Tenike G Miranda, Hairil Akbar, Riswan</i>	
119.	Validity and Reliability of the Questionnaire Assessing the Healthcare Delivery Status in Iran's Prisons	717
	<i>Ahmad Hajebrahimi, Khalil Alimohammadzadeh, Seyed Mojtaba Hosseini, Ali Maher, Mohammadkarim Bahadori</i>	
120.	Outcome of Anterior Cruciate Ligament Reconstruction using Femoral Intrafix for a Single Tunnel Double Bundle Anterior Cruciate Ligament Reconstruction	724
	<i>Ahmed Mahmoud Gad, Mohamed Kamal Ismail, Ramy Salah Morsy</i>	
121.	PON1 Q192R Gene Polymorphism and Pesticide Exposure Status of Rice Farmers, Suphan Buri, Thailand.....	731
	<i>Airada Sridon, Kampol Ruchiwit, Kowit Suwannahong, Wanida Pongstaporn, Yuttana Sudjaroen</i>	
122.	Effect of Oral Contraceptive Pills on Levels of Calcium and Vitamin D in women in Al-Najaf Province	739
	<i>Alaa Abd Kadim, Suha Hussein Mohammed</i>	
123.	Clinical study on the therapeutic effects of <i>Quercus infectoria</i> galls as oral powder in gingivitis and plaque patients	744
	<i>Ali Hasanain Alhamadani, Hiba Ahmed Saeed, Hussein Asli Khayoon</i>	

The Measurement of Cortisol Levels in the Blood Pregnant Trimester III with ELISA (Enzyme Linked Immunosorbent Assay) Method

Ruqaiyah^{1,2}, Nusratuddin Abdullah³, Mochammad Hatta⁴, Nasrudin A Mappeware⁵, Ayatullah Harun², Fatmawati Amir², Alfina Baharuddin⁶

¹ Research in Postgraduate Program, Medical Faculty, Hasanuddin University, Indonesia, ²Lecturer in Akbid Pelamonia Kesdam VII/Wirabuana, Indonesia, ³ Professor Obygn in, Medical Faculty, Hasanuddin University, Indonesia, ⁴Professor Microbiology Department, of Medical Faculty, Hasanuddin University, Indonesia, ⁵Assistant Professor Obygn in Medical Faculty, University Moslem of Indonesia, ⁶Research Environmental health Departement of public health, University Moslem Of Indonesia

Abstract

Background: During pregnancy, maternal cortisol increases 2-4 times. Every type of body response in the form of stress, both physical and psychological stress can increase ACTH secretion, which in turn can increase cortisol levels.

Methods: This study uses quasi experimental / quasi experimental one group design pre and post test design with group control. The sample in this study was 24 respondents. . Cortisol level examination tool uses a microplate reader with the ELISA examination method.

Results: There was a significant difference ($p = 0.000 < \alpha$) of the average cortisol level in the control group before observation (54.43 ± 25.86) and after observation (459.15 ± 28.08) Whereas in the treatment group showed a significant difference ($p = 0.004 < \alpha$) the average cortisol level before being treated (65.26 ± 28.74) and after being treated (53.65 ± 23.68) Yoga exercises.

Conclusion: There was a significant difference ($p = 0,000 < \alpha$), mean cortisol levels in the control group before observation (54.43 ± 25.86) and observations after (459.15 ± 28.08), whereas in the treatment group there were significant differences ($p = 0.004 < \alpha$) it can be concluded that prenatal yoga reduces cortisol levels before and after treatment.

Keywords: Cortisol, blood level, ELISA, pregnant, Trimester III

Introduction

Cortisol is the main product of the hypothalamus-pituitary-adrenal (HPA) axis, secreted both in response to threats and in a strong and consistent pattern throughout the day (diurnal cortisol rhythm). Cortisol has many effects on the body, including building up energy and shutting down systems that are not essential for dealing with threats, and also affect memory and mood^{1,2}.

Under normal circumstances, stress hormones are released in small amounts throughout the day, but when faced with stress these hormone levels increase dramatically (in the range of 5-20 times). Each type of body response in the form of stress, both physical stress and psychological stress can increase secretion ACTH, which can ultimately increase cortisol levels. In conditions of anxiety, anxiety and depression, cortisol secretion increased^{3,4,5} Increased cortisol affects the function of almost every cell and tissue in the body by activating glucocorticoid receptors (GR) and mineralocorticoid receptors (MR) that modify gene expression programs through direct binding to specific DNA sequences and

Corresponding author:

Alfina Baharuddin

alfina.baharuddin@umi.ac.id

recruitment of transcription complexes^{6,7}.

Several clinical studies have examined stressors that occur naturally and most report insignificant or low relationships. This is especially true for studies that use a single measure of cortisol from amniotic fluid, serum or saliva. Ecological momentary assessments are used to simultaneously assess mood and cortisol 5 times a day for 3 days in 83 women (6-37 weeks gestational age). possible cortisol as a biological relationship between maternal psychological stress during pregnancy and fetal development^{8,9}.

The results from multilevel analysis showed a strong relationship between negative mood and cortisol. For every 1.0% increase in negative moods there is an appropriate 1.9% increase in cortisol. This association was not affected by advanced gestational age cross-sectional studies at the end of pregnancy in 79 women to investigate the relationship between the factors involved in anxiety and depression^{10,11,12}.

Some regression analysis shows that depressed subjective feelings explain more than 50% of the variation of BDI and STAI scores. Plasma cortisol is not a significant predictor of psychometric scores and did not show a significant correlation with them in correlation analysis, and subjects with low and high cortisol levels showed the same psychometric scores. The mechanism that mediates the effects of subjective pressure remains unclear: candidates who might include monoamine neurotransmission and / or changes caused by stress in the expression or distribution of glucocorticoid receptors^{13,14}..

Research to test one potential mechanism by examining the relationship between pregnancy anxiety and maternal salivary cortisol on 4 occasions during pregnancy in a sample of 448 women. Strong evidence from animal models shows that maternal pressure predicts increased fetal glucocorticoid exposure through synthesis and release of maternal glucocorticoids but similar evidence in humans that links pregnancy anxiety (or general maternal pressure) with increased levels of maternal cortisol during pregnancy is still lacking^{5,9}

During pregnancy, maternal cortisol increases 2-4-fold, and crosses the placenta but is limited because the pathway is regulated by the enzyme 11 β -hydroxysteroid

dehydrogenase (11 β -HSD2: Maternal cortisol accounts for about 30-40% of the variability in fetal cortisol concentration. Physiological increase in glucocorticoid levels during the pregnancy, facilitating fetal organ maturation by promoting cell differentiation Synthetic glucocorticoids (including dexamethasone or betamethasone) are given during the third trimester of pregnancy at risk of preterm delivery to increase the survival rate and maturation of the lungs^{15,16}. Based on these references, researchers are interested in examining cortisol levels in pregnant women with prenatal yoga treatment.

Materials ad Methods

This study used quasi experimental one group design pre and post test design with group control. In this study the researchers' treatment or intervention was prenatal Yoga exercises in third trimester pregnant women (UK 28-40 weeks) ,The sample in this study was 24 respondents.

Cortisol examination procedure;

a) Prepare samples, reagents, main solutions and equipment

b) Let the blood sample freeze for 2 hours at 4 ° C. Do it

c) Centrifugation for 15 minutes, then immediately assay checks or can be divided and stored at temperatures <-20 ° C ($\leq$ 1 month), or -80 ° C ($\leq$ 6 months). Blood tubes must be disposable, sterile.

d) Reagents are prepared at room temperature

e) Dilute 30 ml of WBC to 750 ml of wash buffer with distilled water.

f) If crystalline deposits are found in the Wash Buffer Concentrate, warm at 40 ° C and shake gently until the crystals dissolve.

g) The solution must be cooled to room temperature before use

h) Standard Solution

1. Prepare a standard solution 15 minutes before use.

2. Standard with 1 ml Sample Diluent, allow 10 minutes to dissolve.

3. This reconstruction produces 40 ng / ml stock solution. Then make serial dilutions (40, 20, 10, 5, 2.5, 1.25, 0.63, 0 ng / mL).

i) Procedure:

1. Add 50 µl of standard or sample to each microplate hole.

2. Add 50 µl Biotinylated Detection AB. Incubation 45 minutes at 37 ° C

3. Aspiration and wash 3 times

4. Add 100 µl HRP Conjugate. Incubation 30 minutes at 37 ° C

5. Aspiration and wash 5 times

6. Add 90 µl Substrate Reagent. Incubation for 15 minutes at 37 ° C

7. Add 50 µl Stop Solution. Read at 450 nm right away.

8. Calculation of results in the form of OD values. Real values or concentrations are generated by entering OD values in standard curves.

Result

Table 1 Characteristics of education level and type of work

Variable	Group		p-value
	Control (%)	Treatment (%)	
Level education			0.590
- SMA	1 (8.3%)	3 (25%)	
- D3	11 (91.7%)	9 (75%)	
Type work			1.000
- House wife	10 (83.3%)	11 (91.7%)	
- Wiraswasta	2 (16.7%)	1 (8.3%)	

The results level education test analysis on the distribution of data on the education level of pregnant women in Table 3 showed no significant difference ($p = 0.590 > \alpha$) which was divided into 2 levels of education. Namely: high school and D3. Whereas there was no significant difference in the type of work ($p = 1.00 > \alpha$)

Table 2: The Comparison Of Cortisol Levels Before And After Treatment

Group observation	Before Rerata $\pm$ SD	After Rerata $\pm$ SD	p-value
Control	54.43 $\pm$ 25.86	459.15 $\pm$ 28.08	0.000
Treatment Yoga	65.26 $\pm$ 28.74	53.65 $\pm$ 23.68	0.004

Table 2 shows that there is a significant difference ($p = 0,000 < \alpha$) of the average cortisol level in the control group before observation (54.43 ± 25.86) and after observation (459.15 ± 28.08). it means Whereas in the

treatment group showed a significant difference ($p = 0.004 < \alpha$) the average cortisol level before being treated (65.26 ± 28.74) and after being treated (53.65 ± 23.68) Yoga exercises

Table 3 The Correlation Test Results Between Variables In The Group

Treatment After Yoga Exercises

Correlation of variable	n	Koefisien korelasi (r)	p-value
Level anxiety and kortisol	12	0.853	0.000
Cortisol With FKBP5	12	0.937	0.000

Discussion

In the results of this study the treatment of Yoga exercises in third trimester pregnant women can significantly reduce cortisol levels. So the third minor hypothesis has been proven, that is, there is a difference in effect between non-action and prenatal Yoga exercises on cortisol levels in primigravida trimester III.

In line with previous studies where anxiety disorders were associated with increased levels of basal cortisol in saliva as well as in plasma. Stress conditions during pregnancy will affect hormonal changes and fetal growth, especially in the hormone cortisol and will result in impaired fetal growth such as low birth weight and abnormal body length. Several studies have shown that depression is associated with hyperactivity from the hypothalamic-pituitary-adrenal (HPA) axis which results in high cortisol levels^{17,18,19}.

During pregnancy, maternal cortisol increases 2-4 times. The physiological increase in glucocorticoid levels during pregnancy, facilitates maturation of the fetal organs by promoting cell differentiation. Synthetic glucocorticoids (including dexamethasone or betamethasone) given during the third trimester of pregnancy are at risk of preterm delivery to increase the survival rate and maturation of the lungs^{8,110,114}.

Research conducted on white rats (*Rattus Norvegicus*) that formation of cortisol and cortisol secretion increased six times in a mouse within 4 to

20 minutes after fracture of the two bones in his legs which caused the rat to be stressed. This is in line with the literature that stress can cause an increase in ACTH secretion by anterior pituitary gland followed by an increase in cortisol hormone secretion within a few minutes, prenatal yoga reduces anxiety and decreases cortisol levels.²⁰

Stressors activate the hypothalamic-pituitary-adrenal (HPA) axis, which triggers the secretion of glucocorticoid hormones (cortisol for humans and corticosterone for mice) to drive systemic changes in physiological patterns, which aim to return the body to homeostasis. Exposure to high levels of glucocorticoids can cause changes in fetal programming, as well as affect the development of glucocorticoid receptors in the fetal brain. For mothers, glucocorticoids induce physiological and behavioral changes that can interfere with hereditary care^{4,9,13}.

Initially, stress causes an increase in diurnal cortisol, facilitating the recruitment of energy and coping resources. However, repeated or sustained increases in cortisol can cause damage to the negative feedback system of cortisol secretion, resulting in flat slopes and negative clinical and health outcomes^{13,20}. Heterogeneity in the relationship of chronic stress to diurnal slope indicates that some individuals are more susceptible than others to a disturbance in the healthy relationship between stress and cortisol secretion. One factor that might be a genetic risk.^{21,22}

When cortisol levels begin to fall, the inhibitory effect of cortisol on the hypothalamus and anterior pituitary decreases so that the factors that stimulate increased cortisol secretion (CRH-ACTH) will increase⁹. This system is sensitive because cortisol production or excessive administration of cortisol or other synthetic glucocorticoids can quickly inhibit the hypothalamic-pituitary axis and stop the production of ACTH^{23,24}. This study contradicts previous results, in which researchers investigated whether anxiety disorders were associated with the HPA axis by using a new method of assessing cortisol in hair for a method that was considered to provide a retrospective cortisol index of secretion over a long period of time. The results showed lower hair cortisol levels in anxiety disorder patients than in healthy controls. This interpretation tentatively indicates the condition of anxiety disorders is associated with hypocortisolism^{3,25}.

The hypoactive HPA axis model has been used to explain adrenocortical activity in other psychiatric and psychosomatic disorders as well as in stress illustrating that participants who exhibit hypocortisolism often exhibit a series of symptoms characterized by increased sensitivity to stress, pain and fatigue²⁶, which is characterized by increased sensitivity to stress, pain and fatigue. The pathway that leads to hypocortisolism is likely to result from exacerbated worries, higher tension and anxiety may lead to an initial increase in adrenocortical activity, namely hypercortisolism. With the increase in the mechanism of chronic compensation becomes activated and gradually produces a weakening of cortisol secretion, namely hypocortisolism.^{16,20}

Conclusion

There was a significant difference ($p = 0.000 < \alpha$), mean cortisol levels in the control group before observation (54.43 ± 25.86) and after observations (459.15 ± 28.08), whereas in the treatment group there were significant differences ($p = 0.004 < \alpha$) cortisol levels before being treated (65.26 ± 28.74) and after being treated (53.65 ± 23.68), it can be concluded that prenatal yoga reduces cortisol levels before and after treatment.

Source of Funding - Self-funding

Conflict of Interest- None of the authors has

competing interests

Ethical Clearance- This research was approved by the Research Ethics Commission of the Faculty of Medicine, Hasanuddin University Makassar, (No. 839/UN4.6.4.5.31/PP36/2019), and all research subjects gave written informed consent.

References

1. McDonald, S.W., Lyon, A.W., Benzie, K.M., McNeil, D.A., Lye, S.J., Dolan, S.M., Pennell, C.E., Bocking, A.D., Tough, S.C., The All Our Babies pregnancy cohort: design, methods, and participant characteristics. *BMC Pregnancy Childbirth* 13 (Suppl2013. 1), S2.
2. Neylan, T.C., Brunet, A., Pole, N., Best, S.R., Metzler, T.J., Yehuda, R., Marmar, C.R., PTSD symptoms predict waking salivary cortisol levels in police officers. *Psychoneuroendocrinology* 2005, 30, 373–381.
3. McEwen BS, Stellar E. Stress and the individual. Mechanisms leading to disease. *Arch Intern Med.* 1993;153(18):2093–2101.
4. D. Picard, B. Khursheed, M.J. Garabedian, M.G. Fortin, S. Lindquist, K.R. Yamamoto, Reduced levels of hsp90 compromise steroid receptor action in vivo, *Nature* 348 (6297) (1990) 166–168.
5. Dahlan, M. Sopiudin. Besar Sampel dan Pengambilan Sampel, dalam Penelitian Kedokteran dan Kesehatan. Jakarta: Salemba Medika..2009
6. Daniel A. Ostrovsky, MD, FACP, FAAP. Yoga in the third trimester may Reduce labor pain, duration of labor, and risk Morris, M.C., Compas, B.E., Garber, J., 2012. Relations among posttraumatic stress disorder, comorbid major depression, and HPA function: a systematic review and metaanalysis. *Clin. Psychol.* 2013, Rev. 32, 301–315.
7. Ngai FW, Chan SW. Psychosocial factors and maternal wellbeing: an exploratory path analysis. *Int J Nurs Stud.* 2011;48(6):725–731, <http://dx.doi.org/10.1016/j.ijnurstu.2010.11.002>
8. Nicolas C Nicolaidis and George P Chrousos. Corticotropin-Releasing Hormone (CRH). Nomenclature. *Encyclopedia of Endocrine Diseases*, Second Edition, 2019, Volume 2 doi:10.1016/B978-0-12-801238-3.64324-6nih.gov/pubmed/25287545

[Accessed November 22, 2016].

9. Nirwana, A. B. *Psikologi Kesehatan Wanita*. Yogyakarta : Muha Medika, 2011
10. Liu, R.Y., Unmehopa, U.A., Zhou, J.N., Swaab, D.F., Glucocorticoids suppress vasopressin gene expression in human suprachiasmatic nucleus. *J. Steroid Biochem. Mol. Biol.* 98, 248–253.
11. Pao-Ju Chena,b, Luke Yangc, Cheng-Chen Choud, Chia-Chi Li e, Yu-Cune Changf, Jen-Jiuan Liawe,*; Effects of prenatal yoga on women's stress and immune function across pregnancy-A randomized controlled trial. *Complementary Therapies in Medicine* 206, 31 (2017) 109–117
12. Paquette, A.G., Lester, B.M., Koestler, D.C., Lesseur, C., Armstrong, D.A., Marsit, C.J., Placental FKBP5 genetic and epigenetic variation is associated with infant neurobehavioral outcomes in the RICHS cohort. 2014, *PLoS One* 9 (8), e104913.
13. Pariante, C.M., 2006. The glucocorticoid receptor: part of the solution or part of the problem? *J. Psychopharmacol.* 20, 79–84.
14. Liu, Z., Zhu, F., Wang, G., Xiao, Z., Wang, H., Tang, J., Wang, X., Qiu, D., Liu, W., Cao, Z., Li, W., 2006b. Association of corticotropin-releasing hormone receptor1 gene SNP and haplotype with major depression. *Neurosci. Lett.* 404, 358–362
15. McEwen BS dan Wingfield JC. The concept of allostatis in biology and biomedicine. *Hormones and behavior* 2003; 43:2-15
16. Willour, V.L., Chen, H., Toolan, J., Belmonte, P., Cutler, D.J., Goes, F.S., Zandi, P.P., Lee, R.S., MacKinnon, D.F., Mondimore, F.M., Schweizer, B., DePaulo Jr., J.R., Gershon, E.S., McMahon, F.J., Potash, J.B., Family-based association of FKBP5 in bipolar disorder. *Mol. Psychiatry* 2009, 14, 261—268
17. Wochnik GM, Ruegg J, Abel GA, Schmidt U, Holsboer F, Rein T. FK506-binding proteins 51 and 52 differentially regulate dynein interaction and nuclear translocation of the glucocorticoid receptor in mammalian cells. *The Journal of Biological Chemistry.* 2005; 280:4609–16. [PubMed: 15591061]
18. Wochnik, G.M., Ruegg, J., Abel, G.A., Schmidt, U., Holsboer, F., Rein, T., FK506- binding proteins 51 and 52 differentially regulate dynein interaction and nuclear translocation of the glucocorticoid receptor in mammalian cells. *J. Biol. Chem.* 2005, 280, 4609–4616. <https://doi.org/10.1074/jbc.M407498200>
19. Abelson, J.L., Khan, S., Liberzon, I., Young, E.A., HPA axisactivity in patients with panic disorder: review and synthesis offour studies. *Depress. 207, Anxiety* 24 (1), 66—76.
20. Aguilera G. Regulation of Pituitary ACTH Secretion During Chronic Stress *Frontiers in Neuroendocrinology.* 1994;15:321e350.
21. Dahro, Ahmad. *Psikologi Kebidanan : Analisis Perilaku Wanita Untuk Kesehatan*. Jakarta : Salemba Medika, 2012.
22. Dallman MF, Akana SF, Strack AM, Scribner KS, Pecoraro N, La Fleur SE, Houshyar H, Gomez F. Chronic stress-induced effects of corticosterone on brain: direct and indirect. *Ann NY Acad Sci.* 2004;1018:141e150Aguilera,G., Nikodemova,M., Wynn,P.C., Catt,K.J., Corticotropin releasing hormone receptors: two decades later. *Peptides* 2004, 25, 319–329. <http://dx.doi.org/10.1016/j.peptides.2004.02.002>
23. Wolf, E.J., Mitchell, K.S., Logue, M.W., Baldwin, C.T., Reardon, A.F., Humphries, D.E., Miller, M.W., Corticotropin releasing hormone receptor 2 (CRHR-2) gene is associated with decreased risk and severity of posttraumatic stress disorder in women. *Depress. Anxiety* 2013, 30, 1161–1169.
24. 49. Holsboer, F., Stress, hypercortisolism and corticosteroid receptors in depression: implications for therapy. *J. Affect. Disord.* 2001, 62, 77–91.
25. McEwen BS danGianaros PJ. Stress and allostasis-induced brain plasticity. *Annu. Rev. Med.* 2011;62:431-45
26. Mulder EJ, Robles de Medina PG, Huizink AC, Van den Bergh BR, Buitelaar JK, Visser GH. Prenatal maternal stress: effects on pregnancy and the unborn child. *Early Hum Dev* 2002;70:3e14.